S. I. LIBRARY

Cat. cde 5/28/43

QL
577
B82
pt.9
Ent.

CATALOGUE

OF

COLEOPTEROUS INSECTS

IN

THE COLLECTION

OF THE

BRITISH MUSEUM.

PART IX.

CASSIDIDÆ.

BY

CHARLES H. BOHEMAN,

PROFESSOR OF NATURAL HISTORY, STOCKHOLM.

SMITHSONIAN INSTITUTION 1861

LONDON:

PRINTED BY ORDER OF THE TRUSTEES.

1856.

S. I. 1913
QIM.S.

S. BAGSTER AND SONS, TYP.,
LONDON.

SMITHSONIAN
LIBRARY

595.798
B676
Cent.

PREFACE.

THE object of the present Catalogue is to give a complete list of all the genera and species of Tortoise Beetles (*Cassididæ*) which have been described in various works, and the descriptions of those, not previously noticed in the Author's "Monographia Cassididarum, Holmiæ, 1850–1855," have been added.

The letters B.M. in the margin show the species now contained in the British Museum, and their absence indicates those which are desiderata, and therefore desirable to be procured for the Collection.

JOHN EDWARD GRAY.

Nov. 1856.

CATALOGUE

OF

CASSIDIDÆ.

TRIBUS I.

Genus 1. DELOCRANIA.

Dejean, Cat. ed. 3, p. 391. *Guérin.* Mag. de Zool. 1844, Ins. pl. 131. *Bohem.* Mon. Cassid. i. p. 4.

1. Delocrania cossyphoides. B.M.

Guér. Mag. de Zool. 1844, Ins. pl. 131. *Bohem.* Mon. Cassid. i. p. 5, 1, tab. I. A.

Delocrania hispoides, *Dej.* Cat. ed. 3, p. 367.
Patria. Brasilia. (Tapayos. Coll. *W. B. Bates.*)

Genus 2. SPHÆROPALPUS.

Chevrolat. Dej. Cat. ed. 3, p. 391. *Guér.* Icon. de Règne Anim. ii. p. 485. *Bohem.* Mon. Cassid. i. p. 6.

Imatidium, *Dej.* Cat. ed. 2, p. 115.
Platyauchenia, *Sturm.* Cat. ed. 1843, t. vi. f. 8.

1. Sphæropalpus cinctus. B.M.

Dej. Cat. ed. 3, p. 391. *Guér.* Icon. de Règne Anim. Ins. texte, p. 285. *Bohem.* Mon. Cassid. i. p. 7, 1. tab. I. B.

Platyauchenia limbata, *Sturm.* Cat. 1843, p. 359, t. vi. f. 8.
Var. a. sutura elytrorum, præsertim antice, latius flava.
Patr. Brasilia.

Genus 3. CALOPEPLA.

Hope, Col. Man. iii. p. 152. *Bohem.* Mon. Cassid. i. p. 8.
Imatidium, *Schœnh.* Syn. Ins. ii. p. 134. *Dej.* Cat. ed. 2, p. 115.
Cassida, *Oliv.* Ent. vi. p. 922.
Craspedonta, *Chevrol. Dej.* Cat. ed. 3, p. 367.

1. Calopepla Leayana. B.M.

Bohem. Mon. Cassid. i. p. 9, 1. tab. I. C.
Imatidium Leayanum, *Latr.* Gen. Crust. et Ins. iii. p. 50, 1. t. ii. f. 7. *Schœnh.* Syn. Ins. ii. p. 135, 6. *Dej.* Cat. ed. 2, p. 115.
Cassida Leyana, *Oliv.* Ent. vi. p. 951, 45, 97, t. v. f. 74.
Craspedonta Leyana, *Dej.* Cat. ed. 3, p. 367.
Patr. Bengalia et Assam Indiæ orientalis.

2. Calopepla Reicheana.

Guér. Icon. de Règne Anim. Ins. texte ii. p. 286. *Bohem.* Mon. Cassid. i. p. 11, 2.
Patr. India orientalis.

Genus 4. EPISTICTIA.

Bohem. Mon. Cassid. i. p. 12.

1. Epistictia selecta. B.M.

Bohem. Mon. Cassid. i. p. 13, 1. tab. I. D.
Patr. India orientalis (Bombay Coll. *J. S. Wooley*), Tasmania.?

2. Epistictia Matronula.

Bohem. Mon. Cassid. i. p. 14, 2.
Patr. Ceylon.

3. Epistictia viridimaculata.

Bohem. Mon. Cassid. i. p. 15, 3.
Patr. Nepal Indiæ orientalis.

Genus 5. HOPLIONOTA.

Hope, Coleopt. Man. iii. p. 153. *Guér.* Icon. de Règne Anim Ins. texte ii., p. 287. *Bohem.* Mon. Cassid. i. p. 16.
Cassida, *Fabr.* Syst. El. i. p. 387. *Oliv.* Ent. vi. p. 922. *Herbst* Naturs. viii. p. 204. *Schœnh.* Syn. Ins. ii. p. 209.

Cassida (Imatidium) *Klug*, Ins. v. Madag. p. 122.
Notosacantha, *Chevrol. Dej.* Cat. ed. 3, p. 391.
Thyreomorpha, *Dej.* Cat. ed. 3, p. 367.

I. Elytra sub-quadrata.

A. *Elytra pone medium late rotundato-emarginata, singulum unispinosum.*

1. Hoplionota echinata. B.M.

Bohem. Mon. Cassid. i. p. 18, 1. tab. I. E. 1.
Cassida echinata, *Fabr.* Syst. El. i. p. 390, 12. *Oliv.* Ent. vi. p. 963. 68. 97. t. v. f. 86. *Schœnh.* Syn. Ins. ii. p. 212, 12. *Boisd.* Faune Ent. de l'Océan. ii. p. 542, 1.
Notosacantha echinata, *Dej.* Cat. ed. 3, p. 391.
Patr. Java, Australasia, Madagascar. *J. S. Baly*, Esq.

B. *Elytra lateribus non emarginata, singulum unispinosum.*

2. Hoplionota armigera.

Bohem. Mon. Cassid. i. p. 19, 2.
Cassida armigera, *Oliv.* Ent. vi. p. 962, 67, 97, t. v. f. 85.
Patr. Madagascar.

3. Hoplionota semiviridis.

Guér. Icon. de Règne Anim. Ins. texte ii. p. 287. *Bohem.* Mon. Cassid. i. p. 20, 3.
Patr. Madagascar.

4. Hoplionota pallescens.

Guér. Icon. de Règne Anim. Ins. texte ii. p. 287. *Bohem.* Mon. Cassid. i. p. 21, 4.
Patr. Madagascar.

5. Hoplionota bispinosa.

Bohem. Mon. Cassid. i. p. 21, 5.
Patr. Madagascar.

6. Hoplionota biaculeata.

Sub-quadrata, modice convexa, subopaca; prothorace dorso remote, lateribus crebrius, profundius punctato; elytris crebre, sat profunde, minus regulariter punctato-striatis, dorso viridibus, singulo antice bi-carinato, spina sat longa, acuta instructo, pone illam macula sat magna, testacea, postice carinula transversa, arcuata, extus cum spina connexa. Long. 5, lat. 3¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Madagascar, Mus. Dom. *Deyrolle.*

C. *Elytra lateribus non emarginata, singulum bispinosum vel tuberculatum.*

7. Hoplionota quadrituberculata.

Guér. Icon. de Règne Anim. Ins. texte ii. p. 287. *Bohem.* Mon. Cassid. i. p. 23, 6.
Patr. Madagascar.

8. Hoplionota quadricornis.

Bohem. Mon. Cassid. i. p. 24, 7.
Patr. Madagascar.

9. Hoplionota viriditincta.

Sub-quadrata, modice convexa, testacea, parum nitida; prothorace dorso obsolete, lateribus profunde, sub-remote punctato; elytris minus profunde, irregulariter punctato-striatis, dorso antico parce viridi-signatis, singulo vitta intramarginali fusca notato, bituberculato, tuberculo antico parvo, postico valido, breviter conico, intra apicem transversim irregulariter carinato. Long. 5, lat. 4, millim.
Bohem. Mon. Cassid. Append.
Patr. Madagascar, Mus. Dom. *Deyrolle.*

D. *Elytra lateribus non emarginata, singulum tri-vel quadrispinosum aut tuberculatum.*

10. Hoplionota viridinotata.

Sub-quadrata, modice convexa, nitida, testacea; prothorace dorso nigro-æneo, mediocriter punctato, utrinque oblique impresso, lateribus profunde punctato; scutello ferrugineo; elytris crebre, profunde punctato-striatis, castaneis, singulo macula sat magna basali, viridi, antice tri-carinato, medio tuberculo conico et postice rugula transversa instructo. Long. 4½, lat. 3½ millim.
Bohem. Mon. Cassid. Append.
Patr. Madagascar, Mus. Dom. *Deyrolle.*

11. Hoplionota horrida.

Bohem. Mon. Cassid. i. p. 25, 8.
Patr. Madagascar.

12. Hoplionota paradoxa.

Bohem. Mon. Cassid. i. p. 26, 9.
Patr. Madagascar.

13. HOPLIONOTA RUBROMARGINATA. B.M.

Sub-quadrata, modice convexa, nigro-picea, nitida, undique dilute ferrugineo-marginata, subtus rufo-testacea; prothorace dorso lævi, lateribus profunde, minus crebre punctato; elytris sat profunde, crebre sub-seriatim punctatis, singulo tuberculis anticis tribus parvis et pone medium ruga elevata cruciformi, postice carinula parva, arcuata adjecta; margine profundius, remotius, extrorsum sub-seriatim punctato. Long. 5, lat. 4½ millim.

Bohem. Mon. Cassid. Append.
Patr. Ceylon, Mus. Brit. et Mus. Dom. *Baly.*

14. HOPLIONOTA MACULIPENNIS.

Sub-quadrata, modice convexa, testacea, parum nitida; pectore piceo; prothorace dorso transversim bi-impresso, remote, subtilius, lateribus profunde, crebre punctato, ante basin nigro-bimaculato; elytris mediocriter, minus regulariter punctato-striatis, singulo tuberculis inæqualibus octo nigris instructo; margine sat late explanato, maculis duabus obsoletis, fuscis, una ante, altera pone medium ornato. Long. 5, lat. 4 millim.

Bohem. Mon. Cassid. Append.
Var. a prothorace immaculato.
Patr. India orientalis, Dom. *Westwood,* Mus. Holm.

15. HOPLIONOTA HORRIFICA. B.M.

Sub-quadrata, modice convexa, testacea, parum nitida; prothorace dorso transversim bi-impresso, subtilius, lateribus profunde, crebre punctato; elytris mediocriter, minus regulariter punctato-striatis, parce nigro-variegatis, singulo carinis tuberculisque inæqualibus instructo, tuberculo pone medium reliquis multo validiore; margine sat late explanato, transversim rugoso, ante medium prope discum macula parva, albida, sub-hyalina. Long. 5, lat. 4, millim.

Bohem. Mon. Cassid. Append.
Var. a. supra castanea.
Patr. In Insulis Philippensibus, Mus. Brit.—Ceylon, Mus. Dom. *Dohrn.*—China et Manilla, Mus. Reg. Holm.—*Var. a.* India orient. Mus. Dom. *Saunders.*

16. HOPLIONOTA PLICATA.

Bohem. Mon. Cassid. i. p. 27, 10.
Cassida (Imatidium) plicata, *Klug.* Ins. v. Madag. p. 122, 199. t. v. f. 10. Acta Berol. 1832, p. 210, 199, t. v. f. 10.
Patr. Madagascar.

17. Hoplionota pulchella.

Guér. Icon. de Règne Anim. Ins. texte ii. p. 279. *Bohem.* Mon. Cassid. i. p. 28, 11.
Patr. Madagascar.

18. Hoplionota ochracea.

Bohem. Mon. Cassid. i. p. 29, 12.
Patr. Madagascar.

19. Hoplionota luctuosa.

Bohem. Mon. Cassid. i. p. 30, 13.
Patr. Madagascar.

20. Hoplionota odiosa.

Sub-quadrata, modice convexa, supra cærulescenti-nigra, parum nitida, subtus cum capite testacea; prothorace dorso obsolete, lateribus profunde punctato, antice anguste testaceo-marginato; elytris sat crebre, mediocriter punctato-striatis, singulo seriebus duabus e tuberculis parvis instructo. Long. 4¾, lat. 4¼ millim.

Bohem. Mon. Cassid. Append.
Patr. Java, Mus. Soc. Ind. orient.

21. Hoplionota erythrocata.

Bohem. Mon. Cassid. i. p. 31, 14.
Patr. Madagascar.

E. *Elytra lateribus non emarginata, dorso carinata.*

22. Hoplionota pallens.

Bohem. Mon. Cassid. i. p. 32, 15.
Patr. Madagascar.

23. Hoplionota marginata.

Bohem. Mon. Cassid. i. p. 33, 16.
Patr. Madagascar.

II. Elytra subrotundata.

24. Hoplionota Molucana.

Hombron et *Jaquinot*, Voy. d'Urville, Ins. pl. 18, f. 8. *Bohem.* Mon. Cassid. i. p. 34, 17.
Patr. Amboina.

25. Hoplionota badia.

Bohem. Mon. Cassid. i. p. 35, 18.

Thyreomorpha badia, *Dej.* Cat. ed. 3, p. 391.
Patr. Caput bonæ Spei.

26. HOPLIONOTA INCOMTA. B.M.

Sub-rotundata, paullo convexa, testacea, sub-opaca; antennis articulis quinque ultimis nigris; prothorace dorso obscuriore ibi vix, lateribus vero profunde, sub-seriatim punctato; elytris mediocriter, crebre, sub-seriatim punctatis; singulo carina longitudinalis flexuosa, fusca, ramulos duos transversos ad futuram emittente; margine basi macula oblonga intramarginali pallida, sub-hyalina. Long. 3½, lat. 3 millim.
Bohem. Mon. Cassid. Append.
Patr. Caffraria, Mus. Holm.—Brit. Mus. (Port Natal).

III. Elytra oblongo-quadrata.

27. HOPLIONOTA PARALLELA. B.M.

Bohem. Mon. Cassid. i. p. 36, 19, tab. I. E. 6.
Patr. Caffraria (Port Natal).

28. HOPLIONOTA SERRICOSTA.

Bohem. Mon. Cassid. i. p. 37, 20.
Patr. Java.

29. HOPLIONOTA OCHROLEUCA.

Bohem. Mon. Cassid. i. p. 38, 21.
Patr. India orientalis.

30. HOPLIONOTA GEMMATA.

Bohem. Mon. Cassid. i. p. 39, 22.
Cassida gemmata, *Klug*, Ins. v. Madag. p. 122, 200, t. v. f. 11. Acta Berol. 1832, p. 210, 200, t. v. f. 11.
Patr. Madagascar.

31. HOPLIONOTA TÆNIATA.

Bohem. Mon. Cassid. i. p. 40, 23.
Cassida tæniata, *Fabr.* Syst. El. i. p. 396, 49. *Oliv.* Ent. vi. p. 972, 86, 97, t. vi. f. 102. *Schœnh.* Syn. Ins. ii. p. 220, 69. *Boisd.* Faune Ent. de l'Océan. ii. p. 543, 4.
Patr. Insulæ Maris Pacifici.

Genus 6. MEGAPYGA.

Bohem. Mon. Cassid. i. p. 40.

1. MEGAPYGA CÆRULEOMACULATA. B.M.

Bohem. Mon. Cassid. i. p. 42, 1.
Patr. Manilla (Ins. Philipp.) Mus. Brit. Coll. *Cuming.*

2. MEGAPYGA EXIMIA. B.M.

Bohem. Mon. Cassid. i. p. 43, 2, tab. I. F.
Patr. Manilla (Ins. Philipp.) Mus. Brit. Coll. *Cuming.*

Genus 7. PRIOPTERA.

Hope, Col. Man. iii. p. 176. *Bohem.* Mon. Cassid. i. p. 44.
Cassida, *Fabr.* Syst. El. i. p. 387. *Oliv.* Ent. vi. p. 922. *Herbst.* Naturs. viii. p. 204. *Schœnh.* Syn. Ins. ii. p. 209.
Basiprionota, *Chevrol. Dej.* Cat. ed. 3, p. 367.

I. Elytra antice ad suturam gibbosa.

A. *Prothorax immaculatus.*

1. PRIOPTERA WESTERMANNI. B.M.

Mannerh. Bull. de Moscow, xvii. p. 864. *Bohem.* Mon. Cassid. i. p. 45, 1.
Patr. Assam Indiæ orientalis.

2. PRIOPTERA IMPUSTULATA.

Bohem. Mon. Cassid. i. p. 46, 2, t. ii. A.
Patr. Assam.

3. PRIOPTERA QUADRI-IMPRESSA.

Bohem. Mon. Cassid. i. p. 47, 3.
Basiprionota quadri-impressa, *Buquet, Dej.* Cat. ed. 3, p. 367.
Patr. Java.

4. PRIOPTERA SEXMACULATA. B.M.

Bohem. Mon. Cassid. i. p. 49, 4.
Patr. Assam.

5. PRIOPTERA MACULIPENNIS.

Bohem. Mon. Cassid. i. p. 50, 5.
Patr. Assam.

6. PRIOPTERA OCTONOTATA.

Bohem. Mon. Cassid. i. p. 51, 6.
Prioptera, 8, punctata? *Hope*, Col. Man. iii. t. ii. f. 2.
Patr. Celebes.

7. PRIOPTERA BIMACULATA. B.M.

Bohem. Mon. Cassid. i. p. 52, 7.
Cassida bimaculata, *Thurb.* Nov. Ins. Spec. p. 86, t. v. f. 93. *Schœnh.* Syn. Ins. ii. p. 221, 91.
Cassida bimacula, *Herbst.* Naturs. viii. p. 262, 32, t. cxxxii. f. 4.
Patr. China. Ind. or. (Mus. Brit.)

B. *Prothorax nigro-maculatus.*

8. PRIOPTERA BIPUNCTICOLLIS.

Sub-rotundata, convexa, flavo-testacea, sub-nitida; antennis articulo ultimo apice pectoreque postice fuscis; prothorace sub-lævi, antrorsum haud angustato, ante basin maculis duabus mediocribus, rotundis, nigris; elytris subtiliter, sub-remote punctulatis, immaculatis, basi retusis, antice leviter gibbosis, singulo in disco bi-foveolato. Long. 11, lat. 9, millim.

Bohem. Mon. Cassid. Append.
Patr. Java, Mus. Soc. Ind. Orient.

9. PRIOPTERA OCTOPUNCTATA. B.M.

Bohem. Mon. Cassid. i. p. 53, 8.
Cassida octopunctata, *Fabr.* Ent. Syst. i. p. 296, 21. Mant. Ins. p. 63, 18. Syst. El. i. p. 395, 41. *Linn.* Syst. Nat. Gmel. i. iv. p. 1636, 35. *Oliv.* Ent. vi. p. 946, 37, 97, t. iii. f. 38. *Herbst.* Naturs. viii. p. 334, 96. *Schœnh.* Syn. Ins. ii. p. 218, 60.

Patr. Java, Borneo.

10. PRIOPTERA DECEMPUSTULATA.

Bohem. Mon. Cassid. i. p. 55, 9.
Basiprionota octopunctata, *Dej.* Cat. ed. 3, p. 367.
Patr. Malacca Indiæ orientalis.

11. PRIOPTERA OCTOMACULATA.

Bohem. Mon. Cassid. i. p. 56, 10.
Basiprionota octomaculata, *Dej.* Cat. ed. p. 367.
Patr. Java.

12. PRIOPTERA DECEMSTILLATA.

Sub-rotundata, rufo-testacea, nitida, convexa; antennis articulo ultimo pectoreque postice nigris; prothorace subtiliter, sub-remote punctulato, angulis anticis magis prominulis rotundatis, dorso utrinque macula sat magna, rotunda, nigra; elytris mediocriter, vage crebre punctatis, pone basin leviter gibbosis, posterius valde ampliatis, singulo maculis

quatuor magnis foveolaque profunda disci antici notatæ. Long. 9½, lat. 8 millim.

Bohem. Mon. Cassid. Append.
Patr. India orientalis, Mus. Dom. *Deyrolle.*

13. PRIOPTERA MÆRKELI.

Bohem. Mon. Cassid. i. p. 57, 11.
Patr. Java.

II. Elytra æqualia, non gibbosa.

A. *Prothorax immaculatus.*

14. PRIOPTERA QUADRISIGNATA. B.M.

Bohem. Mon. Cassid. i. p. 58, 12.
Patr. Manilla (Ins. Philipp.) Mus. Brit.

15. PRIOPTERA BINOTATA.

Sub-rotundata, convexa, dilute flavescens, parum nitida; prothorace sub lævi, lateribus ad medium rotundato ampliato, dein recto; elytris crebre subtiliter, vage punctulatis, margine basali angustissime nigris; singulo macula parva in callo suprahumerali nigra. Long. 8½, lat. 6½ millim.

Bohem. Mon. Cassid. Append.
Patr. Ins. Philipp. Mus. Dom. *Waterhouse.*

16. PRIOPTERA OCTOPUSTULATA. B.M.

Sub-rotundata, convexa, dilute flavo-testacea, sub-nitida; prothorace sub-lævi; elytris subtiliter, sat crebre, antice sub seriatim, postice vage punctulatis, basi non retusis, disco bifoveolato, singulo maculis quatuor mediocribus, nigris, duabus ante binisque pone medium, transversim positis. Long. 8¾, lat. 7¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Insul. Philip. Mus. Britann. (Coll. *Cuming.*)

17. PRIOPTERA DECEMNOTATA. B.M.

Bohem. Mon. Cassid. i. p. 59, 13.
Cassida sinuata, *Fabr.* Syst. El. i. p. 398, 60. Ent. Sys. i. p. 298, 29. *Herbst.* Naturs. viii. p. 336, 99. *Schœnh.* Syn. Ins. ii. p. 221, 81. *Oliv.* Encycl. Ins. v. p. 392, 78. Entom. vi. p. 949, 42, 97, t. v. f. 11.

Var. a. genuinæ simillima sed præter maculas ordinarias, macula parva suturali ante apicem, communis elongata.
Var. b. elytrorum maculis parvis, punctiformibus.
Patr. Manilla. *Var. a* et *b.* Insul. Philipp. Mus. Britann.

B. *Prothorax nigro-bimaculatus.*

18. PRIOPTERA DECEMMACULATA. B.M.

Bohem. Mon. Cassid. i. p. 60, 14.
Patr. Ad montes Himalayæ Ind. orientalis.

19. PRIOPTERA PALLIDICORNIS.

Bohem. Mon. Cassid. i. p. 61, 15.
Patr. Tenasserim Indiæ orientalis.

20. PRIOPTERA DECEMSIGNATA.

Bohem. Mon. Cassid. i. p. 62, 16.
Patr. Assam Indiæ orientalis.

21. PRIOPTERA WHITEI. B.M.

Sub-rotundata, convexa, flavo-testacea, nitida; antennis articulis duobus ultimis nigris; prothorace subtilissime, remote punctulato, tenuiter canaliculato, medio maculis duabus magnis, remotis, nigro-fuscis; elytris subtiliter, sub-remote punctulatis, disco antico bi-foveolatis, singulo vitta latissima ante apicem desinente maculaque marginali infra medium, nigris; pectore postice infuscato. Long. 10, lat. 8½ millim.

Bohem. Mon. Cassid. Append.
Patr. China Mus. Britann. (Coll. *R. Fortune*) et Mus. Holm.

Genus 8. HIMATIDIUM.

Illig. Mag. iii. p. 131, 217. *Perty*, Delect. Anim. p. 102. *Bohem.* Mon. Cassid. i. p. 63.
Imatidium, *Fabr.* Syst. El. i. p. 345. *Schœnh.* Syn. Ins. ii. p. 134. *Dej.* Cat. ed. 3. p. 367.
Cassida, *Oliv.* Ent. vi. p. 922. *Herbst.* Naturs. viii. p. 204.

1. HIMATIDIUM FASCIATUM. B.M.

Bohem. Mon. Cassid. i. p. 65, 1, tab. II. B.
Imatidium fasciatum, *Fabr.* Syst. El. i. p. 346. *Illig.* Mag. i. p. 392. 2. *Schœnh.* Syn. Ins. ii. p. 134, 2. *Dej.* Cat. ed. 3, p. 367.
Imatidium comptum, *Perty*, Delect. Anim. Brasil. p. 101, t. xx. f. 8.
Cassida fasciata, *Oliv.* Ent. vi. p. 971, 84, t. vi. f. 100.
Cassida capensis, *Herbst.* Naturs. viii. p. 278, 51, t. cxxxiii. f. 10.

Var. a. prothorace immaculato, elytrorum fascia postica utrinque abbreviata, in marginem non excurrente.
Patr. Cayenna, Amazon (Para) Coll. *H. W. Bates.*

2. Himatidium semifasciatum.

Sub-ovatum, leviter convexum, dilute flavum, nitidum, antennis ferrugineis; capite nigro; prothorace evidenter, remote punctato, basi maculis duabus obliquis scutelloque nigris; elytris mediocriter punctato striatis basi plagis tribus, media communi et pone medium fasciis duabus atris, utrinque abbreviatis, in marginem non excurrentibus. Long. 6, lat. 5 millim.
Bohem. Mon. Cassid. Append.
Patr. Pernambuco. Mus. Dom. *Dohrn.*

3. Himatidium quatuordecimmaculatum.

Bohem. Mon. Cassid. i. p. 66, 2.

Imatidium quatuordecimmaculatum, *Latr.* Humboldt et Bonpland Observ. de Zool. et d'Anat. comp. i. p. 211, t. xv. f. 6, Ann. de Mus. Nat. i. p. 294, t. xxi. f. 10. *Dej.* Cat. ed. 3, p. 367.
Cassida quatuordecimmaculatum, *Schœnh.* Syn. Ins. ii. p. 230, 155.
Patr. Columbia.

4. Himatidium semicirculare. B.M.

Bohem. Mon. Cassid. i. p. 67, 3.
Cassida semicircularis, *Oliv.* Entom. vi. p. 970, 82, 97, t. vi. f. 99.
Imatidium semicirculare, *Latr.* Humboldt et Bonpland Observ. de Zool. et d'Anat. comp. p. 213, t. xv. f. 7. *Dej.* Cat. ed. 3, p. 367.
Imatidium annulum, *Dej.* Cat. ed. 2, p. 115.
Var. a. elytrorum annulo flavo antice utrinque abbreviato, circum semicircularem referente.
Patr. Columbia, *Var. a.* Mus. Dom. *Chevrolat.*

5. Himatidium impurum.

Bohem. Mon. Cassid. i. p. 68, 4.
Patr. Brasilia.

6. Himatidium nimbatum.

Perty, Delect. Anim. Brasil. p. 102, t. xx. f. 10. *Bohem.* Mon. Cassid. i. p. 69, 5.
Patr. Brasilia.

7. HIMATIDIUM THORACICUM. B. M.

Bohem. Mon. Cassid. i. p. 70, 6.
Imatidium thoracicum, *Fabr.* Syst. El. i. p. 346, 3. *Schœnh.* Syn. Ins. ii. p. 134, 3. *Dej.* Cat. ed. 3, p. 367.
Imatidium albicolle, *Latr.* Humboldt et Bonpland Observ. de Zool. et d'Anatom. comp. i. p. 215.
Cassida albicollis, *Oliv.* Ent. vi. p. 974, 86, 97, t. vi. f. 105.
Patr. Brasilia, Columbia (Coll. *Goudot*), Cayenna. Amazon (Para) Coll. *H. W. Bates.*

8. HIMATIDIUM LINEOLA.

Bohem. Mon. Cassid. i. p. 11, 7.
Imatidium Lineola, *Fabr.* Syst. El. i. p. 346, 4. *Schœnh.* Syn. Ins. ii. p. 134, 4. *Dej.* Cat. ed. 3, p. 367.
Imatidium marginicolle, *Dej.* in litt.
Cassida pallipes *Oliv.* Ent. vi. p. 973, 88, 97, t. vi. f. 104.
Cassida collaris, *Herbst.* Naturs. viii. p. 259, 28, t. cxxx. f. 7. *Schœnh.* Syn. Ins. ii. p. 216, 23.
Patr. Cayenna.

9. HIMATIDIUM CYANIPENNE.

Bohem. Mon. Cassid. i. p. 72, 8.
Patr. Brasilia.

10. HIMATIDIUM RUFIVENTRE.

Bohem. Mon. Cassid. i. p. 73, 9.
Patr. Brasilia.

11. HIMATIDIUM CHALYBEUM.

Bohem. Mon. Cassid. i. p. 73, 10.
Patr. Para.

12. HIMATIDIUM RUFOMARGINATUM.

Bohem. Mon. Cassid. i. p. 75, 12.
Patr. Mexico.

13. HIMATIDIUM CINCTUM.

Bohem. Mon. Cassid. i. p. 76, 13.
Imatidium cinctum, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 285.
Patr. Brasilia, Columbia.

14. HIMATIDIUM DISCOIDEUM.

Bohem. Mon. Cassid. i. p. 77, 14.
Patr. Bahia.

15. Himatidium rubrum.

Bohem. Mon. Cassid. i. p. 78, 15.
Patr. Brasilia.

16. Himatidium fulvum. B.M.

Bohem. Mon. Cassid. i. p. 79, 16.
Patr. Columbia.

17. Himatidium marginicolle.

Bohem. Mon. Cassid. i. p. 80, 17.
Patr. Brasilia.

18. Himatidium rubricatum.

Bohem. Mon. Cassid. i. p. 81, 18.
Imatidium rubricatum, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 285.
Patr. Cayenna.

Genus 9. CALLIASPIS.

Dej. Cat. ed. 3, p. 367. *Bohem.* Mon. Cassid. i. p. 81.
Cassida, *Oliv.* Ent. vi. p. 922.
Cephaloleia, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 284.

1. Calliaspis cassidoides.

Cephaloleia cassidoides, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 284.
Calliaspis rufina, *Bohem.* Mon. Cassid. i. p. 83, 1.
Patr. Java.

2. Calliaspis cinnabarina. B.M.

Bohem. Mon. Cassid. i. p. 84, 2, t. ii. c.
Calliaspis cinnabaris, *Lacordaire*, Dej. Cat. ed. 3, p. 367.
Patria. Cayenna, Brasilia (Coll. Maj. Gen. *Hardwicke*).

3. Calliaspis porphyrio.

Bohem. Mon. Cassid. i. p. 85, 3.
Patr. Para.

4. Calliaspis rubra. B.M.

Dej. Cat. ed. 3, p. 367. *Bohem.* Mon. Cassid. i. p. 86, 4.
Cassida rubra, *Oliv.* Ent. vi. p. 974, 90, 97, T. N. f. 106.
Patr. Cayenna.

5. Calliaspis rufula.

Bohem. Mon. Cassid. i. p. 87, 5.
Patr. Cayenna.

6. Calliaspis resplendens.

Himatidium resplendens, *Bohem.* Mon. Cassid. i. p. 74, 11.
Patr. Cayenna.

7. Calliaspis discophora.

Lacordaire, Dej. Cat. ed. 3, p. 367. *Bohem.* Mon. Cassid. i. p. 88, 6.
Patr. Cayenna.

8. Calliaspis funeraria.

Lacordaire, Dej. Cat. ed. 3, p. 367. *Bohem.* Mon. Cassid. i. p. 88, 7.
Patr. Cayenna.

9. Calliaspis bicolor.

Sub-ovata, modice convexa, supra nigra, nitida; antennis, capite, corpore pedibusque testaceis; prothorace remote, evidenter, dorso subtiliter, crebrius punctulato, extus anguste ferrugineo-marginato; elytris subtiliter, regulariter punctato striatis, interstitiis punctis subtilissimis, valde remotis impressis. Long. 4½, lat. 3½ millim.
Bohem. Mon. Cassid. Append.
Patr. Honda Boliviæ. Mus. Dom. *Chevrolat.*

Genus 10. PORPHYRASPIS.

Hope, Coleopt. Man. iii. p. 154. *Bohem.* Mon. Cassid. i. p. 89.
Imatidium, *Fabr.* Syst. El. i. p. 345. *Schœnh.* Syn. Ins. ii. p. 134.
Cassida, *Germ.* Ins. Spec. i. p. 540.
Hemisphærota, *Chevrolat.* Dej. Cat. ed. 3, p. 367.

I. *Elytra inæqualia.*

1. Porphyraspis sanguinea. B.M.

Bohem. Mon. Cassid. i. p. 91, 1.
Imatidium sanguineum, *Fabr.* Syst. El. i. p. 346, 5. *Schœnh.* Syn. Ins. ii. p. 135, 5.
Hemisphærota corallina, *Dej.* Cat. ed. 3, p. 367.
Patr. Cayenna (Tapayos), Coll. H. W. Bates.

2. PORPHYRASPIS COCCINEA. B.M.

Bohem. Mon. Cassid. i. p. 92, 2.
Patr. Para Brasiliæ. Amazon (Tapayos) Coll. *H. W. Bates.*

3. PORPHYRASPIS COSTIPENNIS. B.M.

Bohem. Mon. Cassid. i. p. 93, 3.
Patr. Para Brasiliæ. Amazon (Santarem) Coll. *H. W. Bates.*

4. PORPHYRASPIS CIRCUMDATA.

Bohem. Mon. Cassid. i. p. 94, 4.
Patr. Columbia.

5. PORPHYRASPIS TRISTIS. B.M.

Bohem. Mon. Cassid. i. p. 95, 5.
Hemisphærota tristis, *Dej.* Cat. ed. 3, p. 367.
Patr. Brasilia (Tapayos. Coll. *H. W. Bates*).

6. PORPHYRASPIS INTRICATA. B.M.

Bohem. Mon. Cassid. i. p. 95, 6.
Patr. Brasilia.

7. PORPHYRASPIS RUGOSA.

Bohem. Mon. Cassid. i. p. 96, 7.
Patr. Brasilia.

II. *Elytra æqualia.*

8. PORPHYRASPIS PALMARUM. B.M.

Ovata, convexa, rufo-testacea, sub-nitida; prothorace dorso utrinque impresso, punctato, medio longitudinaliter elevato, canaliculato, lateribus profunde punctato; elytris profunde, rude, minus regulariter striata-punctatis, disco parce fusco-variegatis. Long. 5, lat. 3½ millim.
Bohem. Mon. Cassid. Append.
Porphyraspis Lataniæ. *Sallé MS.*
Patr. St. Domingo.—Dom. *Chevrolat* et *Deyrolle.* Mus. Reg. Holm.—Pernambuco Mus. Dom. *Dohrn.*

9. PORPHYRASPIS CYANEA. B.M.

Imatidium cyaneum, *Say* Journ. Acad. of Philad. iii. p. 435.
Cassida erythrocera, *Germ.* Ins. Spec. p. 540, 731.
Hemisphærota erythrocera, *Dej.* Cat. ed. 3, p. 367.
Porphyraspis erythrocera, *Bohem.* Mon. Cassid. i. p. 97, 8, t. ii. D.
Patr. Florida et Kentucky Americæ septentrionalis (St. John's Bluff) Coll. *Edw. Doubleday.*

10. Porphyraspis xanthocera.

Bohem. Mon. Cassid. i. p. 98, 9.
Patr. Cuba.

11. Porphyraspis mulsanti.

Sub-rotundata, convexa, nitida, supra cyanea, subtus nigra; antennis totis tarsisque flavis; prothorace disco remote, subtiliter, lateribus et antice profundius punctato, angulis anticis anguste luteo-marginatis; elytris profunde, crebre punctato-striatis, singulo basi plaga magna transversa testacea, macula parva cærulea notata. Long. 5, lat. 3¾, millim.
Bohem. Mon. Cassid. Append.
Patr. Brasilia Mus. Dom. *Chevrolat.*

12. Porphyraspis Beskei.

Bohem. Mon. Cassid. i. p. 99, 10.
Patr. Brasilia, prope novum Friburgum.

13. Porphyraspis miniata. B.M.

Breviter rotundata, convexa, dilute miniata, nitida; prothorace convexo, sub-lævi, lateribus profunde punctato; elytris crebre, profunde punctato-striatis, interstitiis angustis, modice convexis; margine transversim rugoso. Long. 4, lat. 3¾ millim.
Bohem. Mon. Cassid. Append.
Patr. Columbia Mus. Britann. (Coll. *Goudot.*)

Genus 11. CALYPTOCEPHALA.

Chevrolat, Dej. Cat. ed. 3, p. 367. *Bohem.* Mon. Cassid. i. p. 100. Cassida, *Germ.* Ins. Spec. i. p. 540.

1. Calyptocephala nigricornis. B.M.

Dej. Cat. ed. 3, p. 367. *Bohem.* Mon. Cassid. i. p. 101, 1, t. ii. E. Cassida nigricornis, *Germ.* Ins. Spec. p. 540, 730.
Patr. Brasilia.

2. Calyptocephala lutea. B.M.

Bohem. Mon. Cassid. i. p. 102, 2.
Patr. Pernambuco Brasiliæ, Cayenna.

3. Calyptocephala brevicornis. B.M.

Bohem. Mon. Cassid. i. p. 103, 3.
Patr. Brasilia Para (Coll. *H. W. Bates*). Venezuela (Coll. *Dyson*).

4. Calyptocephala punctata.

Bohem. Mon. Cassid. i. p. 104, 4.
Patr. Chiquitos.

5. Calyptocephala marginipennis.

Bohem. Mon. Cassid. i. p. 105, 5.
Patr. Mexico.

Genus 12. SPILOPHORA.

Bohem. Mon. Cassid. i. p. 107.
Imatidium, *Fabr.* Ent. Syst. i. p. 345. *Schœnh.* Syn. Ins. ii. p. 134.
Calyptocephala, *Dej.* Cat. ed. 3, p. 367. *Erichs.* Archiv. f. Naturg. xiii. p. 151.

1. Spilophora trimaculata.

Bohem. Mon. Cassid. i. p. 108, 1.
Imatidium trimaculatum, *Fabr.* Syst. El. i. p. 345, 1. *Schœnh.* Syn. Ins. ii. p. 134, 1.
Patr. Surinam.

2. Spilophora trigemina. B.M. var.?

Bohem. Mon. Cassid. i. p. 109, 2.
Calyptocephala trigemina, *Lacordaire*, Dej. Cat. ed. 3, p. 367. *Guér.* Icon. de Règne Anim. Ins. ii. texte, p. 285.
Patr. Cayenna.

3. Spilophora pulchra. B.M.

Sub-triangularis, modice convexa, dilute flavescens, nitida; antennis leniter infuscatis; prothorace parvo, lateribus rotundato, punctato; elytris dorso mediocriter punctato-striatis, extrorsum vage punctatis, basi fascialata, utrinque extrorsum excurrente plagaque postica transversa, communibus, cæruleis. Long. 7, lat. 5½ millim.
Bohem. Mon. Cassid. Append.
Patr. Rio Tapayos Mus. Britann. (Coll. *H. W. Bates.*)

4. Spilophora sellata. B.M.

Sub-triangularis, modice convexa, flavo-testacea, nitida; prothorace parvo, lateribus rotundato, basi marginibusque remote punctato; elytris subtiliter, regulariter punctato-striatis, cyaneis, basi utrinque macula magna, obliqua mar-

gineque postico, sat late inæqualiter flavescentibus. Long. 6½, lat. 5¼ milim.
Bohem. Mon. Cassid. Append.
Patr. Ega Amazon. Mus. Britann. (Coll. *H. W. Bates.*)

5. Spilophora Peruana.

Calyptocephala Peruana. *Erichs.* Archiv. f. Naturg. xiii. p. 157, 1.
Bohem. Mon. Cassid. i. p. 106, 3.
Patr. Peruvia Orientalis "Montana de Vitoc."

TRIBUS II.

Genus 13. TAUROMA.

Hope, Ann. Nat. Hist. iii. p. 97, t. iv. f. 4. *Bohem.* Mon. Cassid. i. p. 113.
Cassida, *Linn.* Syst. Nat. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *Herbst.* Naturs. viii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209.
Omocera Chevrolat. *Dej.* Cat. ed. 3, p. 367.
Polychalca Chevrolat. *Dej.* Cat. ed. 3, p. 368.

I. Elytra sub-quadrata, spina valida humerali literaliter prominula.

A. *Elytra dorso convexa.*

1. Tauroma bicornis. B.M.

Bohem. Mon. Cassid. i. p. 115. 1, t. iii. A.
Cassida bicornis. *Fabr.* Syst. El. i. p. 403, 86. Ent. Syst. i. p. 301, 44. Mant. Ins. i. p. 64, 36. Spec. Ins. i. p. 112, 29. Syst. Ent. p. 91, 18. *Linn.* Syst. Nat. i. ii. p. 576, 8. Ed. Gmel. i. iv. p. 1638, 8. Amoen. Acad. vi. p. 293, 9. Cent. Ins. no. 9. *Herbst.* Naturs. viii. p. 328, 89, t. cxxxvi. f. 6. *Oliv.* Enc. meth. v. p. 587, 48. Entom. vi. p. 934, 16, 97. t. iv. f. 59. *Schœnh.* Syn. Ins. ii. p. 224, 113.
Cassida aculeata, *Voet.* Col. ii. t. xliii. f. 19.
Omocera bicornis, *Dej.* Cat. ed. 3, p. 367.
Patr. Surinam, Cayenna.

2. Tauroma viridicærulea.

Bohem. Mon. Cassid. i. p. 117, 2.
Patr. Para.

3. Tauroma creberrima. B.M.

Bohem. Mon. Cassid. i. p. 118, 3.
Patr. Columbia, (Coll. *Goudot*).

4. Tauroma smaragdina. B.M.

Bohem. Mon. Cassid. i. p. 119, 4.
Patr. (Ega) (Coll. *H. W. Bates.*)

5. Tauroma Ammon.

Oblongo-subquadrata, convexa, smaragdina, sub-nitida, anguste viridi-aureomarginata; prothorace profunde, crebre rugoso-punctato, lateribus oblique ampliato; elytris ad suturam seriatim, antice parcius, disco profunde, crebre, vage punctatis; corpore, femoribus ultra medium tibiisque posterioribus, apice excepto, cyaneis; humeris lateraliter extensis, tenuioribus, compressis, apice reflexis. Long. 11, lat. 8½ millim.
Bohem. Mon. Cassid. Append.
Patr. Moxos Peruviæ. Dom. Pavon. Mus. Berol.

6. Tauroma azureicornis. B.M.

Bohem. Mon. Cassid. i. p. 120, 5.
Omocerus azureicornis, *Chevrol.* Col. de Mexique Cent. ii. fasc. 5, no. 119.
Omocera azureicornis, *Dej.* Cat. ed. 3, p. 367.
Patr. Oaxaca Mexico.

7. Tauroma eximia. B.M.

Bohem. Mon. Cassid. i. p. 121, 6.
Patr. Guyana, Venezuela (Coll. *Dyson* and *Birschell*).

8. Tauroma aureicornis.

Bohem. Mon. Cassid. i. p. 122, 7.
Omocera aureicornis. *Blanch.* Orbign. Voy. de l'Amér. mér. Ins. pl. 24, f. 2.
Patr. Bolivia. St. Crux de la Sierra.

9. Tauroma æruginosa. B.M.

Bohem. Mon. Cassid. i. p. 123, 8.
Patr. Surinam. Para (*Gordon Grahame*, Esq.)

10. Tauroma boliviana.

Sub-quadrata, convexa, fusco-cuprea, supra sub-opaca, subtus nitida; prothorace remote, dorso postico evidentius punctato, medio breviter canaliculato, lateribus ultra medium oblique ampliato, dein fere recto; elytris sat profunde, remote, vage punctatis, dorso antico gibbosis; humeris lateraliter extensis, validis, brevibus, apice non reflexis. Long. 10, lat. 10 millim.
Bohem. Mon. Cassid. Append.
Patr. Bolivia Dom. *Chevrolat.* Mus. Holm.

11. Tauroma Reichei.

Bohem. Mon. Cassid. i. p. 124, 9.
Patr. Para Brasiliæ, Columbia, Chili.

12. Tauroma ignava. B.M.

Bohem. Mon. Cassid. i. p. 125, 10.
Patr. Brasilia, (*T. V. Wollaston*, Esq.)

13. Tauroma Taurus. B.M.

Bohem. Mon. Cassid. i. p. 126, 11.
Cassida Taurus *Fabr.* Syst. El. i. p. 403, 87. Ent. Syst. i. p. 302, 45. Mant. Ins. ii. App. p. 380, 29-30. *Linn.* Syst. Nat. Gmel. i. iv. p. 1643, 33. *Herbst.* Naturs. viii. p. 344, 111. Oliv. Encycl. Ins. vi. p. 387, 49. Entom. vi. p. 935, 17, 97, t. ix. f. 60. a. b. *Schœnh.* Syn. Ins. ii. p. 225, 114.
Omocera Taurus, *Dej.* Cat. ed. 3, p. 367.
Tauroma punctipennis? *Hope*, Ann. Nat. Hist. iii. p. 97.
Patr. Havanna, Para Brasiliæ, Columbia, Cayenna.

14. Tauroma jugata.

Sub-quadrata, convexa, supra æneo-nigra, parum nitida, subtus cuprea, nitida; prothorace crebre, mediocriter, rugoso-punctato, lateribus ultra medium oblique ampliato, dein recto, medio canaliculato; elytris sat profunde, minus crebre, vage, introrsum sub-seriatim, extrorsum subtilius punctatis; humeris lateraliter extensis, validioribus, apice reflexis. Long. 12⅓, lat. 9½ millim.
Bohem. Mon. Cassid. Append.
Patr. Caraccas Mus. Dom. *Chevrolat.*

B. *Elytra dorso parum convexa.*

15. Tauroma truncata. B.M.

Bohem. Mon. Cassid. i. p. 128, 12.
Patr. Brasilia, Bahia.

16. Tauroma coracina.

Bohem. Mon. Cassid. i. p. 129, 13.
Patr. Bahia, Brasilia.

17. Tauroma cornuta. B.M.

Bohem. Mon. Cassid. i. p. 130, 14.
Omocera cornuta, *Dej.* Cat. ed. 3, p. 367.
Patr. Brasilia.

18. Tauroma anchoralis.

Bohem. Mon. Cassid. i. p. 131, 15.

Omocera anchoralis, *Dej.* Cat. ed. 3, p. 367.
Patr. Brasilia.

II. Elytra sub-triangularia, angulo humerali lateraliter parum prominulo.

19. Tauroma antiqua. B.M.

Bohem. Mon. Cassid. i. p. 132, 16.
Cassida antiqua. *Klug.* Preis-verzeichn. 1829. p. 8, 105.
Polychalca antiqua. *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia (St. Paul, St. Catherine).

20. Tauroma scabrosa.

Bohem. Mon. Cassid. i. p. 133. 17.
Patr. Brasilia meridionalis, Minas Geraës.

21. Tauroma gibbera. B.M.

Bohem. Mon. Cassid. i. p. 134, 18.
Patr. Bahia, Brasilia.

22. Tauroma malachitica.

Bohem. Mon. Cassid. i. p. 136, 19.
Cassida malachitica, *Germ.* Ins. Spec. i. p. 534, 722.
Cassida metallica, *Vigor's* Zool. Journ. ii. p. 240, t. ix. f. 9.
Polychalca pertusa, *Dej.* Cat. ed. 3, p. 368.
Patr. Minas Geraës Brasiliæ.

Genus 14. DESMONOTA.

Hope, Ann. Nat. Hist. iii. p. 97, t. iv. f. 5. *Bohem.* Mon. Cassid. i. p. 137.
Cassida, *Fabr.* Syst. El. i. p. 387. *Schœnh.* Syn. Ins. ii. p. 209. *Latr.* in *Humb.* et *Bonpl.* Observ. de Zool. ii. 353. *Germ.* Ins. Spec. i. p. 535.
Himatidium, *Perty* Delect. Animal. Brasil. p. 102.
Polychalca, *Dej.* Cat. ed. 3, p. 368. *Erichs.* Archiv. f. Naturg. xiii. p. 151.
Cyrtonota, *Chevrol. Dej.* Cat. ed. p. 368. ex parte.

I. Elytra rotundata.

1. Desmonota variolosa. B.M.

Bohem. Mon. Cassid. i. p. 139, 1.
Cassida variolosa, *Fabr.* Syst. El. i. p. 407, 111. *Weber.* Observ. Ent. i. p. 50, 1. *Schœnh.* Syn. Ins. ii. p. 228, 139.
Cassida smaragdina, *Vigor's* Zool. Journ. ii. p. 240, t. ix. f. 8.
Polychalca variolosa, *Dej.* Cat. ed. 3, p. 368.
Patr. Bahia, Brasiliæ.

2. Desmonota cariosa.

Bohem. Mon. Cassid. i. p. 140, 2.
Polychalca cariosa, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia meridionalis.

3. Desmonota denticulata. B.M.

Bohem. Mon. Cassid. i. p. 141, 3.
Polychalca denticulata, *Dej.* Cat. ed. 3, p. 368.
Patr. Paraguay, Bolivia ad Santa Cruz de la Sierra.

4. Desmonota salebrosa. B.M.

Bohem. Mon. Cassid. i. p. 142, 4.
Polychalca salebrosa, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia, Bolivia (Coll. *Bridges*).

5. Desmonota peruviana.

Sub-ovata, convexa, fusco-ænea, supra opaca, subtus cuprea, nitida; prothorace parum convexo, subtiliter, remote punctato, medio breviter, tenuiter canaliculato, utrinque foveolato, lateribus longe ultra medium oblique ampliato, dein recto, angulis posticis retrorsum productis, acutis; elytris subtiliter coriaceis, remote, mediocriter, dorso sub-seriatim punctatis, disco antico obsoletius bifoveolatis, postico sub-lævibus; margine obtuse crenulatis. Long. 11¾, lat. 8½ millim.
Bohem. Mon. Cassid. Append.
Patr. Peruvia, Dom. *Deyrolle* Mus. Holm.

6. Desmonota crenulata.

Sub-ovata, convexa, fusco-ænea, supra opaca, subtus cuprea, nitida; prothorace parum convexo, subtiliter, sat remote punctulato, medio breviter, tenuiter canaliculato, utrinque foveolato, lateribus ultra medium oblique ampliato, dein recto, angulis posticis retrorsum productis, acutis; elytris obsolete coriaceis, valde remote, parum profunde punctatis, singulo in disco antico bi-foveolato; margine obtuse crenulato. Long. 12, lat. 9 millim.
Bohem. Mon. Cassid. Append.
Patr. Patagonia, Mus. Dom. *Chevrolat.*

7. Desmonota ærea. B.M.

Bohem. Mon. Cassid. i. p. 143, 5.
Patr. Brasilia (Pernambuco) (Mrs. J. P. George Smith).

8. Desmonota monstrosa.

Bohem. Mon. Cassid. i. p. 144, 6.
Patr. Brasilia.

9. Desmonota decora.

Bohem. Mon. Cassid. i. p. 146, 7.
Himatidium decorum, *Perty*, Delect. Anim. Brasil. p. 102, t. xx. f. 9.
Patr. Minas Geraës Brasiliæ, locis montosis.

II. Elytra sub-ovata vel quadrata.

10. Desmonota turpis.

Bohem. Mon. Cassid. i. p. 147, 8.
Patr. Bahia.

11. Desmonata dentipennis. B.M.

Bohem. Mon. Cassid. i. p. 148, 9.
Patr. Bahia.

12. Desmonota platynota. B.M.

Bohem. Mon. Cassid. i. p. 149, 10, t. iii. B.
Cassida platynota, *Germ.* Ins. Spec. i. 535, 723.
Polychalca nodosa, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

13. Desmonota gravida.

Bohem. Mon. Cassid. i. p. 150, 11.
Patr. Chiquitos.

14. Desmonota multicava. B.M.

Bohem. Mon. Cassid. i. p. 151, 12.
Cassida multicava, *Latr.* Humb. et Bonpl. Observ. de Zool. et d'Anatom. comp. ii. p. 352, t. xxii. f. 8, 9.
Cyrtonota multicava, *Dej.* Cat. ed. 3, p. 368.
Polychalca multicava, *Erichs.* Archiv. f. Naturg. xiii. p. 151, 1.
Patr. Columbia (Coll. *Goudot*), Peruvia. Venezuela (Coll. *Birschell*).

15. Desmonota aurolimbata.

Bohem. Mon. Cassid. i. p. 152, 13.
Patr. Bahia.

16. Desmonota Duponti.

Bohem. Mon. Cassid. i. p. 153, 14.
Patr. Bahia.

17. Desmonota bispinosa. B.M.

Bohem. Mon. Cassid. i. p. 154, 15.
Patr. Brasilia.

18. Desmonota subcornuta. B.M.

Bohem. Mon. Cassid. i. p. 155, 16.
Polychalca subcornuta, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

19. Desmonota retusa.

Bohem. Mon. Cassid. i. p. 156, 17,
Patr. Brasilia.

20. Desmonota laticollis.

Bohem. Mon. Cassid. i. p. 157, 18.
Patr. Caraccas, Bahia.

21. Desmonota bicornuta. B.M.

Bohem. Mon. Cassid. i. p. 158, 19.
Patr. Bahia, Columbia, Venezuela (Coll. *Birschell*).

22. Desmonota quadrinodosa.

Oblongo-quadrata, modice convexa, obscure ænea, supra opaca, subtus nitida; prothorace lato, supra inæquali, subtiliter, confertim coriaceo, multifoveolato, canaliculato, lateribus antice oblique ampliato, dein recto; elytris mediocriter, sub-seriatim punctatis, singulo basi tuberculis duobus compressis, interiore majore, posterius in carinam introrsum flexam continuato, ante medium disci profunde foveolato. Long. 7½, lat. 5 millim.

Bohem. Mon. Cassid. Append.
Patr. — — — Mus. Dom. Dohrn.

23. Desmonota cribripennis. B.M.

Bohem. Mon. Cassid. i. p. 159, 20.
Patr. Bahia. var. Bolivia (Coll. *Bridges*).

24. Desmonota intermedia.

Bohem. Mon. Cassid. i. p. 160, 21.
Patr. Brasilia.

25. Desmonota marginiventris.

Bohem. Mon. Cassid. i. p. 162, 22.
Patr. Brasilia.

26. Desmonota metallica. B.M.

Bohem. Mon. Cassid. i. p. 163, 23.

Cassida metallica, *Klug*, Preis-Verzeichn, 1829, p. 8, 106.
Polychalca metallica, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia (St. Catherine).

27. Desmonota perforata. B.M.

Bohem. Mon. Cassid. i. p. 164, 24.
Patr. Bolivia (Coll. *Bridges*). Brazilia (St. Paul's).

28. Desmonota ruderaria.

Bohem. Mon. Cassid. i. p. 165, 25.
Polychalca ruderaria. *Erichs.* Archiv. f. Naturg. xiii. p. 151, 2.
Patr. Peruvia orientalis.

Genus 15. CANISTRA.

Erichs. Archiv. f. Naturg. xiii. p. 151. *Bohem.* Mon. Cassid. i. p. 166.
Cassida *Klug*, Preis-Verzeichn, 1829, p. 7.
Discomorpha *Chevrol. Dej.* Cat. ed. 3, p. 368.
Oxynodera *Guér.* Icon. de Règne Anim. Ins. ii. p. 289.

I. *Elytra umbonata.*

1. Canistra procera.

Bohem. Mon. Cassid. i. p. 167, 1.
Patr. Brasilia.

2. Canistra varicosa.

Rotundata, convexa, nigra, parum nitida, tenuiter breviter, griseo-pubescens; antennis basi ferrugineis; prothorace subtiliter, remote punctulato; elytris pone basin obtuse conico-gibbosis, remote cicatricoso-punctatis, parce obscure sanguineo reticulatis; margine late explanato, extrorsum remote punctato. Long. 15, lat. 14¾ millim.
Canistra varicosa, *Erichs.* Arch. f. Naturg. xiii. p. 152, 1. *Bohem.* Mon. Cassid. i. p. 175, 8. *Bohem.* Mon. Cassid. Append.
Patr. Peruvia orientalis.

3. Canistra irrorata. B.M.

Bohem. Mon. Cassid. i. p. 168, 2. t. iii. C.
Oxynodera irrorata, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 289.
Patr. Bolivia (Coll. *Bridges*). Brasilia.

4. Canistra nigro-ænea.

Rotundata, convexa, nigro-ænea, supra opaca, subtus parum nitida; prothorace breviter sub-triangulari, vix punctulato, lateribus pone apicem ad basin oblique ampliato; elytris remote, mediocriter, vage punctatis, dorso sat crebre foveolatis, pone basin ad suturam foveola majore insculptis; margine sat late explanato, punctis nonnullis remotis impresso. Long. 14½, lat. 12½ millim.

Bohem. Mon. Cassid. Append.

Patr. Bogota, Columbiæ; Mus. Holm. Dom. *Deyrolle.*

5. Canistra calamitosa.

Bohem. Mon. Cassid. i. p. 169, 3.

Patr. Bahia.

6. Canistra contaminata.

Bohem. Mon. Cassid. i. p. 170, 4.

Patr. Bahia Brasiliæ.

7. Canistra rubiginosa. B.M.

Bohem. Mon. Cassid. i. p. 172, 5.

Oxynodera rubiginosa, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 289.

Discomorpha ferrugata, *Dej.* Cat. ed. 3, p. 368.

Patr. Brasilia, Bolivia (Coll. *Bridges*).

8. Canistra plagosa. B.M.

Bohem. Mon. Cassid. i. p. 173, 6.

Patr. Brasilia.

9. Canistra scoriosa.

Bohem. Mon. Cassid. i. p. 174, 7.

Patr. Rio Grande, Brasiliæ; Montevideo, Bahia.

II. *Elytra leviter convexa, absque umbone.*

10. Canistra carbonaria.

Bohem. Mon. Cassid. i. p. 175, 9.

Cassida carbonaria, *Klug.* Preis-Verzeichn. 1829, p. 7, 89.

Discomorpha carbonaria, *Dej.* Cat. ed. 3, p. 368.

Oxynodera rugosa, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 289.

Discomorpha tenebrosa, *Blanch.* Voy. d'Orbigny, Ins. pl. 24, f. 3.

Patr. Brasilia meridionalis, Montevideo.

Genus 16. DOLICHOTOMA.

Hope, Ann. Nat. Hist. iii. p. 95, t. iv. f. 3. *Bohem.* Mon. Cassid. i. p. 176.
Cassida, *Linn.* Syst. Nat. i. ii. p. 574. *Fabr.* Syst. i. p. 387. *De Geer*, Ins. v. p. 178. *Herbst.* Natürs, xiii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209. *Germ.* Ins. Spec. i. p. 529, 530.
Discomorpha *Chevrol. Dej.* Cat. ed. 3, p. 368.
Oxynodera, *Hope*, Coleopt. Mam. iii. p. 184. *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 289.

I. *Elytra antice in dorso gibbosa.*

1. Dolichotoma lugens. B.M.

Bohem. Mon. Cassid. i. p. 178, 1.
Cassida lugens, *Germ.* Ins. Spec. i. p. 530, 716.
Patr. Bahia, Brasiliæ.

2. Dolichotoma inepta. B.M.

Bohem. Mon. Cassid. i. p. 179, 2.
Patr. Brasilia.

3. Dolichotoma aenea. B.M.

Bohem. Mon. Cassid. i. p. 180, 3.
Cassida ænea, *Germ.* Ins. Spec. i. p. 529, 715.
Discomorpha tristis, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

4. Dolichotoma tristis.

Bohem. Mon. Cassid. i. p. 181, 4.
Oxynodera tristis, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 289.
Patr. Guaragos, Chiquitos.

5. Dolichotoma mitis.

Bohem. Mon. Cassid. i. p. 182, 5.
Patr. Brasilia.

6. Dolichotoma clypeata.

Bohem. Mon. Cassid. i. p. 184, 6.
Discomorpha clypeata, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

7. Dolichotoma spurca.

Rotundata, convexa, nigra, supra sub-opaca, pube silacea inæqualiter adspersa, subtus magis nitida; prothorace lævi, utrinque macula magna, longitudinali, sanguinea notato, lateribus late rotundato-ampliato; elytris subtiliter, remote, disco crebrius punctatis, antice retusis, dein valide gibbosis, plaga basali communi, sutura pone gibbum plagaque maxima irregulari disci, obscure sanguineis, hac postice macula rotunda nigra notata; margine late explanato, lævi. Long. $16\frac{1}{3}$, lat. $14\frac{1}{3}$ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Dom. *Jekel*, Mus. Holm.

8. Dolichotoma variegata. B.M.

Bohem. Mon. Cassid. i. p. 185, 7. t. iii. D.
Cassida variegata, *Fabr.* Syst. El. i. p. 404, 95. Ent. Syst. i. p. 303, 54. Mant. Ins. i. p. 65, 45. Spec. Ins. i. p. 113, 38. Syst. Ent. p. 92, 24. *Linn.* Syst. Nat. i. ii. p. 576, 16. Ed. Gmel. i. ii. p. 1640, 16. *Herbst.* Naturs. viii. p. 286, 57. t. cxxxiv. f. 3, *De Geer*, Mem. Ins. v. p. 178, 2. t. xv. f. 6. *Oliv.* Enc. Ins. v. p. 389, 61. Entom, vi. p. 926, 2, 97, t. ii. f. 22. *Schœnh.* Syn. Ins. ii. p. 226, 121.
Cassida tigrina, *Voet.* Col. ii. t. xlii. f. 8, 9.
Discomorpha variegata, *Dej.* Cat. ed. p. 368.
Cassida retusa, *Herbst.* Naturs. viii. p. 292, 61, t. cxxxiv. f. 6. *Schœnh.* Syn. Ins. ii. p. 230, 162. (Individuum materia aliena impura tinctum.)
Dolichotoma retusa. *Bohem.* Mon. Cassid. i. p. 207, 28.
Patr. Cayenna, Surinam.

9. Dolichotoma Dromedarius.

Bohem. Mon. Cassid. i. p. 186, 8.
Patr. Para.

10. Dolichotoma miniata. B.M.

Bohem. Mon. Cassid. i. p. 187, 9.
Patr. Remedios Novæ Granadæ, Columbia (Coll. *Goudot*).

11. Dolichotoma nigropunctata.

Bohem. Mon. Cassid. i. p. 188, 10.
Patr. Columbia.

12. Dolichotoma Collicullus. B.M.

Bohem. Mon. Cassid. i. p. 189, 11.
Patr. Para, Amazon, (Rio Tapayos,) (Coll. *H. W. Bates.*)

13. Dolichotoma lanuginosa. B.M.

Bohem. Mon. Cassid. i. p. 190, 12.
Patr. Columbia, Venezuela (Coll. *Dyson*, *Birschell*).

14. Dolichotoma mirifica.

Late ovata, nigro-ænea, parum nitida, pube brevissima, pallida adspersa; prothorace breviter sub-triangulari, æneo, subtiliter, sat crebre punctulato, tenuiter canaliculato, lateribus obliquo; elytris antice retusis, obtuse gibbosis, evidenter, crebre rugoso-punctatis, sanguineis, undique anguste, cærulescenti-marginatis, callo suprahumerali, linea longitudinali prope gibbum maculisque disci cæruleis. Long. 14, lat. 11¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Caraccas, Dom. Com. *Mannerheim*, Mus. Holm.

15. Dolichotoma depilata.

Bohem. Mon. Cassid. i. p. 191, 13.
Patr. Guyana.

16. Dolichotoma pubescens.

Rotundata, convexa, viridi-ænea, parum nitida, pube densa pallida obsita; prothorace sub-triangulari, supra paullo convexo, subtiliter, minus crebre punctulato, postice canaliculato antice tenuiter carinato, lateribus obliquo; elytris subtilius, sat crebre rugoso-punctatis, antice retusis, gibbosis; margine obsoletius punctulato. Long. 15, lat. 13½ millim.

Bohem. Mon. Cassid. Append.
Patr. Caraccas Dom. *Chevrolat.* St. Esteban, Columbiæ, Dom. *Dohrn.* Mus. Reg. Holm.

17. Dolichotoma metallica. B.M.?

Bohem. Mon. Cassid. i. p. 192, 14.
Oxynodera metallica, *Guér.* Icon. de Règne Anim. Ins. texte. ii. p. 289.
Patr. Bolivia, Guarayos.

18. Dolichotoma cærulea.

Rotundata, convexa, obscure cærulea, supra sub-opaca, subtus nitida; prothorace breviter sub-triangulari, lævi antice tuberculo parvo, medio canaliculato, lineis circularibus impressis cincto, lateribus obliquo; elytris casi retusis, dein transversim obtuse gibbosis, grosse crebre, sed minus profunde rugoso-punctatis; margine late explanato, parce subtiliterque punctulato. Long. 13, lat. 12 millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia ad fluvium Amazonis, Dom. *Buquet*, Mus. Holm.

19. Dolichotoma chlorosoma.

Rotundata, convexa, supra smaragdina, parce cupreo variegata, subtus cærulescens, nitida; prothorace breviter triangulari, subtilissime, sub-remote punctulato, medio tenuiter canaliculato, ante apicem tuberculo parvo, parum elevato, lineolis circularibus cincto; elytris antice retusis, impressis, dein obtuse gibbosis, crebre grosse, at minus profunde rugoso-punctatis; margine subtiliter, minus crebre punctulato. Long. 11½, lat. 11 millim.

Bohem. Mon. Cassid. Append.
Patr. Para, Mus. Holm.

20. Dolichotoma bisbiplagiata. B.M.

Breviter rotundata, convexa, supra obscure viridi-ænea, subopaca, subtus nigra, nitida; prothorace sublævi, postice canaliculato, antice carinato; elytris subtiliter coriaceis, sat crebre, parum profunde rugoso-punctatis. antice retusis, valide gibbosis; singulo medio maculis duabus magnis, transversim positis, una dorsali, altera marginali, sanguineis; margine latissime explanato, remote, subtilius punctulato. Long. 15½, lat. 15½.

Bohem. Mon. Cassid. Append.
Patr. Guatimala (Coll. *Deby*).

21. Dolichotoma biplagiata. B.M.

Bohem. Mon. Cassid. i. p. 193, 15.
Oxynodera biplagiata. *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 290.
Discomorpha biplagiata. *Dej.* Cat. ed. 3, p. 368.
Var. c. ut genuina, sed singulo elytro pone gibbum macula parva, rotunda, sanguinea.
Patr. Columbia (Coll. *Goudot*).

22. Dolichotoma princeps. B.M.

Bohem. Mon. Cassid. i. p. 195, 16.
Patr. Columbia, Guatimala (Coll. *Scherzer*).

23. Dolichotoma bihamata. B.M.

Late ovata, convexa, supra cæruleo-virescens, parum nitida, pube pallida adspersa, subtus nigra, nitida; prothorace breviter triangulari, subtilissime punctulato, tenuiter cana-

liculato; elytris subtiliter coriaceis, minus crebre punctulatis; margine valde explanato, ante medium vitta longitudinali, obscure sanguinea, intus medio ramulum recurvatum, concolorem emittente. Long. 14, lat. 12 millim.

Bohem. Mon. Cassid. Append.
Patr. Trinidad? Mus. Britann.

24. Dolichotoma peruviana.

Bohem. Mon. Cassid. i. p. 196, 17.
Patr. Peruvia.

25. Dolichotoma binotata.

Bohem. Mon. Cassid. i. p. 197, 18.
Patr. Para.

26. Dolichotoma mirabilis.

Rotundata, obscure ænea, parum nitida; antennis longissimis; prothorace brevi, subtilissime, remote punctulato, antice utrinque leviter rotundato-emarginato, lateribus sensim rotundato-ampliato; elytris basi retusis, valide tuberculatis, crebre, minus profunde punctatis, singulovista lata intramarginali plagaque magna, in dorso postico sanguineis, hac macula ovata ænea notata cum vitta marginali connexa; margine late explanato, prope discum transversim rugoso. Long. 18½, lat. 17½ millim.

Bohem. Mon. Cassid. Append.
Patr. Campo del Sacramento Dom. *Castelnau*, Mus. Paris.

27. Dolichotoma Batesi. B.M.

Rotundata, convexa, parum nitida, parce tenuiter pallido-pubescens, supra viridis, subtus cærulea; prothorace sublævi, maculis obscure cærulescentibus variegato, lateribus obliquis, angulis posticis acutis; elytris sat crebre, profunde rugoso-punctatis, antice retusis, gibbosis, singulo medio plaga maxima sub-triangulari, dilute sanguinea, disco exteriore margineque viridi maculata. Long. 16, lat. 14 millim.

Bohem. Mon. Cassid. Append.
Patr. Ega ad fluvium Amazonum. Dom. *Bates*, Mus. Brit.

28. Dolichotoma Grayi. B.M.

Rotundata, convexa, obscure viridis, parum nitida, subtus nigra, magis nitida; prothorace breviter triangulari, subtilissime punctulato, antice plaga parva rotunda, sub-elevata, nitida notato, lateribus oblique ampliato, angulis posticis

valde acutis; elytris crebre rugoso-punctatis, antice retusis, gibbosis; singulo medio plaga magna sanguinea viridi-quinquemaculata. Long. 14½, lat. 14 millim.

Bohem. Mon. Cassid. Append.
Var. a. plaga elytrorum sanguinea æneo trimaculata.
Patr. Santarem Amazon (Coll. *Bates*), Mus. Britann. *Var. a.* Mus. Dom. *Deyrolle.*

29. Dolichotoma ducalis.

Bohem. Mon. Cassid. i. p. 198, 19.
Patr. Brasilia.

30. Dolichotoma distigma.

Bohem. Mon. Cassid. i. p. 199, 20.
Var. a. macula laterali elytrorum immaculata.
Patr. Para.

31. Dolichotoma tarsata.

Bohem. Mon. Cassid. i. p. 200, 21.
Patr. Bahia.

32. Dolichotoma palliata. B.M.

Bohem. Mon. Cassid. i. p. 201, 22.
Cassida palliata, *Fabr.* Syst. El. i. p. 407, 110. Ent. Syst. i. p. 305, 63. Mant. Ins. ii. p. 380, 43, 44. *Herbst.* Naturs. viii. p. 347, 117. *Linn.* Syst. Nat. Gmel. i. iv. p. 1643, 35. *Schœnh.* Syn. Ins. ii. p. 227, 138.
Cassida marginata, *Voet.* Col. t. xliii. f. 14.
Cassida luctuosa, *Oliv.* Ins. vi. p. 930, 8, 97, t. iv. f. 54.
Cassida longicornis, *Guér.* Icon. de Règne Anim. Ins. p. 48, f. 5.
Discomorpha palliata, *Dej.* Cat. ed. 3, p. 368.
Patr. Cayenna Surinam.

II. *Elytra leviter convexa, absque gibbo.*

33. Dolichotoma strigata.

Bohem. Mon. Cassid. i. p. 202, 23.
Discomorpha strigata, *Hoffmansegg, Dej.* Cat. ed. 3, p. 368.
Cassida rubro-fasciata, *Voet.* Col. t. xlii. f. 10, 11.
Patr. Brasilia.

34. Dolichotoma puberula.

Bohem. Mon. Cassid. i. p. 204, 24.
Patr. Guyana.

35. Dolichotoma sericata. B.M.

Bohem. Mon. Cassid. i. p. 205, 25.
Oxynodera sericata, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 289.
Patr. Bolivia, Guarayos. Para (Coll. *H. W. Bates*).

36. Dolichotoma nigritarsis.

Rotundata, rufo-testacea, sub-opaca, pube brevi, pallida parce adspersa; clypeo, oculis, antennarum articulis, basalibus paginæ inferiori exceptis, tibiis apice tarsisque totis nigris; prothorace obsolete punctulato; elytris crebre, sat rude dorso seriatim, extrorsum vage punctatis, pone basin evidentius gibbosis; singulo in dorso bi-carinato carina interiore ante medium desinente. Long. 11½, lat. 10½ millim.

Bohem. Mon. Cassid. Append.
Patr. Pernambuco Dom. *Dohrn.* Mus. Berol.

37. Dolichotoma perforata.

Bohem. Mon. Cassid. i. p. 206, 26.
Patr. Columbia.

Species mihi ignota.

38. Dolichotoma chloris.

Bohem. Mon. Cassid. i. p. 207, 27. *Hope*, Ann. Nat. Hist. iii. p. 96, t. iv. f. 2.
Patr. Insula St. Vincenti.

Genus 17. CALASPIDEA.

Hope Coleopt. Man. iii. p. 183, t. iii. f. 1. *Bohem.* Mon. Cassid. i. p. 208.
Cassida, *Linn.* Syst. Nat. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *De Geer*, Mem. i. p. 176. *Herbst.* Naturs. viii. p. 204. *Oliv.* Entom. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209.
Eugenysa, *Chevrol. Dej.* Cat. ed. iii. p. 368.
Cyrtonota, *Chevrol. Dej.* Cat. ed. 3, p. 368, ex parte.

1. Calaspidea grossa. B.M.

Bohem. Mon. Cassid. i. p. 209, 1, t. iii. E.
Cassida grossa, *Fabr.* Syst. El. i. p. 405, 103. Ent. Syst. i. p. 304, 58. Mant. Ins. i. p. 65, 48. Spec. Ins. i. p. 113, 40. Syst. Ent. p. 92, 25. *Linn.* Syst. Nat. i. ii. p. 577, 17. Ed. Gmel. i. iv. p. 1641, 17. *Herbst.* Naturs. viii. p. 288, 58. t. cxxx. f. 1. *Oliv.* Encycl. v. p. 389, 64. Entom. vi. p. 926,

t. i. f. a. b. Sulzer, Ins. iii. f. 1. Naturs. vi. p. 89, t. iv. f. 1. *Roem.* Ins. p. 6, 15, t. iii. f. 1. *De Geer*, Mem. v. p. 176, 1, t. xv. f. 5. *Schœnh.* Syn. Ins. ii. p. 226, 130.
Cassida decussata, *Fabr.* Mant. Ins. i. p. 66, 57. *Spec.* Ins. i. p. 115, 48. Syst. Ent. p. 93, 32. *Linn.* Syst. Nat. ed. Gmel. i. iv. p. 1642, 28.
Eugenysa grandis, *Dej.* Cat. ed. 3, p. 368.
Patr. Surinam, Cayenna.

2. Calaspidea venosa. B.M.

Bohem. Mon. Cassid. i. p. 211, 2.
Cassida venosa, *Fabr.* Syst. El. i. p. 405, 99. Suppl. p. 84, 56, 57. *Herbst.* Naturs. viii. p. 437, 116. *Schœnh.* Syn. Ins. ii. p. 226, 125.
Eugenysa grossa, *Dej.* Cat. ed. 3, p. 368.
Patr. Cayenna.

3. Calaspidea columbiana.

Bohem. Mon. Cassid. i. p. 212, 3.
Patr. Columbia.

4. Calaspidea imperialis. B.M.

Rotundata, convexa, obscure cærulea, parum nitida, subtus nigra, magis nitida; prothorace crebre, subtiliter punctulato, lateribus ultra medium magis, dein sensim rotundato; elytris irregulariter elevato-reticulatis, reticulo nitido, punctato, areolas opacas, obsolete punctulatas includente; margine late explanato, subtiliter, sat crebre punctulato, vitta magna, lata, dilute sanguinea, maculis parvis feriatis, rotundis, remotis, impressis, cæruleis notato. Long. 18, lat. 18½ millim.

Bohem. Mon. Cassid. Append.
Var. a. vitta marginali elytrorum immaculata apicem versus vix angustata.
Patr. Ega (Amazon), (Coll. *H. W. Bates*). Mus. Britann. *Var. a.* Mus. Paris.

5. Calaspidea regalis. B.M.

Bohem. Mon. Cassid. i. p. 213, 4.
Patr. Brazil.

6. Calaspidea discors. B.M.

Bohem. Mon. Cassid. i. p. 214, 5.
Cassida discors, *Oliv.* Entom. vi. p. 938, 22, t. iv. f. 63.
Cassida cyanea, *Fabr.* Syst. El. i. p. 402, 84. Ent. Syst. i. p. 302, 46.

Mant. Ins. i. p. 65, 37. Spec. Ins. i. p. 112, 30. Syst. Ent. p. 91, 19. *De Geer*, Mem. v. p. 181, 5. t. xv. f. 9.
Cassida bipustulata, *Herbst.* Naturs. viii. p. 324, 86, t. cxxxvi. f. 3.
Cassida cædemadens Cat. Hamb. p. 40, 491.
Cyrtonota discors, *Dej.* Cat. ed. 3, p. 369.
Patr. Cayenna, Surinam.

7. Calaspidea guianiensis.

Rotundata (*mas*) aut ovata (*femina*), convexa, nitida, supra viridi-cærulea vel cærulea, subtus nigra; antennis basi nigris; prothorace subtiliter, minus crebre punctulato, antice utrinque anguste testaceo-marginato; elytris subtiliter, sat crebre punctulatis irregulariter reticulatis, reticulo tenui modice elevato, nitido, evidentius punctato, areolas mediocres includente; margine basi apiceque non reticulato; humeris antrorsum prominulis, apice leviter rotundatis. Long. *maris* 15, lat. 13½. Long. *feminæ* 18, lat. 14½ millim.

Bohem. Mon. Cassid. Append.
Calaspidea discors *var. a. Bohem.* Mon. Cassid. i. p. 214, 5.
Patr. Guyana.

8. Calaspidea signaticollis. B.M.

Rotundata, modice convexa, supra cærulea, sub-nitida, subtus nigra, magis nitida; antennis articulis tribus primis testaceis; prothorace subtiliter, minus crebre punctulato; margine antico utrinque anguste maculis, que duabus parvis pone medium, transversim positis testaceis; elytris undique irregulariter elevato-reticulatis, reticulo parum elevato, subnitido, punctulato, areolas opacas, obsolete punctulatas includente. Long. 15¼, lat. 15 millim.

Bohem. Mon. Cassid. Append.
Patr. — — — Mus. Britann.

9. Calaspidea Desmaresti.

Bohem. Mon. Cassid. i. p. 215, 6.
Patr. Bolivia.

10. Calaspidea parellina.

Bohem. Mon. Cassid. i. p. 216, 7.
Patr. Buenos Ayres.

Genus 18. MESOMPHALIA.

Hope, Ann. Nat. Hist. iii. p. 94, t. iv. f. 1. *Bohem.* Mon.

Cassid. i. p. 218. Cassida, *Linn.* Syst. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *Herbst.* Naturs. viii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209. *Germ.* Ins. Spec. i. p. 531.
Cyrtonota, *Chevrol. Dej.* Cat. ed. 3, p. 368. *Erichs.* Archiv. f. Naturg. xiii. p. 152.
Chelymorpha, *Chevrol, Dej.* Cat. ed. 3, p. 369, ex parte.

I. *Elytra ante medium dorsi umbonata.*

1. Mesomphalia quadraticollis. B.M.

Bohem. Mon. Cassid. i. p. 219, 1.
Cyrtonota quadricollis, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

2. Mesomphalia laticollis.

Ovata, convexa, nigro-ænea, sub-opaca, subtus nigra nitida; prothorace lato, subtiliter, sub-remote punctulato, parce pallido pubescente, lateribus ultra medium obliquo, dein recto; elytris remote, irregulariter, subtiliter punctulatis, antice retusis, valide gibbosis, dorso dense silaceo-pubescentibus, sutura basi, plaga antica transversa utrinque antice bi-postice uni-ramosa maculaque pone medium disci magna, sub-ovata, denudatis; margine subtilissime coriaceo. Long. 22½, lat. 17½ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Dom. *Virmont.* Mus. Berol.

3. Mesomphalia bituberculata.

Ovata, convexa, nigro-ænea, sub-opaca, subtus nitida, pube brevi silacea parce adspersa; prothorace lato, obsolete punctulato, antice late truncato, dein lateribus obliquis, basin versus sub-rectis; elytris basi retusis, valide obtuse gibbosis, vage, sub-remote punctatis, leviter reticulatis, basi plaga magna communi vittaque intramarginali silaceo-pubescentibus; singulo ante apicem tuberculo magno, obtuso, pubescente instructo; margine sat late explanato, remotius punctulato, antice impresso, deflexo. Long. 19½, lat. 15¼ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Dom. *Buquet.* Mus. Holm.

4. Mesomphalia difformis,

Bohem. Mon. Cassid. i. p. 221, 2.
Patr. Brasilia.

5. Mesomphalia congener.

Bohem. Mon. Cassid. i. p. 222, 3.
Cyrtonota congener, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

6. Mesomphalia gibbosa. B.M.

Bohem. Mon. Cassid. i. p. 223, 4, t. iii. F.
Cassida gibbosa, *Fabr.* Syst. El. i. p. 403, 89. Ent. Syst. i. p. 302, 49. Mant. Ins. i. p. 65, 40. Spec. Ins. i. p. 112, 33. *Linn.* Syst. Nat. ed. Gmel. i. iv. p. 1638, 48. *Oliv.* Ent. vi. p. 928, 5, 97, t. i. f. 6. *Herbst.* Naturs. viii. p. 346, 114. *Schœnh.* Syn. Ins. ii. p. 225, 116.
Cyrtonota gibbosa, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

7. Mesomphalia latipennis.

Breviter rotundata, convexa, nigra, sub-opaca, pube dilute silacea dense obsita; prothorace vix punctato, utrinque depresso, densius pubescente; elytris mediocriter, sub-remote punctulatis, undique foveolis profundioribus, denudatis, dorso sub-seriatis insculptis, basi retusis, antice valide conico-gibbosis; margine late explanato, medio valde rotundato-ampliato. Long. 18½, lat. 18½ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Dom. *Olfers.* Mus. Berol.

8. Mesomphalia tumidula.

Bohem. Mon. Cassid. i. p. 224, 5.
Patr. Brasilia.

9. Mesomphalia turrita. B.M.

Bohem. Mon. Cassid. i. p. 225, 6.
Cassida turrita, *Illig.* Mag. i. p. 180, 12. *Schœnh.* Syn. Ins. ii. p. 230, 157.
Cyrtonota vestita, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

10. Mesomphalia sexmaculosa.

Rotundata, convexa, nigro-ænea, parum nitida, parce silaceo-pubescens; prothorace subtilissime punctulato, medio canaliculato, utrinque macula magna, semilunata, silaceo-pubescente; elytris sat crebre; mediocriter punctatis, dorso discoque confertim leviter elevato-reticulatis, antice retusis, valide sub-conico gibbosis, maculis quatuor magnis, dense silaceo pubescentibus; margine late explanato, profundius,

crebrius punctato, pone basin deflexo; humeris rotundatis. Long. 18¼, lat. 17 millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Berol.

11. MESOMPHALIA ALBOFASCICULATA.

Rotundata, convexa, nigro-ænea, opaca; prothorace lato, subtilissime, remote punctulato, supra utrinque parce pallido-pubescente; elytris sub-remote, postice obsoletius punctulatis, pone basin valide gibbosis; singulo fasciculis septem parvis e pilis albidis ornato pubeque rara, prope marginem et postice ad suturam evidentiore, adsperso; margine ante medium latissime explanato, deflexo, antice leviter impresso. Long. 19, lat. 18¼ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Dom. *Deyrolle.*

12. MESOMPHALIA DENUDATA. B.M.

Bohem. Mon. Cassid. i. p. 227, 7.
Cyrtonota denudata, *Dej.* Cat. ed. 3. p. 368.
Patr. Brasilia.

13. MESOMPHALIA VARIOLARIS. B.M.

Bohem. Mon. Cassid. i. p. 228, 8.
Cyrtonota variolaris, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

14. MESOMPHALIA AMPLIATA.

Bohem. Mon. Cassid. i. p. 229, 9.
Patr. Bahia Brasiliæ.

15. MESOMPHALIA PYRAMIDATA.

Bohem. Mon. Cassid. i. p. 230, 10.
Patr. Columbia.

16. MESOMPHALIA SCROBICULATA.

Bohem. Mon. Cassid. i. p. 231, 11.
Patr. Brasilia.

17. MESOMPHALIA SEXMACULATA.

Bohem. Mon. Cassid. i. p. 232, 12.
Patr. Brasilia.

18. Mesomphalia sublævis.

Bohem. Mon. Cassid. i. p. 233, 13.
Patr. Rio Janeiro, Brasiliæ.

19. Mesomphalia nudicollis. B.M.

Bohem. Mon. Cassid. i. p. 234, 14.
Cyrtonota nudicollis, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

20. Mesomphalia lacunosa.

Bohem. Mon. Cassid. i. p. 235, 15.
Patr. Brasilia.

21. Mesomphalia fasciculata.

Bohem. Mon. Cassid. i. p. 236, 16.
Patr. Brasilia.

22. Mesomphalia impudens.

Rotundata, convexa, nigro-ænea, sub-opaca, parcissime, breviter pallido-pubescens; prothorace vix punctulato, medio distincte canaliculato, ante basin utrinque bi-impresso; elytris parum profunde, minus crebre punctatis, antice retusis, dein valide, obtuse gibbosis, dorso margineque interiore leviter, tenuiter reticulatis, reticulo sub-nitido, confertim punctato, areolas inæquales includente; singulo elytro macula elongata ad scutellum, altera ante apicem sub triangulari annuloque disci exterioris ante medium foveolam profundam cingente, silaceo-pubescentibus; humeris late rotundatis. Long. 16½, lat. 16 millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Holm. et Mus. Berol.

23. Mesomphalia vetula.

Bohem. Mon. Cassid. i. p. 237, 17.
Patr. Bahia, Brasiliæ.

24. Mesomphalia glabricollis.

Bohem. Mon. Cassid. i. p. 239, 18.
Patr. Bahia.

25. Mesomphalia retipennis.

Bohem. Mon. Cassid. i. p. 240. 19.
Patr. Brasilia.

26. Mesomphalia fossulata.

Bohem. Mon. Cassid. i. p. 241, 20.
Patr. Brasilia.

27. Mesomphalia vidua.

Bohem. Mon. Cassid. i. p. 242, 21.
Patr. Buenos Ayres.

28. Mesomphalia diversa.

Bohem. Mon. Cassid. i. p. 243, 22.
Cyrtonota diversa, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

29. Mesomphalia Godetii. B.M.

Bohem. Mon. Cassid. i. p. 244, 23.
Cyrtonota Godetii, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

30. Mesomphalia Stevensi.

Rotundata, convexa, nigro-ænea, sub nitida; prothorace subtilissime punctulato, tenuiter canaliculato, utrinque impresso, macula mediocri, sub-rotunda e pube densa silacea; elytris confertim, vage punctatis, antice retusis, dorso valide obtuse gibbosis, maculis sex magnis silaceo-pubescentibus ornatis; margine late explanato, posterius subtilius punctulato; humeris late rotundatis. Long. 18½, lat. 16 millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Dom. *Jekel*, Mus. Holm.

31. Mesomphalia deplorabunda. B.M.

Ovata, convexa, nigra, opaca, parce tenuiter pallido-pubescens; prothorace remote, parum profunde punctato, medio canaliculato, lateribus longe ultra medium obliquis, dein leviter rotundatis; elytris sat crebre mediocriter punctatis pone scutellum valide, obtuse gibbosis, disco parce, obsolete ferrugineo-rugulosis; margine explanato, confertim rugoso-punctato. Long. 13, lat. 11 millim.

Bohem. Mon. Cassid. Append.
Patr. Bolivia (Coll. *Bridges*), Mus. Britann.

32. Mesomphalia lenis.

Bohem. Mon. Cassid. i. p. 346, 24.
Patr. Brasilia.

33. Mesomphalia conspersa. B.M.
Bohem. Mon. Cassid. i. p. 247, 25.
Cassida conspersa, *Germ.* Ins. Spec. i. p. 534, 721.
Cyrtonota stigmata, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

34. Mesomphalia chalybea.
Bohem. Mon. Cassid. i. p. 248, 26.
Cassida chalybea, *Germ.* Ins. Spec. i. p. 532, 719.
Cyrtonota decorata, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

35. Mesomphalia vulnerata.
Bohem. Mon. Cassid. i. p. 249, 27.
Cyrtonota vulnerata, *Dej.* Cat. ed 3, p. 368.
Var. Cyrtonota subpustulata, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

II. *Elytra ante-medium dorsi leviter gibbosa.*

36. Mesomphalia flavosparsa. B.M.
Rotundata (*mas*) aut ovata (*femina*), convexa, supra obscure nigro-ænea, opaca, subtus nigra, sub-nitida; prothorace subtilissime, sub-remote punctulato, apice sub-truncato, lateribus ultra medium parum obliquis, dein rectis; elytris antice leviter gibbosis, subtiliter sub-remote punctulatis obsolete elevato-reticulatis, disco maculis parvis, sparsis inæqualibus, elevatis, lævibus, dilute flavescentibus; humeris late rotundatis. Long. *maris* 17½, lat. 15¼. Long. *fem.* 20, lat. 17 millim.
Bohem. Mon. Cassid. Append.
Var. a. elytrorum maculis ferrugineis.
Patr. Bolivia (Coll. *Bridges*), Mus. Britann. *Var. a.* Mus. Dom. *Chevrolat.*

37. Mesomphalia Sommeri.
Bohem. Mon. Cassid. i. p. 251, 28.
Patr. Brasilia.

38. Mesomphalia sexstillata.
Sub-ovata, convexa, supra viridi-ænea, sub-opaca, subtus cærulea, nitida; prothorace sat lato, subtilissime remote punctulato, lateribus ultra medium leviter, dein subito rotundato; elytris subtiliter, remote punctulatis, obsolete sub-reticulatis, maculis sex mediocribus fulvo-pubescentibus ornatis, dorso leviter, obtuse gibbosis; margine deflexo; humeris rotundatis. Long. 12¼, lat 9¾ millim.
Bohem. Mon. Cassid. Append.
Patr. Brasilia, Dom. *Deyrolle.* Mus. Holm.

39. Mesomphalia anceps.

Bohem. Mon. Cassid. i. p. 252, 29.
Cyrtonota anceps, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

40. Mesomphalia areolata.

Bohem. Mon. Cassid. i. p. 253, 30.
Cassida areolata, *Germ.* Ins. Spec. i. p. 533, 720.
Cyrtonota blanda, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

41. Mesomphalia sexsignata.

Bohem. Mon. Cassid. i. p. 254, 31.
Cyrtonata sexsignata, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

42. Mesomphalia impluviata.

Bohem. Mon. Cassid. i. p. 255, 32.
Patr. Brasilia.

43. Mesomphalia inornata.

Bohem. Mon. Cassid. i. p. 257, 33.
Patr. St. Catharina, Brasiliæ.

44. Mesomphalia lucida.

Rotundata, convexa, viridi-cyanea, parum nitida; prothorace lato, subtiliter, sub-remote punctulato, antice leviter rotundato-emarginato, lateribus ultra medium obliquis, dein rectis, dorso tenuiter, canaliculato; elytris subtiliter, sub-remote punctatis, disco obsolete foveolatis, pone basin obtuse gibbosis, gibbo nitido, cyaneo-micante; singulo maculis duabus parce lutescenti-pubescentibus, una basali, altera longo ante medium in disco exteriore. Long. 14, lat. 11 millim.

Bohem. Mon. Cassid. Append.
Patr. St. Catharina Brasiliæ, Mus. Dom. *Chevrolat.*

45. Mesomphalia consobrina.

Bohem. Mon. Cassid. i. p. 258, 34.
Patr. Brasilia.

46. Mesomphalia oblita.

Bohem. Mon. Cassid. i. p. 259, 35.
Cyrtonota oblita, *Dej.* Cat. ed. 3. p. 368.
Patr. Brasilia.

47. Mesomphalia implexa.

Bohem. Mon. Cassid. i. p. 260, 36.
Cyrtonota implexa, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

48. Mesomphalia subreticulata.

Bohem. Mon. Cassid. i. p. 261, 37.
Patr. Brasilia.

49. Mesomphalia subnotata.

Bohem. Mon. Cassid. i. p. 262, 38.
Patr. Brasilia.

50 Mesomphalia sexpustulata. B.M.

Bohem. Mon. Cassid. i. p. 263, 39.
Cassida sexpustulata, *Fabr.* Syst. El. i. p. 408, 116. Ent Syst. i. p. 306, 67. Mant. Ins. i. p. 66, 55. Spec. Ins. i. p. 114, 46. *Linn.* Syst. Nat. Gmel. i. iv. p. 1643, 65. *Herbst.* Naturs. viii. p. 348, 118. *Oliv.* Ent. vi. p. 939. 25, 97, t. iii. f. 36. Encycl. Ins. v. p. 392, 75. *Schœnh.* Syn. Ins. ii. p. 229. 148.
Cyrtonota sexpustulata, *Dej.* Cat. ed. 3, p. 368.
Cassida pavonina, *Herbst.* Naturs. viii. p. 322, 85. t. cxxxvi. f. 2. *Schœnh.* Syn. Ins. ii. p. 229, 147.
Var. e. supra nigro-ænea, maculis elytrorum quatuor anterioribus parvis, punctiformibus, duabus posticis deficientibus.
Patr. Brasilia, *Var, e.* St. Catharina, Mus. Dom. *Deyrolle.*

51. Mesomphalia similata.

Bohem. Mon. Cassid. i. p. 265, 40.
Var. a. elytrorum macula basali deficiente.
Patr. Brasilia Bahia.

52. Mesomphalia assimilis. B.M.

Bohem. Mon. Cassid. i. p. 266, 41.
Patr. Brasilia.

53. Mesomphalia ignita. B.M.

Bohem. Mon. Cassid. i. p. 267, 42.
Patr. St. Catharina, Brasiliæ.

54. Mesomphalia illepida.

Sub-rotundata, convexa, supra cyanea, nitida, subtus viridi-ænea; prothorace subtiliter, sub-remote punctulato, longe ultra medium obliquo, dein citius rotundato; elytris basi

retusis leviter gibbosis, distincte, minus crebre punctatis, foveolis numerosis, parum profundis, opacis obsoletius, remotius punctatis, gibbo viridi, nitido; margine sat late explanato, oblique deflexo. Long. 13½, lat. 11¼, millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Dom. *Deyrolle.*

55. Mesomphalia tumulus. B.M.

Bohem. Mon. Cassid. i. p. 268, 43.
Patr. St. Catherina, Brasiliæ.

56. Mesomphalia Silbermanni.

Bohem. Mon. Cassid. i. p. 269, 44.
Patr. Buenos Ayres.

57 Mesomphalia chalcoptera.

Bohem. Mon. Cassid. i. p. 270, 45.
Patr. Columbia.

58. Mesomphalia flavo-reticulata. B.M.

Sub-triangularis, convexa, nigro-ænea, opaca; prothorace subtiliter punctulato, lateribus ultra medium obliquis, dein rectis; elytris basi sub-truncatis, antice retusis, obtuse gibbosis, subtilissime punctulatis, medio fascia lata transversa, communi, inæquali vittaque disci longitudinali cum fascia connexa, flavo-reticulatis. Long. 16, lat. 14 millim.

Bohem. Mon. Cassid. Append.
Patr. Bolivia (Coll. *Bridges*), Mus. Britann.

59. Mesomphalia textilis. B.M.

Bohem. Mon. Cassid. i. p. 271, 46.
Patr. Columbia.

60 Mesomphalia castigata. B.M.

Rotundata, convexa, nigro-ænea, supra opaca, parce breviter cinereo-pubescens, subtus nitida, glabra; prothorace lato, sub-remote punctulato, antice leviter emarginato, anguste testaceo-marginato, lateribus ultra medium obliquis, dein rectis; elytris confertim sanguineo-reticulatis, reticulo parum elevato, confertim punctato, areolas mediocres, remote punctatas includente, pone basin obtuse gibbosis; humeris subangulatis, antice sub-truncatis. Long. 14, lat. 12½ millim.

Bohem. Mon. Cassid. Append.
Patr. Bolivia, Mus. Brit. (Coll. *Bridges*) Dom. *Chevrolat.* Peruvia, Dom. *Deyrolle*, Mus. Holm.

61 MESOMPHALIA CAUDATA. B.M.

Bohem. Mon. Cassid. i. p. 272, 47.
Patr. Columbia (Col.. *Goudot.*)

62. MESOMPHALIA VERRUCOSA.

Sub-rotundata, convexa, supra cærulea, subtus nigra, nitida; prothorace subtilissime, remote punctulato, lateribus longe ultra medium obliquis, dein sub-rectis, dorso tenuiter canaliculato; elytris parum profunde, sat crebre, punctatis, basi obtuse gibbosis; singulo in disco inæqualiter, dilute sanguineo-rugoso; humeris non prominulis, leviter rotundatis; margine subtiliter punctulato. Long. 13½, lat. 11 millim.

Bohem. Mon. Cassid. Append.
Patr. Pernambuco, Dom. *Dohrn.* Mus. Reg. Holm.

63. MESOMPHALIA CRUENTATA.

Ovata, convexa, nigra, opaca, parce cinereo-pubescens, subtus nitida; prothorace mediocriter, minus crebre punctato, canaliculato, lateribus longe ultra medium leviter rotundatis; elytris pone basin retusis; valide, breviter gibbosis, mediocriter, minus crebre punctatis, rugis remotis sanguineis laxe, irregulariter reticulatis; margine modice explanato, oblique deflexo; humeris rotundatis. Long. 12½. lat. 9¾ millim.
Bohem. Mon. Cassid. i. p. 382, 142. *Bohem.* Mon. Cassid. Append.
Cyrtonota cruentata, *Erichs.* Archiv. f. Naturg. xiii. p. 152, 1.
Patr. Peruvia orientalis, Dom. *Tschudi.* Mus. Berol.

64. MESOMPHALIA PLEUROSTICHA. B.M.

Bohem. Mon. Cassid. i. p. 274. 48.
Cyrtonota pleurosticha, *Erichs.* Archiv. f. Naturg. xiii. p. 152, 4.
Patr. Peruvia orientalis, Bolivia (Coll. *Bridges*).

65. MESOMPHALIA FLOCCOSA.

Sub-triangularis, modice convexa, nigro-ænea, opaca, subtus nitida; prothorace subtiliter, minus crebre punctulato, antice utrinque anguste testaceo-marginato, medio canaliculato, utrinque bi-impresso; elytris cyaneo-micantibus, antice, leviter gibbosis, subtiliter, minus crebre punctulatis, leviter, remote reticulatis, parce griseo-pubescentibus, posterius passim fasciculatis, basi fere truncatis; margine late explanato antice impresso; humeris rotundatis. Long. 12. lat. 11 millim.

Bohem. Mon. Cassid. i. p. 383, 144. *Bohem.* Mon. Cassid. Append.
Cyrtonota floccosa, *Erichs.* Archiv. f. Naturg. xiii. p. 152, 3.
Patr. Peruvia orientalis, Dom. *Tschudi.* Mus. Berol.

66. Mesomphalia confusa.

Bohem. Mon. Cassid. i. p. 275, 49.
Cyrtonota confusa, *Dej.* Cat. ed. 3, p. 369.
Patr. Cayenna.

67. Mesomphalia rubicundula.

Breviter rotundata, modice convexa, nigro-cyaneo, opaca; antennis basi ferrugineis; prothorace subtilissime punctulato, antice anguste ferrugineo-marginato, medio canaliculato, lateribus ultra medium obliquis, dein leviter rotundatis; elytris obsolete punctulatis, tenuiter griseo-pubescentibus, tenuiter-reticulatis, reticulo sub-nitido, areolas minores includente; disco medio transversim margineque, basi et apice exceptis, sanguineis, dorso antico obtuse gibbosis; margine late explanato; humeris rotundatis. Long. 17, lat. 18 millim.

Bohem. Mon. Cassid. Append.
Patr. Peruvia, Mus. Dom. *Chevrolat.*

68. Mesomphalia ænea. B.M.

Bohem. Mon. Cassid. i. p. 276, 50.
Cassida ænea, *Oliv.* Entom. vi. p. 937, 20. t. iv. f. 61.
Cyrtonota ænea, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

69. Mesomphalia demissa.

Sub-triangularis, nigro-ænea, supra opaca; antennis articulis 2-4 subtus testaceis; prothorace angustiore, subtiliter, subremote punctulato, lateribus ultra medium obliquis, dein basin versus rectis; elytris tenuiter, remotius reticulatis, reticulo parum elevato, crebre punctato, parum nitido, areolas majores, læves includente; dorso antico evidentius gibbosis, medio valde dilatatis, apice acuminatis; margine late explanato, punctulato, medio intus reticulato, ante medium macula flava e reticulo elevato formata notato. Long. 15-17. lat. 13-15 millim.

Bohem. Mon. Cassid. Append.
Mesomphalia ænea, *Var. a. Bohem.* Mon. Cassid. i. p. 276, 50.
Patr. Bahia, Brasiliæ, Dom. *Schœnherr*, Mus. Holm.

70. Mesomphalia contubernalis.

Sub-triangularis, convexa, supra fusco-cuprea, sub-opaca; subtus obscure ænea, nitida; antennis articulis 2-4 subtus testaceis; prothorace angustiore, subtiliter, sub-remote, punctulato, lateribus ultra medium obliquis, dein basin versus rectis; elytris crassius, irregulariter reticulatis, reticulo parum nitido, areolas mediocres, læves includente, dorso antico leviter gibbosis, medio valde dilatatis, apice acuminatis; margine late explanato, ante medium plaga magna flava, extus multoties sinuato-emarginata, intus sub-truncata, maculis 3-4 parvis, fusco-cupreis notata. Long. 13-14½, lat. 11¾-12 millim.

Bohem. Mon. Cassid. Append.
Mesomphalia ænea, *Var. b. Bohem.* Mon. Cassid. i. p. 276, 50.
Patr. Bahia, Dom. *Dohrn*, et *Deyrolle* Mus. Holm.

71. Mesomphalia blanda.

Bohem. Mon. Cassid. i. p. 277, 51,
Patr. Columbia.

72. Mesomphalia acuminata.

Bohem. Mon. Cassid. i. p. 278, 52.
Cyrtonota acuminata, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

73. Mesomphalia Redtenbacheri. B.M.

Bohem. Mon. Cassid. i. p. 280, 53.
Patr. Brasilia (Rio Grande).

74. Mesomphalia fastuosa.

Sub-triangularis, modice convexa, viridis, opaca, subtus nigro-ænea, nitida; antennis articulis 1 apice, 2-4 subtus testaceis, prothorace mediocriter, sat crebre punctato, cupreo-tincto; elytris minus crebre, parum profunde punctulatis, apice acuminatis totis irregulariter reticulatis, reticulo tenui, nitido, antice areolas parvas, postice majores includente; margine late explanato, intus ante medium macula magna irregulari, flava, viridi-trimaculata notato, limbo cupreo; humeris antrorsum parum dilatatis, rotundatis. Long. 13½, lat. 12 millim.

Bohem. Mon. Cassid. i. p. 383, 146. *Bohem.* Mon. Cassid. Append.
Cyrtonota illustris, *Erichs.* Archiv. f. Naturg. xiii. p. 152, 6.
Patr. Peruvia orientalis, Dom. *Tschudi.* Mus. Berol.

75. Mesomphalia Mannerheimi.

Bohem. Mon. Cassid. i. p. 281, 54.
Patr. Chiquitos Boliviæ.

76. Mesomphalia inæqualis. B.M.

Bohem. Mon. Cassid. i. p. 282, 55.
Cassida inæqualis, *Fabr.* Syst. El. i. p. 406, 108. Ent. Syst. i. p. 305, 61. Mant. Ins. i. p. 65, 51. Spec. Ins. i. p. 114, 42. Syst. Ent. p. 93, 28. *Linn.* Syst. Nat. i. ii. p. 578, 24. Ed. Gmel. i. iv. p. 1642, 24. *Herbst.* Naturs. viii. p. 316, 79, t. cxxx. f. 4. *Oliv.* Ent. vi. p. 936, 18, 97, t. iii. f. 49, a. b. *Schulze.* in Naturs. ix. p. 103. t. ii. f. 6. *Schœnh.* Syn. Ins. ii. p. 227, 136.
Cassida lateralis, *Rœm.* Ins. p. 6, 15, t. iii. f. 2. *Sulz.* Gesch. d. Ins. t. iii. f. 2. a. Mus. Petropol. p. 650, 130.
Cassida bimaculata, *De Geer*, Mem. Ins. v. p. 182, 6. t. xv. f. 10. *Voet.* Coll. ii. t. xli. f. 2.
Cassida lunata, *Thunb.* Nov. Ins. Spec. v. p. 88. t. v. f. 99.
Cyrtonota inæqualis, *Dej.* Cat. ed. 3, p. 369.
Patr. Cayenna, Guyana. Surinam. Amazon (Para) Coll. *H. W. Bates.*

77. Mesomphalia elegans. B.M.

Bohem. Mon. Cassid. i. p. 284, 56.
Patr. Mexico.

78. Mesomphalia Chevrolati.

Bohem. Mon. Cassid. i. p. 285, 57.
Cyrtonota Chevrolati, *Dej.* Cat. ed. 3, p. 369.
Patr. Mexico.

79. Mesomphalia tristigma. B.M.

Bohem. Mon. Cassid. i. p. 286, 58.
Cyrtonota tristigma, *Dej.* Cat. ed. 3, p. 369.
Patr. Mexico. Honduras (Coll. *Dyson*).

80. Mesomphalia punicea. B.M.

Bohem. Mon. Cassid. i. p. 287, 59.
Cyrtonota punicea, *Dej.* Cat. ed. 3, p. 369.
Patr. Mexico (*F. Walker, Esq.*) Honduras (Coll. *Dyson*).

81. Mesomphalia primaria.

Bohem. Mon. Cassid. i. p. 289, 60.
Patr. Mexico.

82. Mesomphalia Lebasii. B.M.

Bohem. Mon. Cassid. i. p. 290, 61.
Cyrtonota Lebasii, *Dej.* Cat. ed. 3, p. 369.
Patr. Columbia (Coll. *Goudot*). *Var. Guatimala* (Coll. *Dr. Scherzer*).

83. Mesomphalia acuta.

Bohem. Mon. Cassid. i. p. 291, 62.
Patr. Brasilia.

84. Mesomphalia antiqua. B.M.

Bohem. Mon. Cassid. i. p. 292, 63.
Cassida antiqua, *Sablb.* Spec. Ins. i. p. 56, 37.
Cassida nodipennis, *Germ.* Ins. Spec. i. p. 532, 718.
Cyrtonota ænea, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia (Maj.-Gen. *Hardwicke*).

85. Mesomphalia modica.

Bohem. Mon. Cassid. i. p. 293, 64.
Patr. Brasilia.

86. Mesomphalia plumbea. B.M.

Bohem. Mon. Cassid. i. p. 294, 65.
Cassida plumbea, *Germ.* Ins. Spec. i. p. 531, 717.
Cyrtonota impustulata, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

87. Mesomphalia flavonotata.

Sub-ovata, convexa, supra obscure ænea, sub-opaca, subtus nigro-ænea, nitida; antennis articulis 2 et 3 subtas ferrugineis; prothorace subtiliter, remote punctulato, medio breviter canaliculato, lateribus vix ad medium obliquis, dein sub-rectis, apice utrinque anguste testaceo-marginato; elytris subtiliter, crebre, disco antico evidentius rugoso-punctatis, basi retusis, obtuse gibbosis; singulo maculis duabus parvis, flavis; margine late explanato, subtilius punctulato. Long. 17, lat. 14 millim.

Bohem. Mon. Cassid. Append.
Var. a. macula postica elytrorum obsoleta, e punctis duabus parvis flavis formata.
Patr. Pernambuco, Mus. Dom. *Dohrn.*

88. Mesomphalia flavoguttata. B.M.

Bohem. Mon. Cassid. i. p. 295, 66.
Cyrtonota flavoguttata, *Dej.* Cat. ed. 3, p. 569.
Patr. Brasilia.

89. Mesomphalia verecunda.

Bohem. Mon. Cassid. i. p. 297, 67.
Patr. Corrientes.

90. Mesomphalia opaca.

Ovata, convexa, nigra, opaca; prothorace obsolete punctulato, antice leviter rotundato-emarginato, utrinque anguste testaceo-marginato, lateribus rotundato; elytris subtiliter, vage, sat crebre punctulatis, postice maculis nonnullis parvis irregularibus, sub-elevatis, sordide ochraceis insignitis, antice ad suturam impressis, dein leviter gibbosis; humeris rotundatis, Long. 16½, lat. 13 millim.

Bohem. Mon. Cassid. Append.
Patr. Bolivia, Mus. Dom. *Chevrolat.*

91. Mesomphalia sexplagiata.

Bohem. Mon. Cassid. i. p. 298, 68.
Patr. Brasilia.

92. Mesomphalia quadriplagiata.

Bohem. Mon. Cassid. i. p. 299, 69.
Var. a. elytrorum maculis dilute flavis.
Patr. Brasilia.

93. Mesomphalia smaragdina. B.M.

Sub-triangularis, convexa, supra smaragdina, opaca, subtus nigra, nitida; undique anguste cupreo-marginata; antennis articulo secundo subtus testaceo; prothorace subtiliter, crebre punctulato, medio cupreo-micante, tenuiter canaliculato; elytris antice retusis, leviter gibbosis, basi truncatis, subtiliter coriaceis, confertim punctulatis, gibbo vittaque disci valde inæquali, cupreis, magis nitidis. Long. 10, lat. 8¾ millim.

Bohem. Mon. Cassid. Append.
Var. a. dorso postico macula parva flavo-testacea.
Patr. Bolivia, Mus. Britann. (Coll. *Bridges*).

94. Mesomphalia Goryi. B.M.

Bohem. Mon. Cassid. i. p. 300, 70.
Patr. Columbia (Coll. *Goudot*).

95. Mesomphalia lateralis. B.M.

Bohem. Mon. Cassid. i. p. 301, 71.
Cassida lateralis, *Fabr.* Syst. El. i. p. 407, 109. Ent. Syst. i.

p. 305, 62. Mant. Ins. i. p. 66, 52. Spec. Ins. i. p. 114, 43. Syst. Entom. p. 93, 29. *Linn.* Syst. Nat. i. ii. p. 578, 26. Ed. Gmel. i. iv. p. 1642, 26. *Herbst.* Naturs, viii. p. 315, 78, t. cxxx. f. 5. *De Geer.* Mem. Ins. v. p. 183, 8, t. xx. f. 12. *Voet.* Cat. ii. t. xliii. f. 24. *Oliv.* Ent. vi. p. 937, 21, 97, t. iv. f. 62. Encycl. meth. v. p. 351, 72. *Schœnh.* Syn. Ins. ii. p. 227, 137.

Cassida ferruginea, *Voet.* Cat. ii. t. xliii. f. 24.

Cyrtonota lateralis, *Dej.* Cat. ed. 3, p. 368.

Patr. Cayenna, Guyana.

96. Mesomphalia bistigma. B.M.

Sub-rotundata, modice convexa, supra smaragdina, parum nitida, subtus nigro-ænea, nitida; antennis articulis 2-4 subtus testaceis; prothorace subtilissime, crebre punctulato, medio cupreo-micante, lateribus basin versus rotundato; elytris subtiliter coriaceis, sat crebre, parum profunde punctatis, dorso antico obsolete gibbosis; margine late explanato, macula magna elongata, dilute flava, profundius punctata notato. Long. 8½, lat. 7½ millim.

Bohem. Mon. Cassid. Append.

Patr. Bolivia, Mus. Britann. (Coll. *Bridges*).

97. Mesomphalia Serinus.

Sub-triangularis, modice convexa, dilute viridi-ænea, nitidula; antennis articulis 4 basalibus testaceis, prothorace subtiliter, crebre punctulato; elytris crebre mediocriter punctatis, basi subtruncatis, pone basin sub-gibbosis, disco maculis duabus longitudinaliter positis plagaque magna marginali flavis; humeris fere rectangulis, apice leviter rotundatis. Long. 9½-10, lat. 7¾ millim.

Bohem. Mon. Cassid. i. p. 383, 147. *Bohem.* Mon. Cassid. Append.

Cyrtonota Serinus, *Erichs.* Arch. f. Naturg. xiii. p. 153, 7.

Patr. Peruvia orientalis, Mus. Berol. Prope fluvium Amazonum, Dom. *Jekel.* Mus. Holm.

98. Mesomphalia Jekeli. B.M.

Sub-rotundata, modice convexa, fusco-cuprea, parum nitida, subtus nigro-ænea, nitida; antennis articulis basalibus subtus testaceis; prothorace subtilissime, sat crebre punctulato, lateribus basin versus rotundato; elytris subtiliter coriaceis, crebre mediocriter punctulatis, dorso antico obtuse, leviter gibbosis; margine late explanato, medio macula brevi, dilute flava, profundius punctata notato. Long, 9-10, lat. 7½-7¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Prope fluvium Amazonum, Dom. *Jekel.* Mus. Holm. Mus. Britann.

99. MESOMPHALIA DIMIDIATA. B.M.

Bohem. Mon. Cassid. i. p. 302, 72.
Patr. Brasilia (Rio Grande).

100. MESOMPHALIA BIPUSTULATA. B.M.

Bohem. Mon. Cassid. i. p. 305, 73.
Cassida bipustulata, *Fabr.* Syst. El. i. p. 407, 113. Ent. Syst. i. p. 306, 65. Mant. Ins. i. p. 66, 54. Spec. Ins. i. p. 114, 45. Syst. Ent. p. 93, 31. *Linn.* Syst. Nat. i. ii. p. 578, 38. Amœn. Acad. vi. p. 392, 8. Ed. Gmel. i. iv. p. 1643, 30. Cent. Ins. No. 8. *Schulz.* Naturs. vi. p. 91, t. iv. f. 2. *Illig.* Mag. i. p. 409, 113. *Oliv.* Ent. vi. p. 938, 23, 97, t. iii. f. 35. Encycl. meth. v. p. 391, 74. *Schœnh.* Syn. Ins. ii. p. 228, 141.
Cassida tetrastigma, *Herbst.* Naturs. viii. p. 318, 80, t. cxxxv. f. 10. *Schœnh.* Syn. Ins. ii. p. 228, 141.
Cyrtonota bipustulata, *Dej.* Cat. ed. 3, p. 369.
Cassida quadripunctata, *Voet.* Cat. ii. t. xlii. f. 6.
Var. Cassida cuprea, *Fabr.* Syst. El. i. p. 408, 115. Ent. Syst. i. p. 306. Mant. Ins. App. p. 308, 45, 46. *Linn.* Syst. Nat. Gmel. i. iv. p. 1643, 36. *Herbst.* Naturs. viii. p. 321, 83, t. cxxxvi. f. 1. *Schœnh.* Syn. Ins. ii. p. 228, 144.
Var. Cassida discors, *Fabr.* Syst. El. i. p. 408, 114. *Schœnh.* Syn. Ins. ii. p. 228, 143.
Cassida phlyctæna, Cat. Hamb. p. 40, 492.
Var. Cassida duodecimpustulata, *Fabr.* Syst. El. i. p. 408, 117. Suppl. p. 85, 67, 68. *Oliv.* Enc. meth. v. p. 393, 34. Entom. vi. p. 940, 27, 97, t. iii. f. 39 (mala). *Herbst.* Naturs. viii. p. 354, 128. *Schœnh.* Syn. Ins. ii. p. 229, 149.
Var. l. supra cærulea, immaculata.
Patr. Cayenna, Guyana.

101. MESOMPHALIA BIFENESTRATA. B.M.

Rotundata, modice convexa, viridi-cærulescens, sub-opaca. subtus nigra, magis nitida; antennis articulis 2-4 subtus testaceis; prothorace subtiliter, minus crebre punctulato, margine apicali utrinque anguste testaceo, lateribus late rotundato-ampliatis; elytris dorso rugoso-punctatis, declivitate posteriore margineque subtiliter punctulatis, pone basin vix gibbosis; singulo ante medium macula intramarginali, mediocri, flava. Long. 10-12¾, lat. 9-11 millim.

Bohem. Mon. Cassid. Append.
Patr. Bolivia, Mus. Britann. (Coll. *Bridges*).

102. Mesomphalia bioculata.

Bohem. Mon. Cassid. i. p. 305, 74.
Patr. Brasilia.

103. Mesomphalia discoides. B.M.

Bohem. Mon. Cassid. i. p. 306, 75.
Cassida discoides, *Linn.* Syst. Nat. i. ii. p. 578, 27. Ed. Gmel. i. iv. p. 1642, 27. *Herbst.* Naturs. viii. p. 319, 81, t. cxxxv. f. 11.
Cassida discoides, *Fabr.* Syst. El. i. p. 407, 112. Ent. Syst. i. p. 306, 64. Mant. Ins. i. p. 66, 53. Spec. Ins. i. p. 114, 44. Syst. Ent. p. 93, 30. *Oliv.* Ent. vi. p. 936, 19, 97, t. ii. f. 18, a, b. Encycl. meth. Ins. v. p. 391, 73. *Schœnh.* Syn. Ins. ii. p. 228, 140.
Cassida quadrimaculata, *De Geer*, Mem. d' Ins. v. p. 183, t. xv. f. 11. *Voet.* Col. ii. t. xli. f. 1.
Cassida Paramaribœnsis, *Voet.* Col. ii. t. xlii. f. 12.
Cyrtonota bipustulata, *var. Dej.* Cat. ed. 3, p. 369.
Patr. Cayenna, Bahia.

104. Mesomphalia funebris. B.M.

Bohem. Mon. Cassid. i. p. 308, 76.
Patr. Corrientes, Paraguay, Bolivia (Coll. *Bridges*).

105. Mesomphalia scoparia.

Ovata convexa, nigro-cærulea opaca, subtus nigra, nitida; prothorace sublævi, medio tenuiter canaliculato, apice utrinque anguste testaceo-marginato, pone medium utrinque impresso; elytris antice leviter gibbosis, subtiliter subremote punctulatis, obsolete reticulatis, parce subtiliterque silaceo-pubescentibus, passim fasciculatis; margine subtilius punctulato; humeris rotundatis. Long. 17½, lat. 14½ millim.

Bohem. Mon. Cassid. i. p. 382, 143. *Bohem.* Mon. Cassid. Append.
Cyrtonota scoparia, *Erichs.* Arch. f. Naturg. xiii. p. 152, 2.
Patr. Peruvia orientalis, Dom. *Tschudi.* Mus. Berol.

106. Mesomphalia tomentosa. B.M.

Bohem. Mon. Cassid. i. p. 309, 77.
Patr. Columbia. Venezuela (Coll. *Dyson*).

107. Mesomphalia puberula.

Rotundata, modice convexa, nigro-ænea, sub-opaca, pube brevissima depressa, pallida parce, in prothorace densius adspersa; antennis articulis tribus primis subtus testaceis;

prothorace angusto, subtiliter, sub-remote punctulato, lateribus ad medium leviter rotundatis, dein sub-rectis; elytris obsolete punctatis, dorso leviter irregulariter reticulatis; humeris non prominulis, apice leviter rotundatis. Long. 16½, lat. 14½ millim.

Bohem. Mon. Cassid. Append.
Patr. Campo del Sacramento, Mus. Paris.

108. Mesomphalia excisa.

Rotundata, convexa, supra nigro-cærulea, opaca, parce albido-puberula, subtus nigra, nitida; prothorace longitudine vix dimidio latiori, obsolete punctulato, utrinque profunde arcuatim impresso, lateribus ad medium obliquis, dein basin versus leviter rotundato-angustatis; elytris subtiliter, subremote, posterius adhuc subtilius punctulatis, basi retusis, impressis, dein valide, obtuse gibbosis; margine late explanato, obsolete punctulato, pone basin profunde impresso, intus foveolato; humeris antrorsum prominulis, rotundatis. Long. 15½, lat. 14 millim.

Bohem. Mon. Cassid. Append.
Patr. Puerto Caballo, Mus. Berol. Columbia, Mus. Dom. *Dohrn.*

III. *Elytra convexa, haud gibbosa.*

109. Mesomphalia Ephippium.

Bohem. Mon. Cassid. i. p. 310, 78.
Cassida Ephippium, *Herbst.* Naturs. viii. p. 310, 74, t. cxxxv. f. 6. *Schœnh.* Syn. Ins. ii. p. 230, 136.
Patr. America borealis.

110. Mesomphalia plagiata.

Bohem. Mon. Cassid. i. p. 319, 79.
Var. a. maculis plagis elytrorum æneis majoribus, marginali reliquis majore, transversa.
Patr. Remedios, Columbiæ. *Var. a.* Surinam, Mus. Dom. *Deyrolle.*

111. Mesomphalia flavomaculata.

Bohem. Mon. Cassid. i. p. 313, 80.
Cassida flavomaculata, *Fabr.* Syst. El. i. p. 406, 106. *Schœnh.* Syn. Ins. ii. p. 227, 134.
Patr. Guyana.

112. Mesomphalia Mellyi.

Bohem. Mon. Cassid. i. p. 314, 81.
Patr. Mexico.

113. Mesomphalia Saundersi.

Rotundata, minus convexa, nigro-ænea, opaca, subtus nitida, pilis cinereis parce adspersa; prothorace inæquali, subtilissime, remote punctulato, medio tenuiter canaliculato; elytris ad scutellum impressis, dein leviter gibbosis, dorso foveis magnis, lævibus impressis, interstitiis angustis, elevatis, nitidis, punctulatis; margine late explanato, dilute flavo, crebre punctulato, intus lineolis tribus brevibus, transversis, nigro-æneis plagæ dorsali adfixis, limbo toto anguste nigro-æneo. Long. 18½, lat 17½ millim.

Bohem. Mon. Cassid. Append.
Patr. Quito, Dom. *Buquet.* Mus. Holm.

114. Mesomphalia elocata. B.M.

Bohem. Mon. Cassid. i. p. 315, 82.
Patr. Columbia (Ibaque), Coll. *Goudot.*

115. Mesomphalia dissecta.

Bohem. Mon. Cassid. i. p. 316, 83.
Patr. Bogota.

116. Mesomphalia reticularis.

Bohem. Mon. Cassid. i. p. 317, 84.
Cassida reticularis, *Fabr.* Syst. El. i. p. 404, 94. Ent. Syst. i. p. 303, 53. Mant. Ins. i. p. 65, 44. Spec. Ins. i. p. 113, 37. Syst. Ent. p. 92, 23. *Linn.* Syst. Nat. i. ii. p. 576, 15. Ed. Gmel. i. iv. p. 1640, 15. *Voet.* Col. ii. t. xli. f. 4. *Herbst.* Naturs. viii. p. 295, 63, t. cxxx. f. 3. *Oliv.* Encycl. Ins. v. p. 389, 60. Entom. vi. p. 927, 3, 97, t. iv. f. 51. *Schœnh.* Syn. Ins. ii. p. 223, 120.
Cassida punctata, *De Geer.* Mem. Ins. v. p. 180, 4, t. xv. f. 8.
Cassida ornata, *Herbst.* Archiv. p. 50, 8, t. xxii. f. 30.
Cyrtonota reticularis, *Dej.* Cat. ed 3, p. 369.
Var. Cassida retiformis, *Fabr.* Syst. El. i. p. 405, 98. Ent. Syst. i. p. 304, 56. Mant. Ins. ii. App. p. 380, 38, 39. *Herbst.* Naturs. viii. p. 347, 116. *Linn.* Syst. Nat. Gmel. i. iv. p. 1643, 34. *Oliv.* Ent. vi. p. 928, 4, 97, t. iv. f. 52. Encycl. meth. v. p. 387, 51. *Schœnh.* Syn. Ins. ii. p. 226, 124.
Var. Cassida annulata, *Fabr.* Syst. El. i. p. 405, 102. Ent. Syst. i. p. 304, 57. Mant. Ins. p. 65. 47. Spec. Ins. i. p. 113, 39. *Herbst.* Naturs. viii. p. 291, 60, t. cxxxiv. f. 5.

Linn. Syst. Nat. Gmel. i. iv. p. 1641, 61. Naturf. ix. t. ii. f. 6. *Schœnh.* Syn. Ins. ii. p. 226, 129.
Var. Cassida hieroglyphica, *Herbst.* Naturs. viii. p. 296, 64, t. cxxxiv. f. 8. *Oliv.* Ent. vi. p. 942, 30, 97, t. ii. f. 27. *Schœnh.* Syn. Ins. ii. p. 230, 164.
Patr. Cayenna.

117. Mesomphalia jucunda.

Bohem. Mon. Cassid. i. p. 319, 85.
Cyrtonota jucunda, *Dej.* Cat. ed. 3, p. 369.
Patr. Cayenna.

118. Mesomphalia rubroreticulata.

Sub-triangularis, modice convexa, ænea, supra sub-opaca, subtus nitida; prothorace minus crebre, parum profunde punctato, medio tenuiter canaliculato, lateribus ad medium obliquis, dein rectis; elytris pone basin vix gibbosis, limbo excepto undique rubro-reticulatis, reticulo parum elevato, punctato, margine crassiore areolas sat magnas, æneas includente. Long. 15, lat. 13½ millim.

Bohem. Mon. Cassid. Append.
Patr. Bogota, Dom. *Deyrolle*, Mus. Holm.

119. Mesomphalia illustris. B.M.

Bohem. Mon. Cassid. i. p. 320, 86.
Cassida illustris, *Chevrol.* Col. de Mexique, f. iii.
Cyrtonota illustris, *Dej.* Cat. ed. 3, p. 369.
Patr. Mexico, *Guatemala.*

120. Mesomphalia decemguttata. B.M.

Bohem. Mon. Cassid. i. p. 321, 87.
Cyrtonota decemguttata, *Dej.* Cat. ed. 3, p. 369.
Patr. Brasilia, Amazon (Para).

121. Mesomphalia pallidoguttata. B.M.

Bohem. Mon. Cassid. i. p. 322, 88.
Cyrtonota pallidoguttata, *Blanch.* Voy. d'Orbigny, pl. xxiv. f. 4.
Patr. Andes Americæ meridionalis, Bolivia (Coll. *Bridges*).

122. Mesomphalia lata.

Bohem. Mon. Cassid. i. p. 323, 89.
Patr. Bolivia.

123. Mesomphalia cyanea. B.M.

Bohem. Mon. Cassid. i. p. 324, 90.
Cassida cyanea, *Linn.* Syst. Nat. i. ii. p. 577, 22. Mus. Ludov. Ulr. p.39. Ed. Gmel. i. iv. p. 1642, 22. *Herbst.* Naturs. viii. p. 344, 112. *Petiv.* Gaz. t. 57, f. 6? *Schœnh.* Syn. Ins. ii. p. 224, 111.
Cyrtonota cyanescens, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

124. Mesomphalia thalassina. B.M.

Bohem. Mon. Cassid. i. p. 326, 91.
Patr. Brasilia.

125. Mesomphalia indigacea. B.M.

Bohem. Mon. Cassid. i. p. 327, 92.
Patr. Bolivia (Coll. *Bridges*).

126. Mesomphalia festiva. B.M.

Bohem. Mon. Cassid. i. p. 328, 93.
Cassida festiva, *Klug*, Preis-Verzeichn. 1829, p. 7, 86.
Cyrtonota festiva, *Dej.* Cat. ed. 3, p. 368.
Patr. Brasilia.

127. Mesomphalia aureomarginata.

Ovata, convexa, nitida, subtus cærulea; prothorace cupreo-æneo, parum nitido, sat profunde, irregulariter punctato, medio canaliculato, lateribus ultra medium obliquo, dein subito rotundato; elytris smaragdinis, sub-opacis, irregulariter cupreo-reticulatis, reticulo elevato, tenui, nitido, lævi, areolas sat magnas, remote punctatas includente; margine modice explanato, deflexo, aureo-cyaneo micante, obsoletius reticulato; humeris rotundatis. Long. 14, lat. 9½ millim.

Bohem. Mon. Cassid. Append.
Patr. Montevideo, Mus. Dom. *Deyrolle.*

128. Mesomphalia vicina.

Bohem. Mon. Cassid. i. p. 329, 94.
Patr. Rio Grande, Brasiliæ.

129. Mesomphalia cærulescens. B.M.

Bohem. Mon. Cassid. i. p. 330, 95.
Patr. Bolivia (Coll. *Bridges*).

130. Mesomphalia gentilis.

Sub-triangularis, modice convexa, nigro-ænea, supra subnitida, glabra; prothorace subtilissime, crebre coriaceo, utrinque bi-impresso, lateribus ultra medium obliquo, dein valde rotundato; elytris non gibbosis, posterius fornicatis, irregulariter reticulatis, reticulo parum elevato, punctato, areolas mediocres opacas, obsolete punctatas includente; margine late explanato, subtiliter, crebre punctulato, parce fulvo-liturato; humeris antrorsum prominulis, obtusis. Long. 14½, lat. 13 millim.

Bohem. Mon. Cassid. i. p. 383, 145. *Bohem.* Mon. Cassid. Append.
Cyrtonota gentilis, *Erichs.* Archiv. f. Naturg. xiii. p. 152, 5.
Patr. Peruvia orientalis, Mus. Berol.

131. Mesomphalia Kollari.

Bohem. Mon. Cassid. i. p. 331, 96.
Patr. Brasilia.

132. Mesomphalia Schaumi.

Bohem. Mon. Cassid. i. p. 332, 97.
Patr. Brasilia.

133. Mesomphalia lineatocollis.. B.M.

Bohem. Mon. Cassid. i. p. 333, 38.
Cyrtonota lineatocollis, *Dej.* Cat. ed. 3. p. 369.
Patr. Brasilia.

134. Mesomphalia brevicollis.

Bohem. Mon. Cassid. i. p. 335, 99.
Patr. Bolivia.

135. Mesomphalia extricata.

Bohem. Mon. Cassid. i. p. 336, 100.
Patr. Bolivia.

136. Mesomphalia hæmatina.

Bohem. Mon. Cassid. i. p. 337, 101.
Patr. Chiquitos.

137. Mesomphalia quatuordecimsignata. B.M.

Bohem. Mon. Cassid. i. p. 338, 102.
Patr. Bolivia (Coll. *Bridges*).

138. MESOMPHALIA RUFOSIGNATA.

Bohem. Mon. Cassid. i. p. 339, 103.
Patr. Bolivia.

139. MESOMPHALIA PICTURATA.

Bohem. Mon. Cassid. i. p. 340, 104.
Patr. Bolivia.

140. MESOMPHALIA RUBROPLAGIATA.

Bohem. Mon. Cassid. i. p. 341, 105.
Patr. Bolivia.

141. MESOMPHALIA ATROGUTTATA.

Bohem. Mon. Cassid. i. p. 342, 106.
Var. h. rufo-testacea, prothorace vitta media lata, antice attenuata, nigra, elytris sutura, fasciis duabus inæqualibus, una non procul a basi utrinque antrorsum ducta in callo humerali desinente, altera in medio ad limbum excurrente, macula humerali, plaga transversa infra medium lineaque laterali versus apicem, nigris; subtus nigra, abdomine late testaceo-marginato.
Patr. Bolivia.

142. MESOMPHALIA OBSOLETA.

Bohem. Mon. Cassid. i. p. 344, 107.
Patr. Bolivia.

143. MESOMPHALIA ATROFASCIATA.

Bohem. Mon. Cassid. i. p. 345, 108.
Patr. Chiquitos, Boliviæ.

144. MESOMPHALIA RUBESCENS.

Bohem. Mon. Cassid. i. p. 346, 109.
Patr. St. Cruz, Boliviæ.

145. MESOMPHALIA FLAVOSIGNATA.

Bohem. Mon. Cassid. i. p. 347, 110.
Patr. Amer. merid. interior.

146. MESOMPHALIA ATRORUBRA.

Bohem. Mon. Cassid. i. p. 349, 111.
Chelimorpha atrorubra, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 290.
Patr. Bolivia.

147. Mesomphalia sexlunata.

Bohem. Mon. Cassid. i. p. 350, 112.
Cassida sexlunata, *Klug*, Preis-Verzeichn. 1829, p. 8, 93.
Chelymorpha sexlunata, *Dej.* Cat. ed. 3, p. 370.
Patr. Brasilia.

148. Mesomphalia tigrina. B.M.

Bohem. Mon. Cassid. i. p. 351, 113.
Patr. Columbia.

149. Mesomphalia Lacordairei. B.M.

Bohem. Mon. Cassid. i. p. 353, 114.
Cyrtonota Lacordairei, *Dej.* Cat. ed. 3, p. 369.
Patr. Buenos Ayres, Corrientes.

150. Mesomphalia subrugosa.

Bohem. Mon. Cassid. i. p. 354, 115.
Cyrtonota subrugosa, *Dej.* Cat. ed. 3, p. 369.
Patr. Buenos Ayres.

151. Mesomphalia adspersa.

Bohem. Mon. Cassid. i. p. 335, 116.
Patr. Brasilia.

152. Mesomphalia Cribrum. B.M.

Bohem. Mon. Cassid. i. p. 356, 117.
Cassida Cribrum, *Klug*, Preis-Verzeichn. 1829, p. 8, 97.
Chelymorpha Cribrum, *Dej.* Cat. ed. 3, p. 370.
Patr. Brasilia meridionalis.

153. Mesomphalia seriata.

Ovata, convexa, dilute flavescens, parum nitida; antennis articulis septem ultimis nigris; prothorace parum convexo, dorso parcius, extrorsum crebre punctato, linea media angusta, longitudinali nigra; elytris mediocriter, crebre punctatis, maculis parvis, sub-remotis, seriatis, nigris decoratis; margine modice explanato, deflexo. Long. 11, lat. 7¼ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Dom. *Deyrolle.*

154. Mesomphalia Augur.

Oblongo-ovata, supra dilute flavescens, subtus nigra; capite, antennis basi, pedibus plagaque magna prothoracis medio

antrorsum dilatata, rufo-ferrugineis; elytris crebre punctatis, sutura rufo-ferruginea. Long. 10½, lat. 7½ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Dom. *Chevrolat.*

155. MESOMPHALIA LUTEIPENNIS.

Bohem. Mon. Cassid. i. p. 357, 118.
Patr. Brasilia.

156. MESOMPHALIA BILINEATA.

Bohem. Mon. Cassid. i. p. 358, 119.
Patr. Brasilia.

157. MESOMPHALIA PLAGICOLLIS.

Bohem. Mon. Cassid. i. p. 360, 120.
Patr. Brasilia.

158. MESOMPHALIA SUSPICIOSA.

Bohem. Mon. Cassid. i. p. 361, 121.
Patr. Rio Janeiro, Brasiliæ.

159. MESOMPHALIA LINEATA.

Bohem. Mon. Cassid. i. p. 361, 122.
Cassidea lineata, *Fabr.* Syst. El. i. p. 406, 104. Ent. Syst. i. p. 305, 59. Mant. Ins. i. p. 65, 50. *Linn.* Syst. Nat. Gmel. i. iv. p. 1641, 62. *Herbst.* Naturs. viii. p. 252, 22, t. cxxxi. f. 7. Dict. Encycl. lit. C. i. p. 134, 68. *Schœnh.* Syn. Ins. ii. p. 227, 132.
Cassida obsoleta, *Oliv.* Ent. vi. p. 948, 41, 97, t. v. f. 70.
Cyrtonota sublineata, *Dej.* Cat. ed. 3, p. 369.
Patr. Cayenna.

160. MESOMPHALIA DELETA. B.M.

Cyrtonota deleta, *Dej.* Cat. ed. 3, p. 393.
Mesomphalia delecta, *Bohem.* Mon. Cassid. i. p. 363, 123.
Patr. Brasilia.

161. MESOMPHALIA FLAVIPENNIS.

Bohem. Mon. Cassid. i. p. 364, 124.
Patr. Brasilia.

162. MESOMPHALIA SUTURALIS. B.M.

Bohem. Mon. Cassid. i. p. 365, 125.
Cassida suturalis, *Fabr.* Syst. El. i. p. 399, 69. Ent. Syst. i.

p. 299, 34. Mant. Ins. i. p. 64, 27. Spec. Ins. i. p. 111, 22. Gen. Ins. p. 219, 7, 8. *Linn.* Syst. Nat. Gmel. i. iv. p. 1640, 56. *Schœnh.* Syn. Ins. ii. p. 222, 94.
Cyrtonota brunnicollis, *Dej.* Cat. ed. 3, p. 393.
Patr. Cayenna, Bahia. Santarem (Amazon), Coll. *H. W. Bates.*

163. MESOMPHALIA VARIANS. B.M.

Bohem. Mon. Cassid. i. p. 366, 126.
Cassida varians, *Fabr.* Syst. El. i. p. 405, 101. *Schœnh.* Syn. Ins. ii. p. 226, 128.
Cassida rufescens, *Herbst.* Naturs. viii. p. 283, 55, t. cxxxiv. f. 1. *Schœnh.* Syn. Ins. ii. p. 219, 66.
Cyrtonota subfasciata, *Dej.* Cat. ed. 3, p. 369.
Var. Cassida hæmatites, *Herbst.* Naturs. viii. p. 280, 53, t. cxxxiii. f. 12. *Schœnh.* Syn. Ins. ii. p. 230, 161.
Patr. Cayenna, Bahia.

164. MESOMPHALIA BICOLORATA.

Bohem. Mon. Cassid. i. p. 368, 127.
Patr. Brasilia.

165. MESOMPHALIA MACULATISSIMA.

Bohem. Mon. Cassid. i. p. 369, 128.
Patr. Brasilia.

166. MESOMPHALIA INCARNATA.

Bohem. Mon. Cassid. i. p. 369, 129.
Patr. Cayenna.

167. MESOMPHALIA DILATATA.

Bohem. Mon. Cassid. i. p. 3.
Cassida dilatata, *Fabr.* Syst. i. p. 399, 68. *Schœnh.* Syn. Ins. ii. p. 222, 93.
Var. a. ut genuina, sed elytrorum plaga flava in medio marginis, macula transversa, obscure ænea.
Patr. Cayenna.

168. MESOMPHALIA BRACHIATA. B.M.

Bohem. Mon. Cassid. i. p. 371, 131.
Cassida brachiata, *Fabr.* Syst. El. i. p. 393, 30. Suppl. p. 80, 16, 17. *Herbst.* Naturs, viii. p. 351, 122. *Schœnh.* Syn. Ins. ii. p. 217, 38.
Var. Cyrtonota transversalis, *Dej.* Cat. ed. 3, p. 369.
Var. Cyrtonota margineguttata, *Dej.* Cat. ed. 3, p. 369.

Var. Cassida quadrinotata, *Herbst.* Naturs. viii. p. 307, 72, t. cxxxv. f. 4.
Patr. Cayenna (*Ent. Club*).

169. Mesomphalia falcifera.

Rotundata, modice convexa, rufo-ferruginea, sub-nitida; antennis articulis quinque ultimis nigris; prothorace subtiliter, sat crebre punctulato, medio linea longitudinali, nigra; elytris dilute flavis vage, mediocriter, sat crebre punctulatis, sutura, margine basali anguste vittaque singuli medio introrsum, apicem versus extrorsum arcuata, nigro æneis; margine modice explanato. Long. 6, lat. 5 millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Reg. Holm.

170. Mesomphalia pertusa.

Bohem. Mon. Cassid. i. p. 373, 132.
Patr. Columbia.

171. Mesomphalia misella.

Bohem. Mon. Cassid. i. p. 374, 133.
Patr. Bahia.

172. Mesomphalia sodalis.

Bohem. Mon. Cassid. i. p. 375, 134.
Patr. Caraccas.

173. Mesomphalia cæruleonotata.

Bohem. Mon. Cassid. i. p. 376, 135.
Patr. Mexico.

174. Mesomphalia pictilis.

Bohem. Mon. Cassid. i. p. 377, 136.
Patr. Mexico, Columbia.

175. Mesomphalia albida.

Bohem. Mon. Cassid. i. p. 378, 137.
Cassida albida, *Germ.* Ins. Spec. i. p. 539, 729.
Chelymorpha quinque punctata, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

176. Mesomphalia Exclamationis. B.M.

Bohem. Mon. Cassid. i. p. 379, 138.

Cassida Exclamationis, *Fabr.* Syst. El. i. p. 406, 105. Ent. Syst. i. p. 305, 60. Mant. Ins. i. p. 65, 49. Spec. Ins. i. p. 114, 41. Syst. Ent. p. 92, 27, *Linn.* Syst. Nat. i. ii. p. 577, 20. Ed. Gmel. i. iv. p. 1641, 20. *Herbst.* Naturs. viii. p. 253, 22, t. cxxxi. f. 8. *Oliv.* Ent. vi. p. 947, 40, 97, t. i. f. 16. Encycl. Ins. 5. p. 350, 65. *Schœnh.* Syn. Ins. ii. p. 227, 133.
Patr. Brasilia, Guyana et Insulæ Antillar. (Martinique).

177. MESOMPHALIA EVANIDA. B.M.

Bohem. Mon. Cassid. i. p. 381, 139.
Chelymorpha evanida, *Dej.* Cat. ed. 3, p. 394.
Patr. Columbia.

Species mihi ignotæ.

178. MESOMPHALIA DECUSSATA.

Bohem. Mon. Cassid. i. p. 382, 140.
Cassida decussata, *Herbst.* Naturs. viii. p. 320, 82, t. cxxxv. f. 12. *Schœnh.* Syn. Ins. ii. p. 227, 131.
Cassida didyma, *Oliv.* Ent. vi. p. 941, 28, 97, t. iv. f. 64.
Patr. India.

179. MESOMPHALIA NITIDULA.

Bohem. Mon. Cassid. i. p. 382, 141.
Cassida nitidula, *Oliv.* Ent. vi. p. 939, 24, 97, t. i. f. 4.
Patr. Guyana.

Genus 19. PŒCILASPIS.

Hope, Col. Man. iii. p. 159. *Bohem.* Mon. Cassid. i. p. 383.
Cassida, *Fabr.* Syst. El. i. p. 387. *Herbst.* Naturs. viii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209. *Germ.* Ins. Spec. i. p. 541. *Perty*, Delect. Animal. p. 102, 103. *Klug*, Preis-Verzeichn. 1829, p. 7, 8.
Botanochara, *Dej.* Cat. ed. 3, p. 393.
Chelymorpha, *Chevrol. Dej.* Cat. ed. 3, p. 393.

I. Elytra elevato-reticulata vel carinata.

A. *Elytra valde convexa.*

1. PŒCILASPIS IMPRESSIPENNIS.

Bohem. Mon. Cassid. i. p. 385, 1.
Patr. Brasilia meridionalis.

2. Pœcilaspis nervosa. B.M.

Bohem. Mon. Cassid. i. p. 386, 2.
Cassida nervosa, *Fabr.* Syst. El. i. p. 405, 100. *Illig.* Mag. v. p. 228, 100. *Schœnh.* Syn. Ins. ii. p. 226, 126.
Botanochara nervosa, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia, Para (Capt. *Sir Everard Home*, Bart.).

3. Pœcilaspis texta. B.M.

Bohem. Mon. Cassid. i. p. 387, 3.
Patr. Brasilia, Corrientes (*Ent. Club*).

4. Pœcilaspis subnervosa.

Bohem. Mon. Cassid. i. p. 388, 4.
Patr. Corrientes.

5. Pœcilaspis ordinata. B.M.

Bohem. Mon. Cassid. i. p. 389, 5.
Botanochara ordinata, *Dej.* Cat. ed. 3, p. 393.
Patr. Columbia (Coll. *Goudot*).

6. Pœcilaspis ruforeticulata. B.M.

Bohem. Mon. Cassid. i. p. 390, 6.
Patr. Brasilia.

7. Pœcilaspis pictipennis.

Bohem. Mon. Cassid. i. p. 391, 7.
Patr. Columbia.

8. Pœcilaspis rubrodelineata.

Bohem. Mon. Cassid. i. p. 392, 8.
Patr. Brasilia.

9. Pœcilaspis macularia.

Bohem. Mon. Cassid. i. p. 393, 9.
Botanochara macularia, *Dej.* Cat. ed. 3, p. 369.
Patr. Buenos Ayres.

10. Pœcilaspis porcata.

Bohem. Mon. Cassid. i. p. 394, 10.
Patr. Buenos Ayres.

11. Pœcilaspis intricata.

Bohem. Mon. Cassid. i. p. 395, 11.
Botanochara intricata, *Klug*, *Dej.* Cat. ed. 3, p. 393.
Var. Botanochara concinna, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia.

12. Pœcilaspis bonariensis. B.M.

Bohem. Mon. Cassid. i. p. 396, 12.
Botanochara bonariensis, *Dej.* Cat. ed. 3, p. 393.
Patr. Buenos Ayres.

13. Pœcilaspis ducalis.

Bohem. Mon. Cassid. i. p. 397, 13.
Botanochara ducalis, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia, Buenos Ayres.

14. Pœcilaspis hæmatodes.

Bohem. Mon. Cassid. i. p. 398, 14.
Cassida hæmatodes, *Perty*, Delect. Anim. p. 102, t. xx. f. 11.
Patr. Brasilia.

15. Pœcilaspis ægrota.

Bohem. Mon. Cassid. i. p. 399, 15.
Botanochara ægrota, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia.

16. Pœcilaspis corticina.

Bohem. Mon. Cassid. i. p. 401, 16.
Patr. Brasilia.

17. Pœcilaspis rubricollis.

Ovata, convexa, sub-nitida, subtus nigra; prothorace rufo-testaceo, obsolete, sub-remote punctulato, apice rotundato; elytris nigro-æneis, opacis, undique irregulariter cupreo-reticulatis, reticulo tenui, modice elevato, nitido, areolas mediocres, profunde punctatas includente; margine modice explanato, deflexo. Long. 13, lat. 9 millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Dom. *Deyrolle.*

18. Pœcilaspis bicruciata.

Ovata, convexa, sub nitida, nigra; antennis articulis 2-5 subtus ferrugineis; prothorace obsoletissime punctulato, medio tenuiter canaliculato, obscure sanguineo, postice plaga media

angulisque nigro-fuscis; elytris sanguineis, mediocriter, sat crebre vage punctatis. elevato-reticulatis, sutura, limbo et cruce singuli nigris. Long. 12½, lat. 8¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Reg. Holm. e Mus. Paris, communicata.

19. Pœcilaspis nigropicta. B.M.

Bohem. Mon. Cassid. i. p. 402, 17.
Var. a. prothorace maculis quatuor nigris, duabus in basi minoribus, magis remotis binisque in dorso antico transversim positis, sub-triangularibus.
Patr. Brasilia. *Var. a.* Mus. Britann. (*Ent. Club*).

20. Pœcilaspis xanthoptera.

Bohem. Mon. Cassid. i. p. 402, 18.
Patr. Brasilia.

B. *Elytra parum convexa.*

21. Pœcilaspis deplanata.

Bohem. Mon. Cassid. i. p. 403, 19,
Patr. Chiquitos.

II. Elytra non reticulata.

22. Pœcilaspis quinquefasciata. B.M.

Bohem. Mon. Cassid. i. p. 405, 20, t. iv. A.
Cassida quinquefasciata, *Perty*, Delect. Anim. p. 103, t. xx. f. 14.
Botanochara Zebra, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia.

23. Pœcilaspis conjuncta.

Bohem. Mon. Cassid. i. p. 406, 21.
Patr. Brasilia, Campos Geraës.

24. Pœcilaspis Erichsoni.

Bohem. Mon. Cassid. i. p. 407, 22.
Patr. Brasilia.

25. Pœcilaspis finitima.

Sub-rotundata, convexa, rufo-testacea, parum nitida; antennis articulis quinque ultimis, prosterno utrinque, pectore toto

abdomineque basi nigro-piceis; prothorace obsolete, lateribus crebrius, subtiliter punctato, ante lobum arcu parvo scutelloque nigris; elytris parum profunde, sat crebre, vage punctulatis, sutura, macula in callo suprahumerali, fascia media transversa, inæquali, communi, plagaque inter medium et apicem nigris. Long. 11¾, lat. 9¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Dom. *Buquet*, Mus. Holm.

26. Pœcilaspis sedecimmaculata.

Bohem. Mon. Cassid. i. p. 408, 23.
Patr. Brasilia.

27. Pœcilaspis pavonia.

Bohem. Mon. Cassid. i. p. 409, 24.
Patr. Brasilia.

28. Pœcilaspis angulata. B.M.

Bohem. Mon. Cassid. i. p. 410, 25.
Cassida angulata, *Germ.* Ins. Spec. i. p. 541, 732.
Botanochara angulata, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia.

29. Pœcilaspis ornata.

Bohem. Mon. Cassid. i. p. 412, 26.
Pœcilaspis pardalina. *Bohem.* Mon. Cassid. i. p. 429, 41.
Patr. Brasilia.

30. Pœcilaspis pantherina.

Bohem. Mon. Cassid. i. p. 413, 27.
Cassida pantherina, *Klug*, Preis-Verzeichn. 1829, p. 7, 91.
Botanochara pantherina, *Dej.* Cat. ed. 3, p. 393.
Patr. Buenos Ayres, Montevideo.

31. Pœcilaspis duodecimmaculata.

Bohem. Mon. Cassid. i. p. 414, 28.
Patr. Brasilia.

32. Pœcilaspis cancellata.

Bohem. Mon. Cassid. i. p. 415, 29.
Botanochara cancellata, *Dej.* Cat. ed. 3, p. 393.
Patr. Tucuman.

33. Pœcilaspis decempustulata. B.M.

Bohem. Mon. Cassid. i. p. 416, 30.
Botanochara decempustulata, *Dej.* Cat. ed. 3, p. 369.
Patr. Tucuman, Rio de la Plata. Bolivia (Coll. *Bridges*).

34. Pœcilaspis octopustulata.

Bohem. Mon. Cassid. i. p. 417, 31.
Cassida octopustulata, *Klug*, Preis-Verzeichn. 1829, p. 7, 90.
Botanochara octopustulata, *Dej.* Cat. ed. 3, p. 369.
Patr. Brasilia meridionalis.

35. Pœcilaspis sedecimpustulata. B.M.

Bohem. Mon. Cassid. i. p. 418, 32.
Cassida sedecimpustulata, *Fabr.* Spec. Ins. i. p. 115, 47. *Linn.* Syst. Nat. Gmel. i. iv. p. 1643, 68. *Oliv.* Encycl. Ins. v. p. 392, 76. Entom. vi. p. 940, 26, 97, t. iii. f. 41 (mala).
Cassida sedecimpunctata, *Fabr.* Syst. El. i. p. 498, 118. Ent. Syst. i. p. 307, 68. Mant. Ins. i. p. 66, 56. *Herbst.* Naturs. viii. p. 349, 119. *Schœnh.* Syn. Ins. ii. p. 229, 150.
Cassida tessellata, *Voet.* Col. ii. t. xliii. f. 20.
Botanochara quatuordecimpustulata, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia.

36. Pœcilaspis quatuordecimsignata.

Rotundata, convexa, nigra, sub-opaca; antennis articulis 2-5 rufo-testaceis; prothorace subtiliter, lateribus crebrius punctulato, antice utrinque macula parva, oblonga, sanguinea; elytris subtiliter, sub-remote punctulatis, singulo maculis septem mediocribus, sanguineis; humeris rotundatis. Long. 10¾, lat. 8½ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Dom. *Jekel*, Mus. Holm.

37. Pœcilaspis quatuordecimplagiata.

Bohem. Mon. Cassid. i. p. 420, 33.
Patr. Chiquitos.

38. Pœcilaspis rubroguttata. B.M.

Bohem. Mon. Cassid. i. p. 421, 34.
Patr. Brasilia meridionalis.

39. Pœcilaspis decemverrucata. B.M.

Bohem. Mon. Cassid. i. p. 422, 35.
Patr. Bolivia (Coll. *Bridges*).

40. Pœcilaspis decemnotata.

Bohem. Mon. Cassid. i. p. 423, 36.
Patr. Brasilia.

41. Pœcilaspis duodecimverrucata.

Bohem. Mon. Cassid. i. p. 424, 37.
Botanochara duodecimverrucata, *Dej.* Cat. ed. 3, p. 393.
Patr. Buenos Ayres.

42. Pœcilaspis duodecimnotata.

Bohem. Mon. Cassid. i. p. 425, 38.
Var. a. elytrorum maculis minoribus.
Var. b. elytrorum maculis duabus posticis inter se conjunctis et interdum cum macula media disci confluentibus.
Patr. Brasilia. *Var. a.* et *b.* Mus. Berol.

43. Pœcilaspis fastidita.

Bohem. Mon. Cassid. i. p. 426, 39.
Botanochara fastidita, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia.

44. Pœcilaspis invasa.

Bohem. Mon. Cassid. i. p. 427, 40.
Patr. Bolivia.

45. Pœcilaspis encausta.

Bohem. Mon. Cassid. i. p. 430, 42.
Botanochara encausta, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia.

46. Pœcilaspis anthracina.

Rotundata, convexa, nigra, sub-nitida; antennis articulis 2-rufo-piceis; prothorace subtiliter, remote, dorso parcius punctulato, antice utrinque macula angusta, flavo-testacea, ramulum obsoletum, concolorem introrsum emittente, dorso minus crebre, mediocriter, margine crebrius punctulatis. Long. 10, lat. 8½ millim.

Bohem. Mon. Cassid. Append.
Var. a. prothorace toto nigro.
Patr. St. Paul, Brasiliæ, Mus. Dom. *Chevrolat.*

47. Pœcilaspis suturella.

Bohem. Mon. Cassid. i. p. 431, 43.
Chelymorpha suturella, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

48. Pœcilaspis Histrio.

Bohem. Mon. Cassid. i. p. 432. 44.
Botanochara Histrio, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia meridionalis.

49. Pœcilaspis duplicata.

Bohem. Mon. Cassid. i. p. 433, 45.
Patr. Brasilia.

50. Pœcilaspis vittifera.

Bohem. Mon. Cassid. i. p. 434, 46.
Patr. Allegretto, Brasiliæ.

51. Pœcilaspis proxima.

Bohem. Mon. Cassid. i. p. 435, 47.
Patr. Brasilia, Montevideo.

Genus 20. ELYTROGONA.

Chevrolat. Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. i. p. 438.
Cassida, *Oliv.* Entom. vi. p. 922. *Latr.* Ann. Mus. d'Hist. Nat. p. 298. *Dalm.* Anal. Entom. p. 82.

1. Elytrogona Ampulla. B.M.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. i. p. 440, 1, t. iv. B.
Cassida Ampulla, *Oliv.* Entom. vi. p. 958, 59, 97, t. v. f. 80.
Cassida quatuordecimmaculata, *Latr.* Ann. d'Hist. Nat. f. i. p. 298, t. xxi. f. 10.
Cassida fornicata, *Dalm.* Anal. Entomologica, p. 82, 85.
Patr. St. Domingo, Cuba.

2. Elytrogona interjecta.

Rotundata, valde convexa, rubra, sub-nitida; antennis flavo-testaceis; prothorace lævi; elytris fornicatis, punctis magnis, profundissimis, confertis velut cribrosis, singulo maculis septem nigris, lateribus ante medium angulatis. Long. 8, lat. 7 millim.

Bohem. Mon. Cassid. Append.
Patr. St. Domingo, Mus. Holm.

3. Elytrogona Bacca. B.M.

Klug, Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. i. p. 441, 2.
Patr. St. Domingo.

Genus 21. CHELYMORPHA.

Chevrolat. Dej. Cat. ed. 3, p. 393. *Bohem.* Mon. Cassid. ii. p. i.
Cassida, *Fabr.* Syst. El. i. p. 387. *Herbst.* Naturs. viii. p. 204. *Oliv.* Entom. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209. *Klug,* Preis-Verzeichn. 1829, p. 7, 8.
Cyrtonota, *Chevrol.* Dej. Cat. ed. 3, p. 392, ex part.
Cyphomorpha, *Hope,* Coleopt. Man. iii. p. 160. *Erichs.* Archiv. f. Naturg. xiii. p. 153.

I. Elytra ante medium dorsi gibbosa.

1. CHELYMORPHA HOEPFNERI. B.M.

Dej. Cat. ed. 3, p. 393. *Bohem.* Mon. Cassid. ii. p. 2, 1.
Patr. Mexico.

2. CHELYMORPHA COMATA. B.M.

Bohem. Mon. Cassid. ii. p. 4, 2.
Patr. Mexico.

3. CHELYMORPHA PUBESCENS. B.M.

Chevrolat, Dej. Cat. ed. 3, p. 393. *Bohem.* Mon. Cassid. ii. p. 5, 3.
Patr. Mexico (*Ent. Club*).

4. CHELYMORPHA BRUNNEA. B.M.

Dej. Cat. ed. 3, p. 393. *Bohem.* Mon. Cassid. ii. p. 6, 4.
Cassida brunnea, *Fabr.* Ent. Syst. Suppl. p. 81, 9, 10. Syst. El. i. p. 391, 17. *Herbst.* Naturs. viii. p. 350, 120. *Oliv.* Entom. vi. p. 959, 60, 97, t. ii. f. 20. *Schœnh.* Syn. Ins. ii. p. 216, 25.
Cassida cincta, *De Geer,* Mem. Ins. v. p. 186, 16, t. xv. f. 14. *Schœnh.* Syn. Ins. ii, p. 230, 156.
Cassida tuberculata, *Oliv.* Encycl. Ins. v. p. 384, 35.
Cassida marginata, *Linné,* Syst. Nat. i. ii. p. 576, 14.
Patr. Cayenna, Bahia, Brasiliæ. Napo. (Coll. *H. W. Bates*).

5. CHELYMORPHA CAVATA. B.M.

Bohem. Mon. Cassid. ii. p. 7, 5.
Patr. Columbia.

6. CHELYMORPHA WOLLASTONI. B.M.

Bohem. Mon. Cassid. ii. p. 8, 6.
Patr. Pernambuco.

7. Chelymorpha gibba. B.M.

Dej. Cat. ed. 3, p. 393. *Bohem.* Mon. Cassid. ii. p. 9, 7, t. iv. C.
Cassida gibba, *Fabr.* Syst. El. i. p. 395, 43. Ent. Syst. Suppl. p. 82, 21, 22. *Herbst.* Naturs. viii. p. 352, 124. *Schœnh.* Syn. Ins. ii. p. 218, 62.
Cassida variolosa, *Oliv.* Encycl. meth. v. p. 385, 40. Entom. vi. p. 958, 58, 97, t. ii. f. 21, t. iii. f. 46.
Cassida cribraria, *Herbst.* Naturs. viii. p. 281, 54, t. cxxx. f. 6.
Patr. Cayenna.

8. Chelymorpha constellata. B.M.

Bohem. Mon. Cassid. i. p. 10, 8.
Cassida constellata, *Klug*, Preis-Verzeichn. 1829, p. 7, 87.
Chelymorpha æneomaculata, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia, Santarem (Coll. *H. W. Bates*).

9. Chelymorpha inflata. B.M.

Bohem. Mon. Cassid. ii. p. 12, 9.
Patr. Brasilia.

10. Chelymorpha distinguenda. B.M.

Dej. Cat. ed. 3. p. 393. *Bohem.* Mon. Cassid. ii. p. 13, 10.
Patr. Brasilia.

11. Chelymorpha socia. B.M.

Bohem. Mon. Cassid. ii. p. 14, 11.
Patr. Brasilia, Bahia.

12. Chelymorpha clivosa.

Bohem. Mon. Cassid. ii. p. 16, 12.
Patr. Brasilia.

13. Chelymorpha infaceta.

Bohem. Mon. Cassid. ii. p. 17, 13.
Patr. Corrientes.

14. Chelymorpha Sturmii.

Dej. Cat. ed. 3, p. 393. *Bohem.* Mon. Cassid. ii. p. 18, 14.
Patr. Brasilia.

15. Chelymorpha bullata.

Bohem. Mon. Cassid. ii. p. 19, 15.
Patr. Brasilia.

16. Chelymorpha subpunctata.

Bohem. Mon. Cassid. ii. p. 20, 16.
Patr. Brasilia.

17. Chelymorpha ignobilis.

Bohem. Mon. Cassid. ii. p. 21, 17.
Patr. Brasilia.

18. Chelymorpha urbana.

Bohem. Mon. Cassid. ii. p. 22, 18.
Patr. Brasilia.

19. Chelymorpha orthogonia.

Bohem. Mon. Cassid. ii. p. 23, 19.
Cyrtonota orthogonia, *Dej.* Cat. ed. 3, p. 393.
Patr. Brasilia.

20. Chelymorpha Catharinensis.

Rotundata (*mas*) aut ovata (*femina*) obscure cæruleo-virescens, sub-nitida, subtus nigra; antennis articulis quatuor basalibus testaceis; prothorace sub-remote punctato, vitta media maculisque duabus testaceis; elytris subtiliter coriaceis, mediocriter, vage punctatis, obsolete rugosis, basi maculis 1 vel 2 parvis, testaceis notatis, dorso antico obtuse gibbosis. Long. *maris*, 9, lat. 7. Long. *feminæ*, 12, lat. 8½ millim.

Bohem. Mon. Cassid. Append.
Var. a. prothoracis vitta media brevi, maculis lateralibus obsoletioribus, elytris immaculatis.
Patr. St. Catharina, Brasiliæ, Dom. *Deyrolle*, Mus. Reg. Holm.

21. Chelymorpha cobaltina. B.M.

Bohem. Mon. Cassid. ii. p. 24, 20.
Patr. Bahia, Brasiliæ.

II. Elytra ante medium dorsi non gibbosa.

A. *Prothorax flavus vel rufo-testaceus, immaculatus.*

22. Chelymorpha flavicollis.

Dej. Cat. ed. 3, p. 393. *Bohem.* Mon. Cassid. ii. p. 25, 21.
Cassida flavicollis, *Klug*, Preis-Verzeichn. 1829, p. 14, 204.
Patr. Cuba.

23. CHELYMORPHA SWARTZII. B.M.

Bohem. Mon. Cassid. ii. p. 26, 22.
Cassida Swartzii, *Thunb.* Schœnh. Syn. Ins. ii. p. 229, 152, t. iv. f. 12.
Patr. Jamaica (Coll. *P. H. Gosse*).

24. CHELYMORPHA PUNCTIPENNIS. B.M.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 27, 23.
Patr. Brasilia.

25. CHELYMORPHA APIATA. B.M.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 28, 24.
Patr. Brasilia.

26. CHELYMORPHA OBDUCTA.

Bohem. Mon. Cassid. ii. p. 29, 25.
Patr. Bolivia, Brasilia.

27. CHELYMORPHA CALVA.

Bohem. Mon. Cassid. ii. p. 30, 26.
Patr. Mexico.

28. CHELYMORPHA PEREGRINA.

Bohem. Mon. Cassid. ii. p. 31, 27.
Patr. Bolivia.

29. CHELYMORPHA RUSSULA.

Bohem. Mon. Cassid. ii. p. 32, 28.
Patr. Brasilia.

30. CHELYMORPHA ALTERNANS. B.M.

Bohem. Mon. Cassid. ii. p. 33, 29.
Patr. Columbia. *Var.* Para, Santarem (Coll. *H. W. Bates*).

31. CHELYMORPHA INFACETA.

Bohem. Mon. Cassid. ii. p. 34, 30.
Patr. Bolivia.

32. CHELYMORPHA PARUMMACULATA.

Bohem. Mon. Cassid. ii. p. 34, 31.
Patr. Brasilia.

B. *Prothorax flavus vel rufo-testaceus, nigro-maculatus vel variegatus.*

33. Chelymorpha catenulata.

Bohem. Mon. Cassid. ii. p. 35, 32.
Patr. Mexico.

34. Chelymorpha Smithii. B.M.

Ovata, valde convexa, rufo-testacea, sub-nitida, pube brevi, pallida, adspersa; antennis extrorsum, prosterno et pectore totis abdomineque basi medio infuscatis; prothorace confertim, mediocriter punctato, ante medium maculis quatuor rotundatis et basi duabus obliquis, approximatis, nigris; elytris sat crebre, mediocriter punctatis, pone scutellum macula magna suturali, singulo maculis duabus, fascia media valde inæquali plagaque postica magna nigris. Long. 10, lat. 8 millim.

Bohem. Mon. Cassid. Append.
Var. a. prothorace maculis tantum duabus ante medium, rotundis, nigris, elytrorum macula basali interiore cum macula suturali connexa, corpore subtus ferrugineo.
Patr. Honduras, Mus. Britann. (Coll. *Dyson*). *Var. a.* Mus. Holm.

35. Chelymorpha undecimmaculata. B.M.

Bohem. Mon. Cassid. ii. p. 37, 33.
Var. b. prothorace nigro-trimaculato, maculis mediocribus, media in medio disci, lateralibus basi adfixis.
Var. c. tota nigra, capite, antennis basi, tibiis anticis totis, posticis apice, prothorace antice maculaque utrinque ovata, ultra medium extensa, rufo-ferrugineis.
Patr. Mexico.

36. Chelymorpha fasciata. B.M.

Oblongo-ovata convexa, rufo-testacea; antennis extrorsum, capite corporeque subtus nigris; abdomine segmentis apice ferrugineis; prothorace mediocriter, minus crebre punctato, medio tenuiter canaliculato, maculis duabus, rotundis transversim positis scutelloque nigris; elytris mediocriter, crebre punctatis, basi maculis duabus sat magnis, macula pone scutellum communi, fascia media sat lata inæquali maculisque duabus inter medium et apicem nigris. Long. 12, lat. 8 millim.

Bohem. Mon. Cassid. Append.
Patr. Mexico, Mus. Britann.

37. Chelymorpha Argus. B.M.

Bohem. Mon. Cassid. ii. p. 38, 34.
Cassida cribraria, *Oliv.* Encyl. Ins. v. p. 383, 27. Ent. vi. p. 956, 54, 97, t. iii. f. 44. *Schœnh.* Syn. Ins. ii. p. 219, 65.
Chelymorpha cribraria, *var.* 17 punctata, *Dej.* Cat. ed. 3, p. 394.
Var. Chelymorpha cribraria, *Dej.* Cat. ed. 3, p. 394.
Var. Cassida Argus, *Herbst.* Naturs. viii. p. 278, 50, t. cxxxiii. f. 9. *Schœnh.* Syn. Ins. ii. p. 230, 160.
Var. Chelymorpha geniculata, *Dej.* Cat. ed. 3, p. 394.
Patr. America borealis (Delaware), *E. Doubleday, Esq.* Cuba.

38. Chelymorpha quatuordecimpunctata.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 40, 35.
Var. a. elytrorum maculis duabus posticis convexis, fasciam brevem, obliquam formantibus.
Var. b. ut var. a sed maculis elytrorum mediis etiam confluentibus.
Patr. Mexico. *Var. a.* et *b,* Mus. Dom. *Deyrolle.*

39. Chelymorpha sedecimstillata. B.M.

Ovata, modice convexa, dilute flavo-testacea, sub-nitida; antennis extrorsum nigris; prothorace subtiliter, sub-remote punctulato, medio canaliculato, maculis duabus ante medium rotundis, nigris, transversim positis; elytris mediocriter, minus crebre punctatis, singulo maculis septem inæqualibus nigris. Long. 9, lat. 6 millim.
Bohem. Mon. Cassid. Append.
Patr. Mexico, Mus. Britann.

40. Chelymorpha duodecimsignata.

Bohem. Mon. Cassid. ii. p. 41, 36.
Patr. Mexico.

41. Chelymorpha tredecimmaculata.

Bohem. Mon. Cassid. ii. p. 42, 37.
Patr. Mexico.

42. Chelymorpha conjugata. B.M.

Bohem. Mon. Cassid. ii. p. 43, 38.
Var. a. maculis nigris prothoracis elytrorumque parvis, præsertim duabus posticis.
Patr. Mexico.

43. Chelymorpha cingulata.

Bohem. Mon. Cassid. i. p. 44, 39.
Patr. Brasilia meridionalis.

44. CHELYMORPHA LONGULA.

Oblongo-ovata, convexa, supra sordide flavo-testacea, subtus nigra; antennis basi, femoribus apice tibiisque anticis flavo-testaceis; prothorace minus crebre, vage punctulato, medio maculis duabus parvis, transversim positis fuscis; scutello nigro; elytris mediocriter, sat crebre punctatis, singulo pone medium obsolete bi-carinato; margine parum explanato. Long. 9¾, lat. 5½ millim.

Bohem. Mon. Cassid. Append.
Patr. Brasilia, Mus. Dom. *Deyrolle.*

45. CHELYMORPHA PUNCTICOLLIS. B.M.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 45, 40.
Patr. Brasilia.

46. CHELYMORPHA PARTITA.

Bohem. Mon. Cassid. ii. p. 46, 41.
Patr. Brasilia.

47. CHELYMORPHA DISCONOTATA.

Bohem. Mon. Cassid. ii. p. 47, 42.
Patr. Cayenna.

48. CHELYMORPHA PRÆTEXTATA. B.M.

Bohem. Mon. Cassid. ii. p. 48, 43.
Patr. Bolivia (Coll. *Bridges*).

49. CHELYMORPHA TESTACEOMARGINATA.

Bohem. Mon. Cassid. ii. p. 49, 44.
Patr. Columbia.

50. CHELYMORPHA PELTATA.

Bohem. Mon. Cassid. ii. p. 50, 45.
Patr. Brasilia.

51. CHELYMORPHA PUNCTIGERA.

Bohem. Mon. Cassid. ii. p. 51, 46.
Patr. Brasilia.

52. CHELYMORPHA PACIFICA.

Bohem. Mon. Cassid. ii. p. 52, 47.
Patr. Bolivia.

53. Chelymorpha personata.

Bohem. Mon. Cassid. ii. p. 53, 48.
Patr. Montevideo.

54. Chelymorpha indigesta.

Bohem. Mon. Cassid. ii. p. 54, 49.
Patr. Montevideo, Buenos Ayres.

55. Chelymorpha cribraria. B.M.

Bohem. Mon. Cassid. ii. p. 55, 50.
Cassida cribraria, *Fabr.* Syst. El. i. p. 396, 46. Ent. Syst. i. p. 297, 23. Mant. Ins. i. p. 63, 20. Spec. Ins. i. p. 110, 16. *Linn.* Syst. Nat. ed. Gmel. i. iv. p. 1636, 33.
Cassida multipunctata, *Oliv.* Encycl. Ins. v. p. 384, 36. Entom. vi. p. 957, 97, t. v. f. 79.
Chelymorpha multipunctata, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia, Para (Coll. *H. W. Bates*), Cayenna, Guyana.

56. Chelymorpha polysticha. B.M.

Bohem. Mon. Cassid. ii. p. 56, 51.
Patr. Guadeloupe, St. Barthelemy.

57. Chelymorpha atomaria.

Bohem. Mon. Cassid. ii. p. 57, 52.
Patr. Columbia.

58. Chelymorpha advena.

Ovata, convexa, sub-nitida, supra testacea, subtus nigra, antennis nigris, articulis 1 apice, 2-5 totis rufo-testaceis; prothorace subtiliter, sub-remote punctulato, fere in medio maculis duabus parvis, transversim positis scutelloque nigris; elytris mediocriter, sub-remote, vage punctulatis, disco postico maculis 2 vel 3 parvis, nigris notatis. Long 7¼, lat. 5½ millim.

Bohem. Mon. Cassid. Append.
Patr. Buenos Ayres, Mus. Dom. *Chevrolat.*

59. Chelymorpha commutabilis.

Bohem. Mon. Cassid. ii. p. 58, 53.
Patr. Brasilia.

60. Chelymorpha patagonica.

Bohem. Mon. Cassid. ii. p. 60, 54.
Patr. Patagonia.

61. CHELYMORPHA VARIABILIS. B.M.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 61, 55.
Patr. Buenos Ayres, Rio Janeiro, Corrientes.

62. CHELYMORPHA VARIANS. B.M.

Ovata, convexa, nitida, supra testacea, subtus nigra; prothorace sub-remote, subtiliter punctulato, plaga magna dorsali maculaque utrinque laterali, sub-triangulari; nigris, illa medio linea longitudinali maculaque antica testaceis; elytris vage, sat crebre, mediocriter punctatis, sutura anguste maculisque numerosis, inæqualibus, nigris; margine parum explanato, nigro-maculato. Long. 9½, lat. 6¾ millim.

Blanchard, Hist. Fisica y Politica de Chile Zool. t. v. p. 530, 1, t. xxi. f. 4.
Bohem. Mon. Cassid. Append.
Var. a. prothorace vittis duabus latis, a basi ultra medium continuatis, antrorsum linea angusta, transversa connexis, macula utrinque laterali, sub-triangulari, sutura maculisque disci exterioris minoribus, nigris.
Var. b. prothorace ut in *var. a.* signato, linea tamen transversa deficiente, sutura vix infuscata, disco exteriore elytrorum parce nigro-maculato.
Var. c. prothorace ut in *b.* macula utrinque laterali parva, punctiformi; elytris immaculatis, puncto parvo, supra humerali nigro notatis.
Patr. Chili, Mus. Paris.

63. CHELYMORPHA HONESTA.

Bohem. Mon. Cassid. ii. p. 62, 56.
Patr. Brasilia.

64. CHELYMORPHA HÆMATURA.

Bohem. Mon. Cassid. ii. p. 63, 57.
Patr. Brasilia.

65. CHELYMORPHA TUCUMANA.

Dej. Cat. ed. 3, p. 370. *Bohem.* Mon. Cassid. ii. p. 64, 58.
Patr. Tucuman.

66. CHELYMORPHA INSIGNIS.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 65, 59.
Cassida insignis, *Klug*, Preis-Verzeichn. 1829, p. 8, 94.
Patr. Brasilia.

67. CHELYMORPHA BIANNULARIS. B.M.

Klug, Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 66, 60.
Patr. Mexico.

68. CHELYMORPHA PRINCEPS.

Bohem. Mon. Cassid. ii. p. 68, 61.
Patr. Bolivia.

69. CHELYMORPHA IMPERIALIS.

Guér. Icon. du Règne Anim. Ins. texte ii. p. 290. *Dej.* Cat. ed. 3, p. 370.
Bohem. Mon. Cassid. ii. p. 69, 62.
Patr. Buenos Ayres, Bolivia.

70. CHELYMORPHA CIRCUMPUNCTATA. B.M.

Bohem. Mon. Cassid. ii. p. 70, 63.
Cassida circumpunctata, *Klug*, Preis-Verzeichn. 1829, p. 8, 95.
Chelymorpha Schüppelii, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

71. CHELYMORPHA NOTATA. B.M.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 72, 64.
Patr. Bahia, Brasiliæ.

C. *Prothorax niger, interdum plus minusve rufotestaceo-marginatus.*

72. CHELYMORPHA PANTHERINA. B.M.

Blanchard, Voy. d'Orbigny, Ins. pl. 24, f. 5. *Bohem.* Mon. Cassid. ii. p. 73, 65.
Patr. Bolivia.

73. CHELYMORPHA BOLIVIANA. B.M.

Bohem. Mon. Cassid. ii. p. 74, 66.
Patr. Bolivia (Coll. *Bridges*).

74. CHELYMORPHA DECEMSTILLATA.

Rotundata, convexa, nigra, sub-nitida, glabra; antennis articulis 2-4 testaceis; prothorace subtiliter, minus crebre margine laterali evidentius punctulato, antice utrinque anguste testaceo-marginato; elytris sat crebre, mediocriter punctatis, singulo maculis quinque magnis, sanguineis, ma-

cula intra-apicali cuneata; margine modice explanato, obliquedeflexo. Long. 10½, lat. 8½.

Bohem. Mon. Cassid. Append.
Patr. Mexico, Mus. Berol.

75. CHELYMORPHA KLUGII.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 75. 67.
Patr. Brasilia meridionalis.

76. CHELYMORPHA INCLUSA.

Bohem. Mon. Cassid. ii. p. 76, 68.
Patr. Novum Friburgum, Brasiliæ.

77. CHELYMORPHA RUFIPENNIS. B.M.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 77, 69.
Var. a. prothorace antice latius testaceo-marginato, lineola brevi, longitudinali, dorsali testacea.
Patr. Brasilia, Venezuela (Coll. *Bridges*).

78. CHELYMORPHA RUFICEPS.

Bohem. Mon. Cassid. ii. p. 78, 70.
Patr. Brasilia.

79. CHELYMORPHA ADNATA.

Bohem. Mon. Cassid. ii. p. 79, 71.
Patr. Brasiliæ.

80. CHELYMORPHA NIGRICOLLIS. B.M.

Dej. Cat. ed. 3. p. 393. *Bohem.* Mon. Cassid, ii. p. 80, 72.
Var. a. prothorace medio linea longitudinali testacea.
Patr. Bahia, Brasiliæ.

81. CHELYMORPHA MULTIGUTTATA.

Bohem. Mon. Cassid. ii. p. 81, 73.
Patr. Bolivia.

82. CHELYMORPHA VERMICULATA.

Bohem. Mon. Cassid. ii. p. 82, 74.
Patr. Brasilia.

83. CHELYMORPHA VENUSTA.

Bohem. Mon. Cassid. ii. p. 83, 75.
Var. a. prothorace sanguineo, maculis sex nigris notato, dua-

bus prope apicem approximatis, pyriformibus, duabus in ipsa basi transversis et una utrinque in medio marginis minore, transversa.
Patr. Brasilia.

D. *Prothorax niger, flavo- vel testaceo-variegatus.*

84. CHELYMORPHA JUVENCA. B.M.

Bohem. Mon. Cassid. ii. p. 84, 76.
Cyrtonota juvenca, *Dej.* Cat. ed. 3, p. 393.
Var. e. elytris totis testaceis, margine anguste anguloque humerali latius nigris.
Patr. Mexico. *Var. e.* Mus. Britann.

85. CHELYMORPHA GRAPHIPTERA.

Bohem. Mon. Cassid. ii. p. 85, 77.
Patr. Montevideo.

86. CHELYMORPHA GUTTIFERA.

Bohem. Mon. Cassid. ii. p. 86, 78.
Patr. Guatimala.

87. CHELYMORPHA AREATA.

Rotundata (*mas*) aut breviter ovalis (*femina*), convexa, nigra, sub-nitida; fronte antennisque basi rufis; prothorace remote, subtiliter punctulato, maculis tribus rufis notato; elytris sat profunde, crebre punctatis, singulo maculis tribus sanguineis, duabus prope suturam, magnis, tertia laterali, mire. Long. *maris*, 8½, lat. 6½. Long. *feminæ*, 10½, lat. 7½ millim.

Bohem. Mon. Cassid. ii. p. 94, 85. *Bohem.* Mon. Cassid. Append.
Cyphomorpha areata, *Erichs.* Arch. f. Naturg. xiii. p. 153, 1.
Var. a. prothorace præter maculam mediam utrinque vitta lata, obliqua, fere ad basin continuata femoribusque anterioribus rufis.
Var. b. macula postica elytrorum parva, rotunda.
Patr. Peruvia orientalis.

88. CHELYMORPHA CRUCIFERA.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 87, 79.
Var. Chelymorpha signaticollis, *Dej.* Cat. ed. 3, p. 370.
Patr. Buenos Ayres.

89. CHELYMORPHA GUTTULA.

Bohem. Mon. Cassid. ii. p. 89, 80.
Var. a. prothorace nigro, antice in dorso macula parva vittaque utrinque angustiore sanguineis, elytrorum maculis duabus anticis prope suturam confluentibus, maculis reliquis minoribus, postica rotundata.
Patr. Montevideo.

90. CHELYMORPHA CINCTIPENNIS.

Bohem. Mon. Cassid. ii. p. 90, 81.
Patr. Brasilia.

91. CHELYMORPHA SCITULA. B.M.

Bohem. Mon. Cassid. ii. p. 91, 82.
Patr. Cayenna.

92. CHELYMORPHA CÆCA.

Bohem. Mon. Cassid. ii. p. 92, 83.
Patr. Brasilia.

93. CHELYMORPHA INFIMA.

Bohem. Mon. Cassid. ii. p. 93, 84.
Patr. Columbia.

Genus 22. SELENIS.

Hope, Ann. Nat. Hist. iii. t. iv. f. 3. *Bohem.* Mon. Cassid. ii. p. 94.
Cassida, *Linn.* Syst. Nat. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *Herbst.* Naturs. viii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209.
Acromis, *Chevrol.* Dej. Cat. ed. 3, p. 370. *Erichs.* Arch. f. Naturg. xiii. p. 153.

1. SELENIS SPINIFEX. B.M.

Bohem. Mon. Cassid. ii. p. 96, 1, t. iv. D.
Cassida Spinifex, *Linn.* Syst. Nat. i. ii. p. 576, 7. Ed. Gmel. i. iv. p. 1638, 7. Amœn. Acad. vi. p. 392, 7. Cent. Ins. No. 7. *Fabr.* Syst. El. i. p. 402, 85. Ent. Syst. i. p. 302, 47. Mant. Ins. i. p. 65, 38. Spec. Ins. i. p. 112, 31. Syst. Ent. p. 91, 20. *Herbst.* Naturs. viii. p. 327, 88, t. cxxxvi. f. 4, 5. *Illig.* Mag. v. p. 227, 85. *Oliv.* Enc. Ins. v. p. 387, 52. Entom. vi. p. 931, 11, 97, t. iv. f. 57, a. b. *Schœnh.* Syn. Ins. ii. p. 224, 112.

Cassida Cerastes, *Voet.* Col. ii. t. xliii. f. 21.
Acromis Spinifex, *Dej.* Cat. ed. 3, p. 394.
Var. Cassida Spinifex, *Fabr.* Syst. El. i. p. 402, 85. Var. β.
Cassida perforata, *Fabr.* Ent. Syst. i. p. 301, 43. Mant. Ins. i. p. 64, 35. Spec. Ins. i. p. 111, 28. Gen. Ins. p. 219, 11, 12. *Pallas,* Spicil. Zool. ix. p. 3, t. i. f. 1. *Pallas,* Naturg. ix. p. 3, t. i. f. 1. *Linn.* Syst. Nat. Gmel. i. iv. p. 1638, 50. *Herbst.* Naturs. viii. p. 343, 110. *Oliv.* Enc. Ins. v. p. 386, 47. Entomol. vi. p. 932, 12, 97, t. iv. f. 58.
Var. Cassida angustata, *Linn.* Syst. Nat. i. ii. p. 578, 31? *Oliv.* Enc. Ins. v. p. 390, 67? Entom. vi. p. 933, 14, 97, t. i. f. 12.
Patr. Cayenna, Brasilia, Surinam, Guyana.

2. Selenis venosa.

Bohem. Mon. Cassid. ii. p. 98, 2.
Acromis venosa, *Erichs.* Archiv. f. Naturg. xiii. p. 153, 1.
Patr. Peruvia orientalis.

3. Selenis nebulosa. B.M.

Bohem. Mon. Cassid. ii. p. 99, 3.
Patr. Bolivia, Chiquitos, Venezuela (Coll. *Dyson*).

4. Selenis sparsa. B.M.

Bohem. Mon. Cassid. ii. p. 100, 4.
Patr. Mexico, Nova Granada, Columbia (Coll. *Goudot*).

Genus 23. OMOPLATA.

Hope, Coleopt. Man. iii. p. 159. *Bohem.* Mon. Cassid. ii. p. 101.
Cassida, *Linn.* Syst. Nat. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *De Geer,* Ins. v. p. 184. *Herbst.* Naturs. viii. p. 204. *Oliv.* Entom. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209. *Germ.* Ins. Spec. i. p. 727.
Echoma, *Chevrol,* Dej. Cat. ed. 3, p. 394.

I. *Prothorax breviter triangularis, lateribus basin versus oblique ampliatus.*

1. Omoplata aulica. B.M.

Bohem. Mon. Cassid. ii. p. 103, 1.
Echoma aulica, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

2. Omoplata normalis. B.M.

Bohem. Mon. Cassid. ii. p. 104, 2.

Cassida normalis, *Germ.* Ins. Spec. i. p. 538, 727.
Cassida suturalis, *Herbst.* Naturs. viii. p. 339, 104. *Oliv.* Enc. Ins. v. p. 385, 38. Entomol. vi. p. 942, 31, 97, t. iv. f. 65. *Schœnh.* Syn. Ins. ii. p. 222, 94.
Echoma suturalis, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

3. OMOPLATA TRANSVERSALIS.

Bohem. Mon. Cassid. ii. p. 105, 3.
Patr. Brasilia.

4. OMOPLATA TERSA.

Bohem. Mon. Cassid. ii. p. 106, 4.
Patr. Brasilia.

5. OMOPLATA DICHROA. B.M.

Bohem. Mon. Cassid. ii. p. 107, 5.
Cassida dichroa, *Germ.* Ins. Spec. i. p. 538, 728.
Echoma basalis, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

6. OMOPLATA GENEROSA.

Bohem. Mon. Cassid. ii. p. 126, 23.
Patr. Brasilia.

7. OMOPLATA NUPTA.

Bohem. Mon. Cassid. ii. p. 109, 6.
Patr. Guyana Britannica.

8. OMOPLATA FLAVA.

Bohem. Mon. Cassid. ii. p. 110, 7.
Cassida flava, *Linn.* Syst. Nat. i. ii. p. 576, 11. Ed. Gmel. i. iv. p. 1639, 11. *Fabr.* Syst. El. i. p. 403, 91. Ent. Syst. i. p. 303, 51. Mant. Ins. i. p. 65, 42. Spec. Ins. i. p. 113, 35. Syst. Ent. p. 92, 21. *Herbst.* Naturs. viii. p. 299, 66, t. cxxxiv. f. 10. *Oliv.* Encycl. Ins. v. p. 388, 58. Entom. vi. p. 951, 46, 97, t. v. f. 75. *Schœnh.* Syn. Ins. ii. p. 225, 118.
Cassida cinerea, *Voet.* Col. ii. p. xliv. f. 8.
Patr. Surinam, Bolivia, Brasilia.

9. OMOPLATA HYALINA.

Bohem. Mon. Cassid. ii. p. 111, 8.
Echoma hyalina, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia, Buenos Ayres, Chiquitos.

10. Omoplata Solieri.

Bohem. Mon. Cassid. ii. p. 112, 9.
Echoma Solieri, *Dej.* Cat. ed. 3, p. 394.
Patr. Bahia.

11. Omoplata collata. B.M.

Bohem. Mon. Cassid. ii. p. 113, 10.
Patr. Brasilia.

12. Omoplata decipiens.

Bohem. Mon. Cassid. ii. p. 115, 11.
Echoma decipiens, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

13. Omoplata jugata.

Bohem. Mon. Cassid. ii. p. 116, 12.
Patr. Brasilia.

14. Omoplata trichroa.

Bohem. Mon. Cassid. ii. p. 117, 13.
Patr. St. Cruz, Boliviæ.

15. Omoplata pallidipennis. B.M.

Bohem. Mon. Cassid. ii. p. 118, 14.
Echoma pallidipennis, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia (Maj.-Gen. *Hardwicke*).

16. Omoplata sublucida.

Bohem. Mon. Cassid. ii. p. 119, 15.
Patr. Brasilia.

17. Omoplata tricolorata.

Bohem. Mon. Cassid. ii. p. 119, 16.
Patr. Brasilia.

18. Omoplata nigrolineata.

Bohem. Mon. Cassid. ii. p. 120, 17.
Patr. Brasilia.

19. Omoplata innocens.

Bohem. Mon. Cassid. ii. p. 121, 18.
Patr. Brasilia.

II. *Prothorax semicircularis, lateribus rotundatus.*

20. OMOPLATA SEMILINEATA.

Bohem. Mon. Cassid. ii. p. 122, 19.
Patr. Bahia, Brasiliæ.

21. OMOPLATA FLAVATA.

Bohem. Mon. Cassid. ii. p. 123, 20.
Patr. Brasilia.

22. OMOPLATA AXILLARIS. B.M.

Bohem. Mon. Cassid. ii. p. 124, 21, t. iv. E.
Cassida axillaris, *Sahlb.* Ins. Spec. p. 57, 38, t. iii. f. 4.
Cassida basalis, *Germ.* Ins. Spec. i. p. 537, 726.
Echoma axillaris, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

23. OMOPLATA SCAPULOSA.

Bohem. Mon. Cassid. ii. p. 125, 22.
Patr. Brasilia.

24. OMOPLATA SCAPULARIS. B.M.

Bohem. Mon. Cassid. ii. p. 127, 24.
Patr. Bolivia (Coll. *Bridges*).

25. OMOPLATA HÆMATIDEA.

Bohem. Mon. Cassid. ii. p. 128, 25.
Patr. Brasilia.

26. OMOPLATA LINEIGERA.

Bohem. Mon. Cassid. ii. p. 129, 26.
Patr. Brasilia.

27. OMOPLATA SQUALIDA.

Bohem. Mon. Cassid. ii. p. 130, 27.
Patr. Brasilia.

28. OMOPLATA BRUNNEOSIGNATA.

Bohem. Mon. Cassid. ii. p. 131, 28.
Patr. Brasilia.

29. OMOPLATA HEMISPILA.

Bohem. Mon. Cassid. ii. p. 132, 29.
Patr. Brasilia.

30. Omoplata sobrina.

Bohem. Mon. Cassid. ii. p. 133, 30.
Echoma sobrina, *Dej.* Cat. ed. 3, p. 394.
Patr. Bahia, Brasiliæ.

31. Omoplata trivialis.

Bohem. Mon. Cassid. ii. p. 134, 31.
Patr. Brasilia.

32. Omoplata puella.

Bohem. Mon. Cassid. ii. p. 135, 32.
Patr. Brasilia.

33. Omoplata albiventris.

Bohem. Mon. Cassid. ii. p. 136, 33.
Patr. Mexico.

34. Omoplata maculicollis.

Bohem. Mon. Cassid. ii. p. 137, 34.
Echoma maculicollis, *Dej.* Cat. ed. 3, p. 395.
Patr. Mexico.

35. Omoplata calcarata.

Bohem. Mon. Cassid. ii. p. 138, 35.
Patr. Mexico.

III. *Prothorax angustior; lateribus ultra medium oblique ampliatus, dein rectus.*

36. Omoplata marginata. B.M.

Bohem. Mon. Cassid. ii. p. 139, 36.
Cassida marginata, *Linn.* Syst. Nat. i. ii. p. 578, 23. Ed. Gmel. i. iv. p. 1642, 23. *Fabr.* Syst. El. i. p. 400, 76. Ent. Syst. i. p. 300, 39. Mant. Ins. i. p. 64, 31. Spec. Ins. i. p. 111, 25. Syst. Ent. p. 91, 16. *Herbst.* Naturs. viii. p. 325, 87, t. cxxx. f. 2. Archiv. p. 166, 11, t. 45, f. 1. *Oliv.* Encycl. Ins. v. p. 385, 42. Entom. vi. p. 941, 29, 97, t. i. f. 11. *De Geer*, Ins. v. p. 185, 10. *Schultz*, in Naturf. ix. p. 103, t. ii. f. 3. *Fuesly.* Archiv. Ins. t. xlv. f. 1. *Schœnh.* Syn. Ins. ii. p. 222, 101.
Cassida clypeata, *Voet.* Col. ii. p. xlii. f. 5.
Echoma marginata, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia, Cayenna, Surinam, Guyana.

37. OMOPLATA DISTINCTA.

Bohem. Mon. Cassid. ii. p. 141, 37.
Echoma distincta, *Dej.* Cat. ed. 3, p. 395.
Patr. Insulæ Guadeloupe et St. Martinique.

38. OMOPLATA BONFILSII.

Bohem. Mon. Cassid. ii. p. 141, 38.
Echoma Bonfilsii, *Dej.* Cat. ed. 3, p. 395.
Patr. Peruvia.

39. OMOPLATA BIPLAGOSA.

Bohem. Mon. Cassid. ii. p. 142, 39.
Patr. Cayenna.

40. OMOPLATA IRRORATA. B.M.

Bohem. Mon. Cassid. ii. p. 143, 40.
Cassida irrorata, *Fabr.* Syst. El. i. p. 406, 107. *Voet.* Col. ii. t. xli. f. 1. *Schœnh.* Syn. Ins. ii. p. 227, 135.
Echoma irrorata, *Dej.* Cat. ed. 3, p. 395.
Patr. Cayenna.

Genus 24. OMASPIDES.

Chevrol. Dej. Cat. ed. 3, p. 395. *Hope,* Col. Man. iii. p. 158. *Bohem.* Mon. Cassid. ii. p. 144. *Erichs.* Archiv. f. Naturg. xiii. p. 153.
Cassida, *Linn.* Syst. Nat. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *De Geer,* Mem. v. p. 179. *Herbst.* Naturs. viii. p. 204. *Oliv.* Entom. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209.

1. OMASPIDES CLATHRATA. B.M.

Bohem. Mon. Cassid. ii. p. 146, 1, t. iv. F.
Cassida clathrata, *Linn.* Syst. Nat. i. ii. p. 577, 18. Ed. Gmel. i. iv. p. 1641, 18. *De Geer,* Ins. v. p. 179, 3, t. xv. f. 7. *Voet.* Col. ii. t. xli. f. 3. *Oliv.* Encycl. v. p. 390, 66. Entom. vi. p. 930, 9, t. iv. f. 55.
Cassida transversa, *Fabr.* Syst. El. i. p. 404, 93. Suppl. p. 84, 52, 53. *Herbst.* Naturs. viii. p. 298, 65, t. cxxxiv. f. 9. *Schœnh.* Syn. Ins. ii. p. 223, 103.
Cassida quadripustulata, *Fabr.* Syst. El. i. p. 401, 77. Entom. Syst. i. p. 301, 40. Mant. Ins. i. p. 64, 32. Spec. Ins. i. p. 111, 26. *Linn.* Syst. Nat. viii. p. 341, 108. *Schœnh.* Syn. Ins. ii. p. 222, 102.
Omaspides transversa, *Dej.* Cat. ed. 3, p. 395.
Patr. Cayenna, Santarem (Coll. *H. W. Bates*).

2. Omaspides trifasciata.

Bohem. Mon. Cassid. ii. p. 147, 2. *Dej.* Cat. ed. 3, p. 395.
Cassida trifasciata, *Fabr.* Syst. El. i. p. 404, 97. Ent. Syst. i. p. 304, 55. Mant. Ins. p. 65, 46. *Herbst.* Naturs. viii. p. 290, 59. *Linn.* Syst. Nat. ed. Gmel. i. iv. p. 1640, 60. *Oliv.* Encycl. Ins. v. p. 389, 62. Entom. vi. p. 929, 6, 97, t. ii. f. 28. *Schœnh.* Syn. Ins. ii. p. 226, 124.
Patr. Cayenna.

3. Omaspides augusta.

Rotundata, modice convexa, obscure cærulea, sub-opaca; antennis nigris; prothorace subtilissime punctulato, ante lobum transversim impresso, medio tenuiter, breviter canaliculato; elytris dorso subtiliter, irregulariter reticulatis, reticulo tenui parum elevato, areolas parvas includente; margine late explanato, plaga maxima oblonga, dilute sanguinea, maculis plurimis cærulescentibus notata. Long. 14, lat. 13¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Campo del Sacramento, Dom. *Castelnau*, Mus. Paris.

4. Omaspides basilica. B.M.

Rotundata, modice convexa, obscure cærulea, parum nitida; antennis basi subtus ferrugineis; prothorace subtilissime, sub-remote punctulato, medio tenuiter canaliculato; elytris subtiliter, sat crebre punctulatis, singulo in dorso pone medium macula parva, elongata plagaque magna oblongo-ovata, intramarginali, dilute sanguineis, hac medio maculis tribus longitudinaliter positis, cærulescentibus notata. Long. 14, lat. 13¼ millim.

Bohem. Mon. Cassid. Append.
Patr. Ega, Brasiliæ, Mus. Britann. (Coll. *H. W. Bates*).

5. Omaspides lunulata.

Bohem. Mon. Cassid. ii. p. 148, 3.
Patr. Para.

6. Omaspides specularis.

Rotundata, parum convexa, nigra, nitida, supra obscure ænea; antennis articulis secundo toto, tertio quartoque subtus testaceis; prothorace triangulari, subtiliter, minus crebre punctulato; elytris vage, sat crebre punctulatis, plaga magna oblonga, intramarginali, flava. Long. 12, lat. 11½ millim.

Erichs. Archiv. f. Naturg. viii. p. 153, 1. *Bohem.* Mon. Cassid. ii. p. 149, 4. *Bohem.* Mon. Cassid. Append.
Patr. Peruvia, Mus. Berol.

TRIBUS III.

Genus 25. BATONOTA.

Hope Ann. Nat. Hist. iii. p. 98, t. iv. f. 6. *Bohem.* Mon. Cassid. ii. p. 153.
Cassida, *Fabr.* Syst. El. i. p. 387. *Herbst.* Naturs. viii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209. *Germ.* Ins. Spec. i. p. 536.
Dorynota, *Chevrolat,* Dej. Cat. ed. 3, p. 394.
Omoteina, *Chevrolat,* Dej. Cat. ed. 3, p. 398.

I. *Elytra ante medium spina erecta armata.*

1. BATONOTA BIDENS. B.M.

Bohem. Mon. Cassid. ii. p. 155, 1.
Cassida bidens, *Fabr.* Syst. El. i. p. 403, 88. Ent. Syst. i. p. 302, 48. Mant. Ins. i. p. 65, 39. Spec. Ins. i. p. 112, 32. *Linn.* Syst. Nat. ed. Gmel. i. iv. p. 1638, 49. *Herbst.* Naturs. viii. p. 345, 113. *Oliv.* Ent. vi. p. 934, 15, 97, t. i. f. 10. *Schœnh.* Syn. Ins. ii. p. 225, 115.
Cassida atra, *Oliv.* Enc. meth. Ins. v. p. 387, 53.
Dorynota bidens, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

2. BATONOTA NIGRA.

Sub-triangularis, convexa, nigra, parum nitida, brevissime parce albido-pubescens; antennis basi testaceis; prothorace subtiliter punctulato, extus anguste testaceo marginato; elytris nigro-æneis, subtilissime coriaceis, mediocriter, irregulariter punctatis, disco antico bi-carinulatis, dorso spina mediocriter longa instructis; humeris lateraliter extensis, supra carinatis; margine extus et ultra medium subtilius punctulato. Long. 9½, lat. 9 millim.

Bohem. Mon. Cassid. Append.
Var. a. corpore subtus testaceo-variegato, prothorace evidentius testaceo-marginato, elytris cærulescenti-nigris.
Patr. Peruvia, Dom. *Deyrolle,* Mus. Holm. *Var. a.* Caraccas, Mus. Dom. *Sallé.*

3. BATONOTA MONOCEROS. B.M.

Bohem. Mon. Cassid. ii. p. 156, 2.
Cassida Monoceros, *Germ.* Ins. Spec. i. p. 536, 724.
Dorynota militaris, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

4. Batonota Ballista. B.M.

Bohem. Mon. Cassid. ii. p. 157, 3.
Patr. Brasilia.

5. Batonota pugionata. B.M.

Bohem. Mon. Cassid. ii. p. 158, 4.
Cassida pugionata, *Hoffmansegg.* Germ. Ins. Spec. i. p. 537, 725.
Dorynota armata, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia (St. Paul).

6. Batonota Gladiator.

Sub-triangularis, dilute ferruginea, sub-opaca; antennis articulis 6 ultimis tibiisque extus nigris; prothorace sub-lævi; elytris crebre, vage punctatis, dorso punctato-striatis, limbo antice et postice maculisque nonnullis disci nigris, basi porrectis, ante medium spina longa erecta, suturali; humeris lateraliter extensis, apice sub-obtusis. Long. 9, lat. 8½ millim.

Bohem. Mon. Cassid. Append.
Patr. Guayra, Dom. *Deyrolle,* Mus. Holm.

7. Batonota bellicosa.

Bohem. Mon. Cassid. ii. p. 159, 5.
Patr. Brasilia.

8. Batonota nodosa. B.M.

Bohem. Mon. Cassid. ii. p. 160, 6.
Patr. Columbia.

9. Batonota pugnax.

Bohem. Mon. Cassid. ii. p. 161, 7.
Patr. Columbia.

10. Batonota cornigera.

Bohem. Mon. Cassid. ii. p. 162, 8.
Dorynota cornigera, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia.

11. Batonota mucronata.

Bohem. Mon. Cassid. ii. p. 164, 9.
Dorynota Monocera, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia, Bolivia.

12. Batonota parallela.

Bohem. Mon. Cassid. ii. p. 165, 10.
Dorynota parallela, *Blanchard*, Voy. d'Orbigny, Ins. pl. 24, f. 6.
Patr. Brasilia.

13. Batonota ensifera.

Bohem. Mon. Cassid. ii. p. 166, 11.
Patr. Corrientes.

14. Batonota gregaria. B.M.

Bohem. Mon. Cassid. ii. p. 167, 12.
Dorynota gregaria, *Dej.* Cat. ed. 3, p. 394.
Patr. Brasilia (St. Paul).

15. Batonota spinosa.

Bohem. Mon. Cassid. ii. p. 168, 13.
Patr. Brasilia, Bolivia, Chiquitos.

16. Batonota apiculata. B.M.

Bohem. Mon. Cassid. ii. p. 169, 14.
Patr. Brasilia (St. Paul).

17. Batonota aculeata. B.M.

Bohem. Mon. Cassid. ii. p. 170, 15.
Dorynota aculeata, *Dej.* Cat. ed. 3, p. 394.
Patr. St. Domingo.

18. Batonota turrifera.

Bohem. Mon. Cassid. ii. p. 171, 16.
Patr. St. Domingo.

II. *Elytra basi retusa, ante medium gibbosa.*

19. Batonota funesta.

Bohem. Mon. Cassid. ii. p. 172, 17.
Patr. Cayenna.

20. Batonota truncata. B.M.

Bohem. Mon. Cassid. ii. p. 173, 18, t. v. B.
Cassida truncata, *Fabr.* Syst. El. i. p. 403, 90. Ent. Syst. i. p. 303, 50. Mant. Ins. i. p. 65, 41. Spec. Ins. i. p. 112, 34. *Linn.* Syst. Nat. ed. Gmel. i. iv. p. 1638, 47. *Herbst.* Naturs. viii. p. 284, 56, t. cxxxiv. f. 2. *Oliv.* Enc. meth. Ins. v.

p. 388, 57. Entom. vi. p. 933, 13, 97, t. i. f. 9. *Schœnh.* Syn. Ins. ii. p. 225, 117.
Dorynota truncata, *Dej.* Cat. ed. 3, p. 394.
Patr. Cayenna, Columbia.

21. Batonota insidiosa. B.M.

Bohem. Mon. Cassid. ii. p. 174, 19.
Patr. Columbia, Guatemala.

22. Batonota tenebrosa. B.M.

Bohem. Mon. Cassid. ii. p. 175, 20.
Patr. Chiquitos. Var. Villa Nova, Amazon (Coll. *H. W. Bates*).

23. Batonota exaltata.

Bohem. Mon. Cassid. ii. p. 176, 21.
Cassida exaltata, *Fabr.* Syst. El. i. p. 402, 83. *Schœnh.* Syn. Ins. ii. p. 224, 110.
Patr. America meridionalis.

24. Batonota Dejeanii.

Bohem. Mon. Cassid. ii. p. 178, 22.
Dorynota Dejeanii, *Dej.* Cat. ed. 3, p. 394.
Patr. Cayenna.

25. Batonota bivittipennis. B.M.

Sub-triangularis, convexa, nigra, supra parum, subtus magis nitida; prothorace subtiliter, sat crebre punctulato, tenuiter carinato; elytris viridibus, mediocriter, irregulariter punctatis, ad suturam bi-seriatim punctato-striatis, dorso reticulatis, reticulo modice elevato; basi retusis, transversim gibbosis; margine late explanato, rufo-testaceo, parce æneo-punctato, limbo toto viridi. Long. 14½, lat. 14 millim.

Bohem. Mon. Cassid. Append.
Patr. Ega, Brasiliæ, Mus. Britann. (Coll. *H. W. Bates*). Prope fluv. Amazonum, Mus. Dom. *Deyrolle.*

26. Batonota Kiesenwetteri.

Bohem. Mon. Cassid. ii. p. 179, 23.
Patr. Columbia.

27. Batonota Kunzei.

Bohem. Mon. Cassid. ii. p. 180, 24.
Patr. In confiniis fluminis Amazonum.

28. Batonota viridisignata.

Bohem. Mon. Cassid. ii. p. 181, 25.
Patr. Bolivia, Buenos Ayres.

29. Batonota illudens.

Bohem. Mon. Cassid. ii. p. 182, 26.
Patr. Brasilia.

30. Batonota Lerouxii.

Bohem. Mon. Cassid. ii. p. 183, 27.
Patr. Insula Cuba.

31. Batonota humeralis. B.M.

Bohem. Mon. Cassid. ii. p. 184, 28.
Cassida humeralis, *Oliv.* Entom. vi. p. 964, 70, 97, t. v. f. 88.
Omoteina humeralis, *Dej.* Cat. ed. 3, p. 398.
Patr. Insula St. Domingo.

Genus 26. BASIPTA.

Chevrolat, Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 185.

1. Basipta stolida. B.M.

Bohem. Mon. Cassid. ii. p. 186, 1, t. v. C.
Patr. Caffraria, Caput Bonæ Spei.

2. Basipta pallens.

Klug, Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 187, 2.
Patr. Caput Bonæ Spei.

3. Basipta luteocincta.

Bohem. Mon. Cassid. ii. p. 188, 3.
Patr. Caput Bonæ Spei.

4. Basipta pilosella.

Bohem. Mon. Cassid. ii. p. 189, 4.
Patr. Caffraria.

Genus 27. PHYSONOTA.

Chevrolat, Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 190.
Echoma, *Dej.* Cat. ed. 3, p. 394 (ex parte).
Deloyala, *Dej.* Cat. ed. 3, p. 395 (ex parte).

I. Corpus ovatum, plus minusve convexum.

a. *Elytra apice late rotundata.*

* *Prothorax nigro- vel ferrugineo-notatus.*

1. PHYSONOTA ALUTACEA. B.M.

Klug, Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 191, 1.
Patr. Mexico (Vera Cruz), Columbia, Venezuela (Coll. *Dyson*).

2. PHYSONOTA CYRTODES. B.M.

Bohem. Mon. Cassid. ii. p. 192, 2.
Patr. Mexico, Columbia, Guatemala (Coll. *Scherzer*).

3. PHYSONOTA LUTARELLA. B.M.

Sub-rotundata, valde convexa, flava, nitida, albido-marginata; antennis extrorsum nigris; prothorace lævi, ante basin maculis tribus parvis, obsoletis, brunneis; scutello nigro; elytris dorso minus crebre, mediocriter, margine magis explanato remotius, profundius punctatis; pone basin valide, obtuse gibbosis; corpore subtus nigro-variegato; femoribus supra medio macula magna, nigra, Long. 15, lat. 12½ millim.

Bohem. Mon. Cassid. Append.
Patr. Caraccas, Mus. Britann.

4. PHYSONOTA PICTICOLLIS. B.M.

Bohem. Mon. Cassid. ii. p. 193, 3.
Patr. Mexico.

5. PHYSONOTA VITTICOLLIS.

Bohem. Mon. Cassid. ii. p. 194, 4.
Patr. Mexico.

6. PHYSONOTA OVALIS. B.M.

Bohem. Mon. Cassid. ii. p. 195, 5.
Patr. Mexico.

7. PHYSONOTA HELIANTHI. B.M.

Bohem. Mon. Cassid. ii. p. 196, 6.
Cassida Helianthi, *Randall*, Bost. Journ. of Nat. Hist. ii. p. 30, 48.
Patr. Maine, America borealis.

8. Physonota turgida.

Bohem. Mon. Cassid. ii. p. 197, 7.
Patr. Mexico.

9. Physonota bicostata.

Bohem. Mon. Cassid. ii. p. 198, 8.
Patr. Brasilia.

10. Physonota gigantea. B.M.

Bohem. Mon. Cassid. ii. p. 199, 9, t. v. D.
Patr. Guatemala, Chili, Valparaiso, W. Indies (*C. D. E. Fortnum, Esq.*).

11. Physonota Cumingii.

Bohem. Mon. Cassid. ii. p. 201, 10.
Patr. Chili.

12. Physonota Spencei.

Bohem. Mon. Cassid. ii. p. 202, 11.
Patr. Chili.

13. Physonota breviuscula.

Rotundata, modice convexa, supra sordide flavo-testacea, nitida, subtus nigra; antennis articulis 1 apice, 2-5 subtus flavescentibus; prothorace lævi, medio linea brevi, antice magis, postice minus abbreviata scutelloque nigris; elytris remote, parum profunde, disco exteriore minus regulariter, seriatim punctatis; margine late explanato, vage, minus crebre punctato, ante medium transversim impresso. Long. 12, lat. 10¾ millim.

Bohem. Mon. Cassid. Append.
Patr. — — — Mus. Dom. *Deyrolle.*

14. Physonota incrustata. B.M.

Bohem. Mon. Cassid. ii. p. 203, 12.
Patr. Mexico.

15. Physonota stigmatilis.

Bohem. Mon. Cassid. ii. p. 204, 13.
Patr. Mexico.

16. Physonota separata.

Bohem. Mon. Cassid. ii. p. 205, 14.
Patr. Mexico.

17. Physonota Mexicana.

Bohem. Mon. Cassid. ii. p. 206, 15.
Deloyala Mexicana, *Dej.* Cat. ed. 3, p. 395.
Patr. Mexico.

18. Physonota disjuncta.

Bohem. Mon. Cassid. ii. p. 207, 16.
Cassida disjuncta, *Chevrol.* Coleopt. de Mexique, fasc. iii.
Patr. Mexico.

19. Physonota fuscata. B.M.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 208, 17.
Cassida fuscata, *Klug*, Preis-Verzeichn. 1829, p. 8, 96.
Var. Physonota quadrilineata, *Dej.* Cat. ed. 3, p. 398.
Patr. Brasilia.

20. Physonota atomaria.

Bohem. Mon. Cassid. ii. p. 209, 18.
Patr. Brasilia.

** *Prothorax immaculatus.*

21. Physonota flavago.

Bohem. Mon. Cassid. ii. p. 210, 19.
Patr. Mexico.

22. Physonota languida.

Bohem. Mon. Cassid. ii. p. 211, 20.
Patr. Brasilia.

23. Physonota rubronotata.

Bohem. Mon. Cassid. ii. p. 212, 21.
Patr. Brasilia.

24. Physonota punctulata.

Cassida punctulata, *Klug*, Preis-Verzeichn. 1829, p. 8, 98.
Chelymorpha punctulata, *Dej.* Cat. ed. 3, p. 394.
Pœcilaspis punctulata, *Bohem.* Mon. Cassid. i. p. 436, 48.
Patr. Brasilia.

25. Physonota testudinaria.

Pœcilaspis testudinaria, *Bohem.* Mon. Cassid. i. p. 437, 49.
Patr. Buenos Ayres.

26. PHYSONOTA EXARATA. B.M.

Bohem. Mon. Cassid. ii. p. 213, 22.
Patr. Brasilia.

27. PHYSONOTA PROLIXA.

Bohem. Mon. Cassid. ii. p. 214, 23.
Patr. Buenos Ayres.

28. PHYSONOTA SULCIPENNIS.

Bohem. Mon. Cassid. ii. p. 215, 24.
Patr. Brasilia, Montevideo.

29. PHYSONOTA INGRATA.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 216, 25.
Patr. Tucuman.

30. PHYSONOTA CANDIDA.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 217, 26.
Patr. Brasilia.

31. PHYSONOTA NIGRITARSIS.

Bohem. Mon. Cassid. ii. p. 218, 27.
Patr. Chiquitos.

32. PHYSONOTA CITRINELLA.

Bohem. Mon. Cassid. ii. p. 219, 28.
Patr. Mexico.

33. PHYSONOTA OVATA.

Bohem. Mon. Cassid. ii. p. 219, 29.
Deloyala ovata, *Dej.* Cat. ed. 3, p. 395.
Patr. Mexico.

b. *Elytra apice attenuata.*

* *Prothorax nigro-notatus.*

34. PHYSONOTA MACULIPES.

Bohem. Mon. Cassid. ii. p. 220, 30.
Patr. Bogota.

35. PHYSONOTA CAUDATA. B.M.

Bohem. Mon. Cassid. ii. p. 222, 31.
Deloyala caudata, *Dej.* Cat. ed. 3. p. 395.
Patr. Mexico.

36. Physonota attenuata.

Bohem. Mon. Cassid. ii. p. 223, 32.
Patr. Mexico.

37 Physonota plana.

Bohem. Mon. Cassid. ii. p. 224, 33.
Patr. Bogota, Columbiæ.

38. Physonota humilis.

Ovata, parum convexa, dilute flavo-testacea, nitida, albido-marginata; antennis nigris, articulis 2-5 subtus testaceis; prothorace lævi, ante basin lineola brevi, longitudinali, nigra; elytris subtiliter, sub-remote, irregulariter ad suturam sub-seriatim punctatis; prosterno utrinque, pectore medio, margine subtus basi macula parva triangulari suturaque apice nigris. Long. 11¼, lat. 7½ millim.

Bohem. Mon. Cassid. Append.
Patr. Mexico, Dom. *Lacordaire*, Mus. Holm.

** *Prothorax immaculatus.*

39. Physonota translucida.

Bohem. Mon. Cassid. ii. p. 225, 34.
Patr. Mexico.

40. Physonota flaveola. B.M.

Bohem. Mon. Cassid. ii. p. 226, 35.
Patr. Mexico.

II. Corpus rotundatum.

a. *Elytra non gibbosa.*

** *Tibiæ extus apice calcaratæ.*

41. Physonota calcarata.

Bohem. Mon. Cassid. ii. p. 227, 36.
Patr. Mexico.

42. Physonota spinipes.

Bohem. Mon. Cassid. ii. p. 228, 37.
Patr. Mexico ad Oaxaca.

** *Tibiæ extus apice muticæ.*

43. Physonota maculiventris.

Bohem. Mon. Cassid. ii. p. 229, 38.

Echoma maculiventris, *Dej.* Cat. ed. 3, p. 395.
Patr. Mexico.

b. *Elytra dorso antico gibbosa.*

44. PHYSONOTA NITIDICOLLIS.

Bohem. Mon. Cassid. ii. p. 231, 39.
Patr. Mexico.

45. PHYSONOTA CITRINA. B.M.

Bohem. Mon. Cassid. ii. p. 231, 40.
Patr. Mexico.

46. PHYSONOTA CEREA.

Bohem. Mon. Cassid. ii. p. 232, 41.
Patr. Mexico ad Oaxaca.

47. PHYSONOTA XERAMPELINA.

Bohem. Mon. Cassid. ii. p. 233, 42.
Patr. Brasilia.

48. PHYSONOTA DUBIA.

Bohem. Mon. Cassid. ii. p. 235, 43.
Patr. Brasilia.

Genus 28. EURYPEPLA.

Bohem. Mon. Cassid. ii. p. 236.
Cassida, *Fabr.* Syst. El. i. p. 345. *Schœnh.* Syn. Ins. ii. p. 134.

1. EURYPEPLA JAMAICENSIS. B.M.

Bohem. Mon. Cassid. ii. p. 237, 1, t. v. E.
Cassida jamaicensis, *Linn.* Syst. Nat. i. ii. p. 577, 21. Ed. Gmel. i. iv. p. 1641, 21. *Fabr.* Syst. El. i. p. 401, 81. Ent. Syst. i. p. 301, 42. Mant. Ins. i. p. 64, 34. Spec. Ins. i. p. 111, 27. Syst. Ent. p. 91, 17. *Sloan.* Hist. Jamaic. ii. p. 208, t. ccxxiii. f. 27, 28. *Schœnh.* Syn. Ins. ii. p. 223, 108.
Patr. Insul. Americæ meridionalis.

2. EURYPEPLA BREVILINEATA. B.M.

Bohem. Mon. Cassid. ii. p. 238, 2.
Patr. Campeachy, Yucatan.

3. Eurypepla vitrea.

Bohem. Mon. Cassid. ii. p. 239, 3.
Patr. Cuba.

Genus 29. PLATYCYCLA.

Bohem. Mon. Cassid. ii. p. 240.
Deloyala Chevrolat, *Dej.* Cat. ed. 3, p. 395.

1. Platycycla deruta. B.M.

Bohem. Mon. Cassid. ii. p. 241, 1, t. v. F.
Deloyala deruta, *Dej.* Cat. ed. 3, p. 395.
Patr. Mexico.

Genus 30. ASPIDOMORPHA.

Hope, Coleopt. Man. iii. p. 158. *Bohem.* Mon. Cassid. ii. p. 242.
Cassida, *Linn.* Syst. Nat. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *De Geer*, Ins. v. p. 178. *Herbst.* Naturs. viii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209. *Dej.* Cat. ed. 3, p. 374.
Deloyala, *Chevrolat*, Dej. Cat. ed. 3, p. 395.

I. *Elytra antice non gibbosa.*

1. Aspidomorpha chlorotica.

Bohem. Mon. Cassid. ii. p. 244, 1.
Cassida chlorotica, *Oliv.* Ent. vi. p. 931, 10, 97, t. iv. f. 56.
Patr. Guinea.

2. Aspidomorpha spectabilis.

Bohem. Mon. Cassid. ii. p. 245, 2, t. vi. A.
Patr. Guinea.

3. Aspidomorpha puncticosta. B.M.

Bohem. Mon. Cassid. ii. p. 246, 3.
Patr. Caffraria (*Dr. A. Smith*), Port Natal.

4. Aspidomorpha Palleago. B.M.

Bohem. Mon. Cassid. ii. p. 247, 4.
Patr. Caffraria (*Earl of Derby*).

5. ASPIDOMORPHA INDISTINCTA. B.M.

Bohem. Mon. Cassid. ii. p. 248, 5.
Deloyala indistincta, *Dej.* Cat. ed. 3, p. 395.
Patr. Senegallia, Gambia (*Ent. Club*).

6. ASPIDOMORPHA INNOCUA. B.M.

Bohem. Mon. Cassid. ii. p. 249, 6.
Patr. Senegallia.

7. ASPIDOMORPHA QUINQUEFASCIATA. B.M.

Bohem. Mon. Cassid. ii. p. 250, 7.
Cassida quinquefasciata, *Fabr.* Syst. El. i. p. 401, 79. *Schœnh.* Syn. Ins. ii. p. 309, 105. *Klug,* Ermann Reise Ins. p. 47, 174.
Cassida pectoralis? *Oliv.* Ent. vi. p. 960, 62, t. v. f. 82.
Deloyala quadriremis, *Dej.* Cat. ed. 3, p. 395.
Patr. Senegallia, Gambia (*Ent. Club*), Congo (*A. Curror, Esq.*, R.N.)

8. ASPIDOMORPHA CINCTA. B.M.

Bohem. Mon. Cassid. ii. p. 251, 8.
Cassida cincta, *Fabr.* Syst. El. i. p. 392, 22. Ent. Syst. i. p. 295, 13. Mant. Ins. i. p. 63, 12. Spec. Ins. i. p. 109, 9. *Linn.* Syst. Nat. Gmel. i. iv. p. 1637, 41. *Herbst.* Naturs. viii. p. 331, 92. *Schœnh.* Syn. Ins. ii. p. 216, 30. *Klug,* Ermann Reise Ins. p. 47, 173.
Cassida quadriremis, *Schœnh.* Syn. Ins. ii. p. 223, 107.
Deloyala quadriremis, *Dej.* Cat. ed. 3, p. 395.
Deloyala consanguinea, *Dej.* Cat. ed. 3, p. 395.
Patr. Senegallia, Guinea (Fort Accra).

9. ASPIDOMORPHA ISPARETTA. B.M.

Bohem. Mon. Cassid. ii. p. 252, 9.
Patr. Senegallia, Guinea (Whydah), Coll. *Fraser.*

10. ASPIDOMORPHA WAHLBERGI.

Bohem. Mon. Cassid. ii. p. 253, 10.
Patr. Caffraria.

11. ASPIDOMORPHA WESTWOODII. B.M.

Bohem. Mon. Cassid. ii. p. 254, 11.
Patr. Nova Hollandia (Clarence River).

12. ASPIDOMORPHA ORBICULARIS. B.M.

Bohem. Mon. Cassid. ii. p. 255, 12.
Patr. Phil. Ins. Manilla (Coll. *Cuming*), India orientalis.

13. ASPIDOMORPHA CONFINIS. B.M.

Bohem. Mon. Cassid. ii. p. 256, 13.
Cassida confinis, *Dej.* Cat. ed. 3, p. 398. *Klug*, Ermann Reise Ins. v. p. 47, 181.
Patr. Caffraria, Senegallia, Guinea (*Ent. Club*).

14. ASPIDOMORPHA APICALIS.

Bohem. Mon. Cassid. ii. p. 257, 14.
Cassida apicalis, *Klug*, Ins. von Madag. p. 122, 198.
Patr. Madagascar.

15. ASPIDOMORPHA VIRIDIDORSATA.

Bohem. Mon. Cassid. ii. p. 258, 15.
Patr. Guinea.

16. ASPIDOMORPHA CHLORINA. B.M.

Bohem. Mon. Cassid. ii. p. 259, 16.
Patr. Africa meridionalis, et A. occidentalis (Coll. *Fraser*).

17. ASPIDOMORPHA AMPLISSIMA. B.M.

Bohem. Mon. Cassid. ii. p. 260, 17.
Patr. Manilla, in Ins. Philippinis (Coll. *Cuming*).

18. ASPIDOMORPHA MILIARIS. B.M.

Bohem. Mon. Cassid. ii. p. 261, 18. *Hombr.* et *Jaquin.* Voy. d'Urville, t. xviii. f. 9, var.
Cassida miliaris, *Fabr.* Syst. El. i. p. 400, 71. Ent. Syst. i. p. 300, 36. Mant. Ins. i. p. 64, 29. Spec. Ins. i. p. 111, 24. Syst. Ent. p. 91, 15. *Linn.* Syst. Nat. Gmel. i. iv. p. 1640, 58. *Oliv.* Enc. meth. v. p. 385, 40. Entom. vi. p. 943, 33, 97, t. ii. f. 25. *Schœnh.* Syn. Ins. ii. p. 222, 96.
Deloyala miliaris, *Dej.* Cat. ed. 3, p. 395.
Var. c. elytris dorso plaga nigra, vittam dorsalem communem flavam gerente utrinque ramulos duos concolores emittente, unum basalem, postice ampliatum limbum macula parva flava notatum, alterum infra medium sat latum.
Var. d. elytris nigris, sutura ultra medium anguste, macula parva disci, altera majore transversa humerali, plaga laterali media sinuata introrsum maculaque ante apicem flavis.
Var. e. elytris nigris, macula sub-triangulari pone scutellum

plagaque magna laterali in medio antrorsum arcuatim ducta flavis.

Var. f. flavo-testacea, elytris dorso plaga maxima nigra, utrinque ramulos duos in marginem emittente, anteriorem latiorem, extrorsum postice valde rotundato-ampliatum, macula ovata humerali, flava notatum, sutura ultra medium anguste vittaque longitudinali dorsi antici, undulata, flavo-testaceis.

Patr. India orientalis (*Walter Elliott*, Esq.), Java, Celebes, China, Hong-Kong (*J. C. Bowring*, Esq.), *Var. c.* et *e.* Insul. Philipp. Mus. Britann.

19. ASPIDOMORPHA WESTERMANNI.

Bohem. Mon. Cassid. ii. p. 262, 19.
Deloyala Westermanni, *Dej.* Cat. ed. 3, p. 395.
Patr. Guinea.

20. ASPIDOMORPHA QUADRIMACULATA. B.M.

Bohem. Mon. Cassid. ii. p. 263, 20.
Cassida quadrimaculata, *Oliv.* Ent. vi. p. 945, 36, t. iv. f. 68.
Dej. Cat. ed. 3, p. 374.
Cassida nigropunctata, *Dej.* Cat. ed. 2.
Patr. Senegallia, Gambia (*Ent. Club*). Congo (*A. Curror, Esq.*, R.N.).

21. ASPIDOMORPHA ORIENTALIS.

Rotundata (*mas*) vel sub-ovata (*femina*), modice convexa, flavo-testacea; antennis articulo ultimo, pectore medio abdomineque plaga magna nigris; prothorace lævi, antice rotundato, basi utrinque levissime sinuato; elytris regulariter, parum profunde, dorso remotius seriatim punctatis, maculis numerosis, inæqualibus, nigris notatis, bis disco exteriore plus minusve confluentibus; margine modice explanato, lævi, nigro-bimaculato. Long. $8\frac{1}{4}$-$9\frac{1}{2}$, lat. $6\frac{3}{4}$-7 millim.

Bohem. Mon. Cassid. Append.
Var. a. maculis elytrorum minoribus, haud confluentibus, pectore utrinque testaceo-marginato.
Var. b. ut var. *a.* sed macula marginali basali deficiente.
Var. c. antennis articulo ultimo haud infuscato, maculis elytrorum parvis paucioribus, margine immaculato; pectore utrinque testaceo-marginato.
Var. d. elytro singulo maculis quatuor parvis, nigris notato, margine immaculato; pectore utrinque testaceo-marginato.
Patr. India orientalis, Mus. Holm. a Dom. *Saunders*, communicata.

II. *Elytra basi leviter gibbosa.*

22. Aspidomorpha adhærens.

Bohem. Mon. Cassid. ii. p. 264, 21.
Cassida adhærens, *Fabr.* Syst. El. i. p. 400, 72. Weber. Observ. Ent. i. p. 51, 2. *Schœnh.* Syn. Ins. ii. p. 222, 97. *Boisd.* Faun. Ent. de l'Ocean. ii. p. 539, 5. *Hombr.* et *Jaquin.* Voy. d'Urville, t. xviii. f. 11.
Deloyala adhærens, *Dej.* Cat. ed. 3, p. 395.
Patr. Insulæ maris pacifici, Nova Guinea.

23. Aspidomorpha bimaculata.

Bohem. Mon. Cassid. ii. p. 266, 22.
Cassida bimaculata, *Fabr.* Syst. El. i. p. 400, 73. Ent. Syst. i. p. 300, 37. *Herbst.* Naturs. viii. p. 340, 106. *Schœnh.* Syn. Ins. ii. p. 222, 98.
Deloyala Troberti, *Dejean.*
Patr. Guinea, Bissagos.

24. Aspidomorpha frenata.

Bohem. Mon. Cassid. ii. p. 267, 23.
Cassida fenestrata, *Oliv. Beauv.* Ins. d'Afrique, p. 255, t. xxxviii. f. 8.
Patr. Guinea.

25. Aspidomorpha biguttata.

Bohem. Mon. Cassid. ii. p. 267, 24.
Cassida biguttata, *Fabr.* Syst. El. i. p. 399, 70. Ent. Syst. i. p. 299, 35. Mant. Ins. i. p. 64, 23. Spec. Ins. i. p. 111, 23. Syst. Ent. p. 91, 14. *Linn.* Syst. Nat. Gmel. i. iv. p. 1640, 57. *Herbst.* Naturs. viii. p. 340, 105. *Oliv.* Encycl. meth. v. p. 385, 39. Entomol. vi. p. 946, 38, 97, t. iii. f. 40. *Schœnh.* Syn. Ins. ii. p. 222, 95.
Patr. Guyana? Cayenna?

26. Aspidomorpha fenestrata. B.M.

Bohem. Mon. Cassid. ii. p. 269, 25.
Cassida fenestrata, *Oliv.* Ent. vi. p. 947, 39, 97, t. iv. f. 69.
Patr. Guinea.

27. Aspidomorpha galamensis.

Bohem. Mon. Cassid. ii. p. 270, 26.
Deloyala galamensis, *Buquet,* Dej. Cat. ed. 3, p. 395.
Patr. Guinea.

28. Aspidomorpha Schœnherri. B.M.

Bohem. Mon. Cassid. ii. p. 370, 27.
Deloyala Schœnherri, *Dej.* Cat. ed. 3, p. 395.
Patr. Sierra Leona (Rev. *D. F. Morgan*), Whydah (Guinea), (Coll. *Fraser*).

29. Aspidomorpha Afzelii. B.M.

Bohem. Mon. Cassid. ii. p. 372, 28.
Patr. Sierra Leona, Guinea, (Whydah, Coll. *Fraser*), "Georgia"?

30. Aspidomorpha funerea.

Bohem. Mon. Cassid. ii. p. 273, 29.
Patr. Sierra Leona.

31. Aspidomorpha stolata.

Bohem. Mon. Cassid. ii. p. 274, 30.
Patr. Guinea.

32. Aspidomorpha Madagascarica.

Bohem. Mon. Cassid. ii. p. 275, 31.
Patr. Madagascar.

33. Aspidomorpha tecta. B.M.

Bohem. Mon. Cassid. ii. p. 276, 32.
Deloyala quinquefasciata var. tecta, *Klug*, Dej. Cat. ed. 3, p. 395.
Patr. Caput Bonæ Spei, Caffraria (Dr. *A. Smith*). Port Natal, Guinea (Whydah), (Coll. *Fraser*).

34. Aspidomorpha silacea. B.M.

Bohem. Mon. Cassid. ii. p. 277, 33.
Patr. Caput Bonæ Spei, Caffraria (Dr. *A. Smith*). W. Africa (Whydah), (Coll. *Fraser*).

35. Aspidomorpha fusconotata. B.M.

Bohem. Mon. Cassid. ii. p. 279, 34.
Patr. India orientalis, Insul. Philipp. (Coll. *Cuming*).

36. Aspidomorpha sugillata.

Bohem. Mon. Cassid. ii. p. 280, 35.
Patr. Cap. Palmas.

37. ASPIDOMORPHA PUNCTUM.

Bohem. Mon. Cassid. ii. p. 281, 36.
Cassida Punctum, *Fabr.* Syst. El. i. p. 404, 96. *Oliv.* Ent. vi. p. 954, 50, 97, t. v. f. 76. *Schœnh.* Syn. Ins. ii. p. 226, 122. *Boisd.* Faun. Ent. de l'Océan. ii. p. 591, 9? *Hombr.* et *Jacquin.* Voy. d'Urville, t. xviii. f. 12.
Patr. Nova Guinea, Insulæ oceani pacifici.

38. ASPIDOMORPHA DIVISA.

Bohem. Mon. Cassid. ii. p. 282, 37.
Cassida divisa, *Boisd.* Faun. Entom. de l'Océan. ii. p. 540, 8.
Deloyala divisa, *Dej.* Cat. ed. 3, p. 395.
Patr. Nova Guinea.

39. ASPIDOMORPHA DOUEI.

Rotundata, modice convexa, dilute flavescens, nitida; antennis articulis duobus ultimis nigro-fuscis; prothorace lævi, elytris mediocriter, minus crebre punctato-striatis, antice leviter retusis et gibbosis, dorso plaga maxima, communi, nigra, medio posticeque profunde emarginata, gibbo, macula sat magna suturali pone medium punctoque infra callum humeralem, flavis; margine late explanato, hyalino, confertim reticulato. Long. 8¾, lat. 7¼ millim.
Bohem. Mon. Cassid. Append.
Patr. Nova Hollandia, Mus. Dom. *Doué.*

40. ASPIDOMORPHA BOISDUVALI.

Bohem. Mon. Cassid. ii. p. 283, 38.
Deloyala Punctum, *Dej.* Cat. ed. 3, p. 395.
Patr. Nova Hollandia.

III. *Elytra basi distincte gibbosa.*

41. ASPIDOMORPHA HEROINA.

Bohem. Mon. Cassid. ii. p. 284, 39.
Patr. — — —

42. ASPIDOMORPHA LIMBIPENNIS.

Bohem. Mon. Cassid. ii. p. 285, 40.
Cassida flava, *De Geer*, Mem. Ins. v. p. 184, 9, t. xv. f. 13.
Patr. Surinam.

43. ASPIDOMORPHA PONTIFEX.

Bohem. Mon. Cassid. ii. p. 286, 41.
Patr. Madagascar.

44. Aspidomorpha St. Crucis. B.M.

Bohem. Mon. Cassid. ii. p. 287, 42, t. vi. B.
Cassida St. Crucis, *Fabr.* Syst. El. i. p. 401, 80. Ent. Syst. iv. Append. p. 446, 41, 42. *Herbst.* Naturs. viii. p. 301, 68, t. 134, f. 12? *Illig.* Mag. v. p. 227, 80. *Schœnh.* Syn. Ins. ii. p. 223, 106.
Deloyala orientalis, *Dej.* Cat. ed. 3, p. 395.
Patr. India orientalis (Assam, Bengal).

45. Aspidomorpha bajula.

Bohem. Mon. Cassid. ii. p. 288, 43.
Deloyala Jamaicensis, *Dej.* Cat. ed. 3, p. 371.
Patr. America?

46. Aspidomorpha lobata. B.M.

Bohem. Mon. Cassid. ii. p. 289, 44.
Cassida Jamaicensis, *Herbst.* Naturs. viii. p. 303, 69, t. cxxxv. f. 1.
Patr. India orientalis (N. Bengal).

47. Aspidomorpha elevata. B.M.

Bohem. Mon. Cassid. ii. p. 291, 45.
Cassida elevata, *Fabr.* Syst. El. i. p. 399, 65. *Schœnh.* Syn. Ins. ii. p. 221, 87.
Deloyala elevata, *Dej.* Cat. ed. 3, p. 395.
Deloyala subcincta, *Dej.* Cat. ed. 3, p. 395.
Patr. Java, Sumatra (Coll. *Raffles*), Assam, Ind. orient.

48. Aspidomorpha quadrilobata. B.M.

Rotundata, convexa, flavo-testacea, nitida; antennis articulo ultimo apice nigro; prothorace lævi, antice late rotundato, lateribus posterius angustatis, sinuato-emarginatis; elytris subtiliter striato-punctatis, pone scutellum valide obtuse gibbosis, dorso plaga maxima, communi, rufo-castanea, utrinque ramulos duos in marginem emittente; humeris antrorsum valde prominulis, apice truncatis. Long. 12, lat. 11½ millim.

Bohem. Mon. Cassid. Append.
Patr. Manilla, Ins. Philipp. Mus. Holm. Mus. Brit. (Coll. *Cuming*).

49. Aspidomorpha bilobata. B.M.

Rotundata, modice convexa, dilute flavescens, nitida, subtus flavo-testacea, prothorace dorso saturatiore, subtilissime, sub-remote punctulato; elytris subtiliter striato-punctatis,

dorso plaga magna, communi, dilute rubro-castanea, utrinque ramulum basalem sat latum, postice medio ampliatum emittente; pone basin valide, breviter gibbosis; margine late explanato, confertim reticulato. Long. 10½, lat. 9¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Manilla, Mus. Reg. Holm. (Coll. *Cuming*).

50. ASPIDOMORPHA BIRADIATA. B.M.

Bohem. Mon. Cassid. ii. p. 292, 47.
Patr. Manilla, Ins. Philipp. (Coll. *Cuming*).

51. ASPIDOMORPHA SUBCRUCIATA. B.M.

Bohem. Mon. Cassid. ii. p. 293, 48.
Patr. Insul. Philipp. (Coll. *Cuming*).

52. ASPIDOMORPHA STICTICA.

Bohem. Mon. Cassid. ii. p. 294, 49.
Patr. — — —

53. ASPIDOMORPHA LIMBATA.

Bohem. Mon. Cassid. ii. p. 295, 50.
Cassida limbata, *Goldfuss.* Enum. Ins. Eleuth. p. 41, t. i. f. 4. *Schœnh.* Syn. Ins. ii. p. 221, 88.
Patr. Caput Bonæ Spei.

54. ASPIDOMORPHA DORSATA. B.M.

Bohem. Mon. Cassid. ii. p. 296, 51.
Cassida dorsata, *Fabr.* Syst. El. i. p. 401, 78. Ent. Syst. i. p. 301, 41. Mant. Ins. i. p. 64, 33. *Linn.* Syst. Nat. ed. Gmel. i. iv. p. 1641, 63. *Herbst.* Naturs. viii. p. 342, 109. *Schœnh.* Syn. Ins. ii. p. 223, 104.
Deloyala dorsata, *Dej.* Cat. ed. 3, p. 395.
Aspidomorpha elevata, var. *Hombr.* et *Jacquin.* Voy. d'Urville, t. xviii. f. 10.
Patr. India orientalis, Java, Sumatra.

55. ASPIDOMORPHA CALLIGERA. B.M.

Bohem. Mon. Cassid. ii. p. 297, 52.
Patr. Bengalia, Ceylon. Ins. Philipp. (Coll. *Cuming*).

56. ASPIDOMORPHA FUSCOPUNCTATA.

Bohem. Mon. Cassid. ii. p. 298, 53.
Patr. Java.

57. ASPIDOMORPHA QUADRIRADIATA. B.M.

Bohem. Mon. Cassid. ii. p. 300, 54.
Patr. Nova Hollandia ad Swan River, Port Essington.

58. ASPIDOMORPHA INUNCTA.

Bohem. Mon. Cassid. ii. p. 301, 55.
Patr. Malabar.

59. ASPIDOMORPHA DIAPHANA. B.M.

Bohem. Mon. Cassid. ii. p. 302, 56.
Cassida diaphana, *Sahlb.* Spec. Ins. 58, 39.
Deloyala diaphana, *Dej.* Cat. ed. 3, p. 395.
Patr. Sierra Leona, Guinea. Old Calabar (*A. Murray*).

60. ASPIDOMORPHA IMPICTA.

Bohem. Mon. Cassid. ii. p. 303, 57.
Patr. Gabon.

61. ASPIDOMORPHA NATALENSIS.

Bohem. Mon. Cassid. ii. p. 303, 58.
Patr. Caffraria ad Portum Natalensem.

62. ASPIDOMORPHA QUADRIRAMOSA.

Bohem. Mon. Cassid. ii. p. 305, 59.
Patr. Guinea.

63. ASPIDOMORPHA ICTERICA.

Bohem. Mon. Cassid. ii. p. 306, 60.
Patr. Caffraria interior, Guinea.

64. ASPIDOMORPHA NOVÆGUINEENSIS. B.M.

Bohem. Mon. Cassid. ii. p. 307, 61.
Cassida Novæguineensis, *Boisd.* Faun. Ent. de l'Océan. ii. p. 537, 3.
Aspidomorpha Astrolabiana, *Hombr.* et *Jacquin.* Voy. d'Urville, Ins. t. xviii. f. 13.
Patr. Nova Guinea. Solomon I. (Wanderer Bay).

65. ASPIDOMORPHA AUSTRALASIÆ.

Bohem. Mon. Cassid. ii. p. 308, 62.
Cassida Australasiæ, *Boisd.* Faun. de l'Océan. ii. p. 537, 2.
Patr. Nova Hollandia.

66. ASPIDOMORPHA SOCIA.

Rotundata, modice convexa, dilute flavescens, nitida; antennis articulis duobus ultimis nigro-fuscis; prothorace lævi, antice rotundato; elytris basi retusis, dein gibbo valido, brevi instructis, subtiliter punctato-striatis, singulo basi macula sub-triangulari, vitta disci exterioris inæquali, medio introrsum flexa nigro-piceis, hac ramulos duos latos in marginem et unum pone medium angustum ad suturam emittente; margine late explanato, hyalino, confertim reticulato. Long. 8, lat. 7½ millim.

Bohem. Mon. Cassid. Append.
Patr. Nova Guinea, Mus. Dom. *Lucas.*

67. ASPIDOMORPHA INQUINATA.

Bohem. Mon. Cassid. ii. p. 309, 63.
Patr. Java.

68. ASPIDOMORPHA RUBRODORSATA.

Bohem. Mon. Cassid. ii. p. 310, 64.
Patr. Java.

69. ASPIDOMORPHA MUTATA.

Bohem. Mon. Cassid. ii. p. 311, 65.
Deloyala tuberculata, *Dej.* Cat. ed. 3, p. 395.
Patr. Sierra Leona, Guinea.

70. ASPIDOMORPHA DILECTA.

Bohem. Mon. Cassid. ii. p. 312, 66.
Patr. Caffraria in tractibus fluvii Gariepis.

71. ASPIDOMORPHA MICANS. B.M.

Bohem. Mon. Cassid. ii. p. 313, 67.
Cassida micans, *Fabr.* Syst. El. i. p. 398, 64. *Schœnh.* Syn. Ins. ii. p. 221, 86.
Cassida furcata, *Thunb.* Nov. Ins. Spec. iv. p. 87, t. v. f. 96. *Herbst.* Naturs. viii. p. 265, 35, t. cxxx. f. 7?
Cassida dorsata, *Oliv.* Encycl. Ins. v. p. 386, 48. Entom. vi. p. 961, 64, 97, t. iii. f. 45?
Patr. India orientalis, China (Hong Kong), (*J. C. Bowring*), Java, Sumatra, Siam.

72. ASPIDOMORPHA ASSIMILIS.

Bohem. Mon. Cassid. ii. p. 314, 68.
Deloyala assimilis, *Dej.* Cat. ed. 3, p. 395.
Patr. Java.

73. Aspidomorpha amabilis. B.M.

Bohem. Mon. Cassid. ii. p. 315, 69.
Deloyala amabilis, *Dej.* Cat. ed. 3, p. 395.
Patr. Ind. orient. "Bot. Gardens, Calcutta" (Maj.-Gen. *Hardwicke*). Java, Assam.

74. Aspidomorpha mutilata.

Bohem. Mon. Cassid. ii. p. 316, 70.
Deloyala micans, *Dej.* Cat. ed. 3, p. 395.
Patr. Java.

75. Aspidomorpha egena.

Bohem. Mon. Cassid. ii. p. 317, 71.
Patr. Bengalia.

76. Aspidomorpha Indica.

Bohem. Mon. Cassid. ii. p. 318, 72.
Patr. Almorah, Indiæ orientalis borealis.

77. Aspidomorpha biremis.

Bohem. Mon. Cassid. ii. p. 319, 73.
Patr. Java.

Genus 31. ISCHYROSONYX.

Chevrolat. Dej. Cat. ed. 394, p. 3. *Bohem.* Mon. Cassid. ii. p. 321.

1. Ischyrosonyx oblonga.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 321, 1, t. vi. C.
Patr. Brasilia.

2. Ischyrosonyx peltoides.

Dej. Cat. ed. 3, p. 394. *Bohem.* Mon. Cassid. ii. p. 323, 2.
Patr. Brasilia.

3. Ischyrosonyx inanis.

Bohem. Mon. Cassid. ii. p. 224, 3.
Patr. Brasilia.

4. Ischyrosonyx discipennis. B.M.

Bohem. Mon. Cassid. ii. p. 325, 4.
Patr. Columbia.

5. Ischyrosonyx marginicollis.

Bohem. Mon. Cassid. ii. p. 326, 5.
Var. c. elytris rufo-testaceis immaculatis.
Patr. Columbia.

6. Ischyrosonyx nigrosignata. B.M.

Bohem. Mon. Cassid. ii. p. 327, 6.
Patr. Columbia.

Genus 32. CASSIDA.

Linn. Syst. Nat. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *De Geer,* Ins. v. p. 178. *Herbst.* Naturs. viii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209. *Dej.* Cat. ed. 3, p. 374. *Bohem.* Mon. Cassid. ii. p. 329.
Deloyala, *Dej.* Cat. ed. 3, p. 395.
Coptocycla, *Dej.* Cat. ed. 3, p. 396.

I. Unguiculi pectinati.

A. *Elytra cicatricosa, interstitiis dorsalibus elevatis.*

1. Cassida clathrata. B.M.

Cassida clathrata, *Fabr.* Syst. El. i. p. 396, 45. Suppl. p. 83, 22. *Herbst.* Naturs. viii. p. 303, 71, t. cxxxv. f. 3. *Schœnh.* Syn. Ins. ii. p. 219, 64. *Dej.* Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 330, 1.
Cassida sulcata, *Oliv.* Ent. vi. p. 950, 44, 97, t. v. f. 73.
Patr. India orientalis (Madras, *Walter Elliott,* Esq.).

B. *Elytra punctato-striata, interstitiis nodosis vel rugosis.*

2. Cassida nodulosa. B.M.?

Bohem. Mon. Cassid. ii. p. 331, 2.
Patr. Caffraria.

3. Cassida foveolata.

Ovata convexa, atra, sub-nitida; prothorace supra inæquali, dorso irregulariter, subtiliter aciculato, lateribus rugoso-punctato, antice utrinque macula parva sub-cuneata, pallida notato; elytris minus regulariter, mediocriter punctato-striatis, dorso profunde foveolatis, foveolis reticulo elevato, nitido inclusis; margine modice explanato, nonnihil deflexo, crebre ruguloso-punctato, ante medium foveola profunda impresso. Long. 8, lat. 5¾ millim.

Bohem. Mon. Cassid. Append.
Patr. India orientalis, Dom. *Downes,* Mus. Soc. Ind. Orient.

4. Cassida Abyssinica.

Oblongo-quadrata, convexa, atra, sub-opaca, setulis brevissimis, pallidis parce adspersa; antennis subtus basi ferrugineis; prothorace inæquali, undique profunde, crebre, irregulariter aciculato, apice utrinque flavo-testaceo, marginato; elytris vage, sat crebre rugoso-punctatis, dorso foveolatis, irregulariter reticulatis; margine parum explanato, profunde rugoso et foveolato; corpore subtus rude punctato. Long. 7½, lat. 4¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Abyssinia, Mus. Paris.

5. Cassida tetrica. B.M.

Ovata, convexa, flavo-testacea, nitida; antennis articulis duobus ultimis supra leviter infuscatis; prothorace lævi, basi utrinque arcu angusto maculaque parva media irregulari, obsoleta, brunneis; elytris sat profunde punctato-striatis, dorso plaga maxima brunnea, rugulis elevatis vario modo confluentibus, flavo-testaceis variegata, longe pone medium ramulum angustum in marginem emittente; margine modice explanato, limbo reflexo; humeris angulatis. Long. 7¾, lat. 6 millim.

Bohem. Mon. Cassid. Append.
Patr. Nova Hollandia, Mus. Britann.

6. Cassida Olivieri.

Bohem. Mon. Cassid. ii. p. 332, 3.
Patr. Oriens.

7. Cassida MacLeayi. B.M.

Rotundata valde convexa, dilute flavescens, nitida; antennis articulis tribus ultimis, prosterno, pectore abdomineque medio nigris; prothorace lævi, basi inæqualiter nigro medio vitta concolore, ante apicem arcu transverso retrorsum flexo conjuncta; scutello nigro; elytris nigris, mediocriter, irregulariter punctato-striatis, dorso rugulis flavis, irregularibus, vario modo confluentibus variegatis; margine modice explanato, lævi, limbo tenuiter reflexo. Long. 11, lat. 9 millim.

Bohem. Mon. Cassid. Append.
Patr. Nova Hollandia borealis, Mus. Britann.

8. Cassida maculatissima. B.M.

Rotundata, convexa, rufo-testacea, nitida; antennis apice corporeque subtus nigris, hoc late testaceo-marginato; pro-

thorace lævi, postice utrinque macula transversa, litura media triradiata, literam T fere referente maculaque parva anteapicali et scutello nigris; elytris irregulariter, mediocriter punctato-striatis, dorso nigris, rugulis parum elevatis, ramosis, sublævibus, rufo-testaceis variegatis; margine testaceo, maculis duabus magnis, una ante altera pone medium suturaque apice nigris; humeris antrorsum prominulis, antice extrorsum leviter rotundatis. Long 10½, lat. 8¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Cape Upstart, Novæ Hollandiæ (*J. Macgillivray*).

C. *Elytra punctato-striata, interstitiis vix vel parum convexis.*

a. *Prothorax nigro-vel fusco-maculatus.*

9. CASSIDA DEUSTA. B.M.

Fabr. Syst. El. i. p. 396, 44. Ent. Syst. i. p. 297, 22. Spec. Ins. i. p. 109, 15. Mant. Ins. i. p. 63, 19. Syst. Entom. p. 89, 8. *Linn.* Syst. Nat. Gmel. i. iv. p. 1636, 34. *Herbst.* Naturs. viii. p. 334, 97. *Oliv.* Encycl. meth. Ins. v. p. 382, 26. Entom. vi. p. 954, 51, 97, t. i. f. 17. *Schœnh.* Syn. Ins. ii. p. 218, 63. *Boisd.* Faun. Entom. de l'Océan. ii. p. 543, 3. *Bohem.* Mon. Cassid. ii. p. 333, 4.
Cassida angulifera, *Hombr.* et *Jaquin*, Voy. d'Urville, t. xviii. f. 17.
Cassida collina, *Mac Leay*, Boisd. Faun. Ent. de l'Océan. ii. p. 541, 10.
Patr. Nova Hollandia, Makassas, "Port Essington," Tasmania.

10. CASSIDA PRÆCOX.

Bohem. Mon. Cassid. ii. p. 334, 5.
Patr. Caffraria.

11. CASSIDA TIGRINA. B.M.

Oliv. Ent. vi. p. 957, 56, 97, t. v. f. 78. *Bohem.* Mon. Cassid. ii. p. 336, 6.
Cassida Solani, *Klug*, Dej. Cat. ed. 3, p. 398.
Patr. Caput Bonæ Spei, Caffraria (Dr. *A. Smith*). Port Natal (Coll. *Gueinzius*).

12. CASSIDA BIPUNCTICOLLIS.

Bohem. Mon. Cassid. ii. p. 337, 7.
Patr. Caffraria.

13. Cassida hybrida. B.M.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 338, 8.
Patr. Caput Bonæ Spei, Caffraria (*Earl of Derby*).

14. Cassida sedecimmaculata.

Sub-rotundata, convexa, testacea, nitida; antennis articulis ultimis leviter infuscatis; pectore abdomineque medio nigris; prothorace dorso punctato, ante basin maculis duabus sat magnis, sub-rotundis, nigris; elytris mediocriter, dorso minus crebre punctato-striatis, disco crebre, irregulariter punctatis, maculis quatuordecim inæqualibus, nigris notatis, binis suturalibus. Long. 7, lat. 5 millim.

Bohem. Mon. Cassid. Append.
Patr. India orientalis, Mus. Holm.

b. *Prothorax immaculatus.*

15. Cassida interrupta.

Ovata, convexa, testacea, nitida; antennis articulo ultimo nigro; prothorace lævi, immaculato, angulis posticis minus rotundatis; elytris subtiliter, seriatim punctatis, maculis duabus suturalibus, nonnullis disci, vitta disci exterioris medio interrupta nigris, hac basi et pone medium ramulum latum, concolorem in marginem emittente; humeris acutis. Long. 12, lat. 9½ millim.

Cassida interrupta, *Fabr.* Syst. El. i. p. 395, 40. Ent. Syst. i. p. 296, 20. Mant. Ins. i. p. 63, 17. Spec. Ins. i. p. 109, 14. Syst. Ent. p. 89, 7. *Linn.* Syst. Nat. Gmel. i. iv. p. 1636, 36. *Oliv.* Entom. v. p. 953, 49, 97, t. ii. f. 34. *Herbst.* Col. viii. p. 333, 95. *Schœnh.* Syn. Ins. ii. p. 218, 59. *Boisd.* Faun. Ent. de l'Océan. ii. p. 542, 2. *Bohem.* Mon. Cassid. Append.
Patr. Nova Hollandia, Mus. Dom. *Banks.*

16. Cassida nigrodorsata. B.M.

Ovata, modice convexa, rufo-testacea, nitida; antennis articulis quatuor ultimis nigris; prothorace lævi, immaculato; elytris mediocriter, minus regulariter, remote punctato-striatis, nigris, singulo in disco antico fovea sat profunda insculpto; margine maculis duabus oblongis flavo-testaceis; humeris antrorsum prominulis, acutis. Long. 9½, lat. 7 millim.

Bohem. Mon. Cassid. Append.
Patr. Nova Hollandia ad Port Essington, Mus. Britann.

17. Cassida illota.

Bohem. Mon. Cassid. ii. p. 339, 9.
Patr. Caput Bonæ Spei.

18. Cassida obovata.

Klug, Ermann Reise Ins. p. 48, 180. *Bohem.* Mon. Cassid. ii. p. 340, 10.
Patr. Mauritius.

19. Cassida nigrovittata.

Bohem. Mon. Cassid. ii. p. 341, 11.
Patr, India orientalis ad Calcutta.

D. *Elytra vage punctulata.*

20. Cassida hepatica. B.M.

Bohem. Mon. Cassid. ii. p. 342, 12.
Patr. Sierra Leona, Gambia (Coll. *Rendall*). Old Calabar (Coll. *Murray*).

21. Cassida spilota. B.M.

Bohem. Mon. Cassid. ii. p. 343, 13.
Patr. Sierra Leona (Rev. *D. F. Morgan*). Old Calabar (Coll. *Murray*).

22. Cassida lyncea.

Bohem. Mon. Cassid. ii. p. 344, 14.
Patr. Senegallia.

23. Cassida signatipennis. B.M.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 345, 15.
Patr. Sierra Leona, Senegallia. Guinea (*Whydah*).

24. Cassida Mouffleti.

Bohem. Mon. Cassid. ii. p. 346, 16.
Patr. Gabon.

25. Cassida hebes.

Bohem. Mon. Cassid. ii. p. 347, 17.
Patr. Caffraria.

26. Cassida punctata. B.M.

Fabr. Syst. El. i. p. 400, 74. Ent. Syst. i. p. 300, 38. Mant. Ins. i. p. 64, 30. *Herbst.* Naturs. viii. p. 341, 107. *Linn.*

Syst. Nat. ed. Gmel. i. iv. p. 1640, 59. *Schœnh.* Syn. Ins. ii. p. 222, 99. *Bohem.* Mon. Cassid. ii. p. 348, 18, t. vi. D.
Cassida maculipennis, *Dej.* Cat. ed. 3, p. 398.
Patr. Caput Bonæ Spei, Caffraria (Dr. *A. Smith*), (*Earl of Derby*). Port Natal (Coll. *Gueinzius*).

27. CASSIDA PARUMMACULATA. B.M.

Bohem. Mon. Cassid. ii. p. 349, 19.
Patr. Caput Bonæ Spei, Caffraria. Port Natal (Coll. *Gueinzius*).

28. CASSIDA ADSPERSA.

Fabr. Syst. El. i. p. 400, 75. *Illig.* Mag. i. p. 407, 75. *Schœnh.* Syn. Ins. ii. p. 222, 100. *Bohem.* Mon. Cassid. ii. p. 350, 20.
Patr. Caput Palmas.

29. CASSIDA NIGROSPARSA.

Bohem. Mon. Cassid. ii. p. 351, 21.
Patr. Guinea.

30. CASSIDA NILOTICA.

Bohem. Mon. Cassid. ii. p. 352, 22.
Var. b. prothorace maculis tribus parvis, rotundis, brunneis, media reliquis majore, singulo elytro maculis parvis numerosis, inæqualibus circiter 21-24, nigro-fuscis adsperso.
Var. c. elytris rufo-testaceis, prothorace ut in genuina aut ut in var. *b.* abdomine fere toto nigro.
Patr. Nubia.

31. CASSIDA LUTEICOLLIS. B.M.

Bohem. Mon. Cassid. ii. p. 353, 23.
Patr. Africa australis, Port Natal.

II. Unguiculi mutici.

A. *Elytra pone basin non gibbosa.*

a. *Elytra punctato-striata, non costata.*

* *Margo elytrorum explanatus, limbo reflexo.*

32. CASSIDA VIGINTIMACULATA. B.M.

Thunb. Nov. Ins. Spec. v. p. 86, f. 95. *Herbst.* Naturs. viii. p. 264, 34, t. cxxxii. f. 6. *Schœnh.* Syn. Ins. ii. p. 219, 67. *Dej.* Cat. ed. 3. p. 398. *Bohem.* Mon. Cassid. ii. p. 354, 24.
Cassida nigronotata, *Dej.*
Patr. Caput Bonæ Spei, Caffraria (Dr. *A. Smith*). Port Natal (Coll. *Gueinzius*).

33. CASSIDA SCRIPTA.

Fabr. Syst. El. i. p. 395, 42. Suppl. p. 82, 21, 22. *Herbst.* Naturs. viii. p. 351, 123. *Schœnh.* Syn. Ins. ii. p. 218, 61. *Bohem.* Mon. Cassid. ii. p. 356, 25.
Patr. Caput Bonæ Spei, Caffraria.

34. CASSIDA AUSTRIACA. B.M.

Fabr. Syst. El. i. p. 390, 10. Ent. Syst. i. p. 393, 6. *Herbst.* Naturs. viii. p. 254, 24, t. cxxxi. f. 9. *Panz.* Faun. Germ. 98, t. 2. Ent. Germ. p. 104, 5. *Oliv.* Ent. vi. p. 980, 99, 97, t. vi. f. 108. *Schœnh.* Syn. Ins. ii. p. 212, 11. *Dej.* Cat. ed. 3, p. 398. *Suffr.* Stett. Ent. Zeit. v. p. 95, 3. *Redtenb.* Faun. Austr. p. 520. *Bach.* Stett. Ent. Zeit. xii. p. 158. *Bohem.* Mon. Cassid. ii. p. 357, 26.
Cassida canaliculata, *Laich.* Tyrol. Ins. i. ii. p. 109, 1.
Cassida sanguinolenta, *Schrank.* En. Ins. p. 48, 88. Beytr. p. 61, 241.
Cassida speciosa, *Brahm.* in Naturs. p. 29, 166. Ins. Kalend. i. p. 74, 241. *Schœnh.* Syn. Ins. ii. p. 214, 20.
Var. Cassida Illyrica, *Dej.* Cat. ed. 3, p. 398.
Var. Cassida immutabilis, *Ziegl.* Dej. Cat. ed. 3, p. 374.
Patr. Europa media et australis.

35. CASSIDA MURRÆA. B.M.

Linn. Syst. Nat. i. ii. p. 575, 2, ed. Gmel. i. iv. p. 1635, 2. *Fabr.* Syst. El. i. p. 390, 14. Ent. Syst. i. p. 294, 9. Mant. Ins. i. p. 62, 8. Spec. Ins. i. p. 108, 6. Syst. Ent. p. 90, 10. *Illig.* Káf. Preuss. i. p. 476, 1, var. β. *Payk.* Faun. Suec. ii. p. 48, 5 a. *Oliv.* Ent. vi. p. 977, 95, 97, t. i. f. 7. Encycl. meth. v. p. 379, 5. *Herbst.* Naturs. viii. p. 240, 17, t. cxxx. f. 12, 13. *Panz.* Ent. Germ. i. p. 154, 8. *Rossi,* Faun. Etr. i. p. 72, 181. Ed. Hellw. i. p. 77, 181. *Preysl.* Bohm. Ins. p. 90, 84. *Scriba,* Journ. p. 179, 132. *Thunb.* Ins. Suec. viii. p. 104. *Blumenb.* Handb. ed. 7, p. 325, 2. *Marsh.* Ent. Britann. i. p. 147, 10. *Brahm,* Ins. Kal. i. p. 181, 600. *Vill.* Ent. i. p. 91, 2. *Geoffr.* Ins. i. p. 314, 5. *Schœnh.* Syn. Ins. ii. p. 213, 17. *Gyllenh.* Ins. Suec. iii. p. 334, 1. *Dej.* Cat. ed. 3, p. 374. *Suffr.* Stett. Ent. Zeit. v. p. 101, 5. *Redtenb.* Faun. Austr. p. 520. *Pflümer,* Stett. Ent. Zeit. viii. p. 71, ix. p. 91. *Bohem.* Mon. Cassid. ii. p. 358, 27.
Var. Cassida maculata. *Linn.* Syst. Nat. i. ii. p. 575, 6. Ed. Gmel. i. iv. p. 1638, 6. *Fabr.* Mant. Ins. i. p. 62, 3. Spec. Ins. i. p. 107, 2. Syst. Ent. p. 82, 2. *De Geer,* Mem. iv. p. 175, 3. *Oliv.* Enc. meth. v. p. 378, 4. *Brahm.* Ins. Kal. i. p. 216, 722. *Kirby,* Acta Soc. Linn. iii. p. 10, 3. *Goeze.* Eur. Faun. viii. p. 312, 3. *Marsh.* Ent. Brit. i. p. 147, 9.

Müll. Linn. Syst. Nat. v. i. p. 92, 6. *Geoffr.* Ins. i. p. 314, 5, t. v. f. 6.
Patr. Europa tota, Dauria et Sibiria.

36. CASSIDA TRILINEATA. B.M.

Ovalis, modice convexa, rufo-testacea, nitida; antennis extrorsum, prosterno pectore abdomineque nigris, hoc anguste ferrugineo-marginato; prothorace sub-lævi, dorso vittis tribus, lateralibus antice conjunctis, macula utrinque laterali scutelloque nigris; elytris inæqualiter, dorso postico subtilius punctato-striatis, sutura, vitta disci inæquali, ramulos tres extrorsum emittente nigris; margine paullo explanato, nonnihil deflexo, reticulato. Long. 5½. lat. 4 millim.

Cassida trilineata, *Hope*, Gray Zool. Miscell. 1831, p. 30. *Bohem.* Mon. Cassid. Append.

Var. a. prothorace ut in genuina, elytrorum vitta ramulisque nigris latioribus, his vario modo confluentibus.

Var. b. prothorace ut in genuina, elytris dorso nigris, maculis 12 inæqualibus flavis notato.

Var. c. prothorace toto, apice excepto, nigro, elytris dorso nigris pone medium ramulum concolorem in marginem emittente.

Patr. India orientalis (Gen. *Hardwicke*), Dom. *Saunders*, Mus. Holm. *Var. b.* Mus. Dom. *Waterhouse.*

37. CASSIDA VITTATA. B.M.

Fabr. Syst. El. i. p. 391, 15. Suppl. p. 81, 9, 10. *Illig.* Mag. iii. p. 157, 15. *Herbst.* Naturs. viii. p. 353, 127. *Oliv.* Entom. vi. p. 982, 103, 97, t. vi. f. 111. *Steph.* Illustr. iv. p. 365, 1, t. xxiii. f. 1. *Schœnh.* Syn. Ins. ii. p. 214, 18. *Dej.* Cat. ed. 3, p. 399. *Suffr.* Stett. Ent. Zeit. v. p. 98. 4, vii. p. 361. *Redtenb.* Faun. Austr. p. 520. *Bohem.* Mon. Cassid. ii. p. 360, 28.
Cassida ocellata, *Herbst.* Naturs. viii. p. 244, 18, t. cxxxi. f. 4 b.
Cassida fastuosa, *Schall.* Act. Hal. i. p. 259? *Scriba*, Beytr. i. p. 86, 17, t. viii. f. 3. *Linn.* Syst. Nat. Gmel. i. iv. p. 1639, 52? *Panz.* Ent. Germ. i. p. 155, 13?
Cassida variabilis, *Dahl.* in litt.
Var. Cassida vittata, *Ahrens*, Neue Hall. Schrift. ii. p. 2, 31, t. ii. f. 6.
Var. e. supra flavo-testacea, prothoracis signaturis obsoletis, maculis elytrorum nigris ut in var. *b.*
Patr. Europa media: Anglia, Germania, Austria, Hungaria, Podolia et Russia meridionalis. *Var. e.* Mongolia.

38. Cassida nigripes.

Oliv. Enc. meth. v. p. 384, 34. Entom. vi. p. 959, 61, 97, t. v. f. 81. *Bohem.* Mon. Cassid. ii. p. 362, 29.
Coptocycla rufa, *Dej.* Cat. ed. 3, p. 397.
Patr. Georgia Americæ borealis, Brasilia?

39. Cassida novemmaculata.

Mannerh. Bull. de Moscou, xxi. p. 308. *Bohem.* Mon. Cassid. ii. p. 363, 30.
Patr. California.

** *Margo elytrorum explanatus, limbo non reflexo.*

40. Cassida areata. B.M.

Klug, Ermanns Reise Ins. p. 48, 179. *Bohem.* Mon. Cassid. ii. p. 364, 31.
Cassida Dregei, *Dej.* Cat. ed. 3, p. 398.
Patr. Caput Bonæ Spei, Caffraria (Dr. *A. Smith*).

41. Cassida Moori.

Ovata, modice convexa, nigra, sub-nitida; prothorace crebre punctulato, rufo-testaceo, basi plaga magna oblongo-quadrata, transversa, nigra, antice leviter rotundato-emarginata; elytris dilute sanguineis, crebre, irregulariter punctato-striatis, macula pone scutellum communi, singulo maculis duabus basalibus, plaga magna media fasciaque postica inæquali, communi, nigris; margine modice explanato, parum deflexo, rugoso-punctato. Long. 9, lat. 6½ millim.
Bohem. Mon. Cassid. Append.
Patr. India orientalis, Mus. Soc. Ind. Orient.

42. Cassida coronula.

Bohem. Mon. Cassid. ii. p. 365, 32.
Patr. Madagascar, Senegallia.

43. Cassida obscuripennis.

Ovata, convexa, rufo-testacea, sub-nitida; prothorace sublævi, ante basin maculis quatuor rotundis, nigris; elytris subtiliter punctato-striatis, nigris, limbo anguste rufo-testaceo. Long. 7, lat. 5¼ millim.
Bohem. Mon. Cassid. Append.
Patr. Have-Blanche, Mus. Paris.

44. Cassida lurida.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 366, 33.
Patr. Senegallia.

45. Cassida mutabilis. B.M.

Klug, Ermanns Reise Ins. p. 47, 178. *Dej.* Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 367, 34.
Patr. Senegallia.

46. Cassida æthiopica.

Bohem. Mon. Cassid. ii. p. 368, 35.
Patr. Nubia.

47. Cassida sanguinosa. B.M.

Creutz, Mus. Berol. *Suffr.* Stett. Ent. Zeit. v. p. 105, 6. *Redtenb.* Faun. Austr. p. 520. *Bohem.* Mon. Cassid. ii. p. 369, 36.
Cassida prasina, *Herbst.* Naturs. viii. p. 221, 4, t. cxxix. f. 6, d. *Schœnh.* Syn. Ins. ii. p. 211, 4.
Cassida rubiginosa, *Gyllenh.* Ins. Suec. iii. p. 457, 3, var. *c.* iv. p. 643, 3, Obs.
Cassida thoracica, *Steph.* Illustr. iv. p. 366, 4.
Cassida sanguinolenta, var. *Dej.* Cat. ed. 3, p. 398.
Patr. Europa media et borealis.

48. Cassida denticollis.

Suffr. Stett. Ent. Zeit. v. p. 186, 11. *Redtenb.* Faun. Austr. p. 521. *Bohem.* Act. Holm. 1851, p. 97. *Cornelius*, Stett. Entom. Zeit. viii. p. 359. *Bohem.* Mon. Cassid. ii. p. 371, 37.
Patr. Europa borealis et media.

49. Cassida seladonia.

Gyllenh. Ins. Suec. iv. p. 644, 3, 4. *Bohem.* Mon. Cassid. ii. p. 372, 38.
Patr. Scania ad Esperód Sueciæ.

50. Cassida rubiginosa. B.M.

Illig. Káf. Preuss. i. p. 479, 4. Mag. i. p. 93. *Herbst.* Naturs. viii. p. 223, 5, t. cxxix. f. 7. *Müll.* Zool. Dan. Pr. p. 65, 612. *Gyllenh.* Ins. Suec. iii. p. 437, 3, excl. var. *c. d.* *Steph.* Illustr. iv. p. 366, 3. *Suffr.* Stett. Ent. Zeit. v. p. 135, 7. *Redtenb.* Faun. Austr. p. 520. *Elditt.* Stett. Ent. Zeit. xii. p. 352. t. ii. A. B. *Bohem.* Mon. Cassid. ii. p. 373, 39.
Cassida viridis, *Fabr.* Syst. El. i. p. 387, 1. Ent. Syst. i. p. 272, 1. Mant. Ins. i. p. 62, 1. Spec. Ins. i. p. 107, 1. Syst. Ent. p. 88, 1. *Payk*, Faun. Suec. ii. p. 45, 2 a. Act. Holm. 1801, p. 119 a. *Latr.* Gen. Crust. et Ins. iii. p. 53. *Oliv.* Ent. vi. p. 975, 91, 97, t. ii. f. 29. Encycl. meth. v. p. 378, 1, var. fem. nigris. *Panz.* Faun. Germ. 96, t. iv. Ent. Germ. i.

p. 153, 1. *Cederh.* Faun. Ingr. No. 172. *Rossi,* Faun. Etr. i. p. 72, 180. Ed. Hellw. i. p. 77, 180. *Fisch.* Naturg. v. Livl. p. 132, 258. *Schwarz.* Nomencl. i. p. 48, 20. *Blumenb.* Handb. ed. 7, p. 324, 1. *Scriba,* Journ. p. 176, 128, var. *Laich.* Tyrol. Ins. i. ii. p. 111, 3, var. β. *Barb.* Gen. Ins. p. 34, t. iii. *Kirby,* Act. Soc. Linn. iii. p. 9, 2? *Fourcr.* Ent. Par. i. p. 140, 1. *Dej.* Cat. ed. 3, p. 398. *Schœnh.* Syn. Ins. ii. p. 210, 1. *Zetterst.* Ins. Lapp. p. 219, 1.

Cassida Cardui, var. *De Geer,* Mem. v. p. 174, 2.

Cassida similis, *Marsh.* Ent. Brit. i. p. 144, 2. *Roes.* Ins. ii. t. 6. *Reaum.* Ins. iii. t. xviii. *Merian,* Ins. Europ. t. cxv. *Schœff.* Icon. t. xxxvi. f. 2. *Harrer,* Beschreib, No. 153.

Cassida nigra, *Herbst.* Naturs. viii. p. 258, 27, t. cxxxi. f. 12. *Schœnh.* Syn. Ins. ii. p. 212, 14. *Suffr.* Stett. Ent. Zeit. v. p. 142, 7 (*b*). (Individuum materia aliena nigra tinctum.)

Patr. Europa tota.

51. CASSIDA LATA.

Suffr. Stett. Ent. Zeit. v. p. 138, a. *Bohem.* Mon. Cassid. ii. p. 375, 40.

Patr. Turcia, Sicilia.

52. CASSIDA THORACICA. B.M.

Fabr. Syst. El. i. p. 388, 2. *Illig.* Káf. Preuss. i. p. 478, 3. *Herbst.* Naturs. viii. p. 228, 10, t. cxxix. f. 11, i. *Fourcr.* Ent. Paris. i. p. 141, 4. *Panz.* Faun. Germ. 38, t. 24. *Oliv.* Ent. vi. p. 976, 93, 97, t. iii. f. 48. *Schœnh.* Syn. Ins. ii. p. 210, 2. *Gyllenh.* Ins. Suec. iii. p. 439, 4. *Dej.* Cat. ed. 3, p. 398. *Suffr.* Stett. Ent. Zeit. v. p. 142, 8. *Redtenb.* Faun. Austr. p. 520. *Bohem.* Mon. Cassid. ii. p. 376, 41.

Cassida viridis, *Payk.* Act. Holm. 1801, p. 120, v. β. Faun. Suec. ii. p. 46, 2, v. β. *Latr.* Gen. Crust. et Ins. iii. p. 52, var. 1. *Geoffr.* Ins. i. p. 314, 4.

Patr. Europa tota et Asia, Podolia et Sibiria.

53. CASSIDA VIBEX. B.M.

Fabr. Syst. El. i. p. 389, 6. Ent. Syst, i. p. 293, 4. Mant. Ins. i. p. 108, 4. Syst. Ent. p. 89, 4. *Linn.* Syst. Nat. i. ii. p. 575, 5. Ed. Gmel. i. iv. p. 1637, 5. *Illig.* Káf. Preuss. i. p. 477, 2. *Payk,* Act. Holm. 1801, p. 118. *Herbst.* Naturs. viii. p. 218, 3, t. cxxix. f. 5, c. *Panz.* Faun. Germ. 96, f. 6. Ent. Germ. i. p. 154, 4. *Oliv.* Ent. vi. p. 977, 94, 97, t. ii. f. 30. Encycl. meth. v. p. 379, 7. *Scriba,* Beitr. ii. p. 80, 15, t. viii. f. 1. Journ. p. 178, 129. *Thunb.* Ins. Suec. viii. p. 104. *Rossi,* Mant. ii. Append. p. 89, 34. *Schrank,* Enum. Ins. p. 48, 89. *Brahm.* Ins. Kal. i. p. 66, 212, et 232, 789. *Vill.* Ent. i. p. 92, 5. *Müll.* Linn. Nat. Syst. v. i. p. 138, 5.

Schœnh. Syn. Ins. ii. p. 211, 5. *Gyllenh.* Ins. Suec. iii. p. 440, 5. *Dej.* Cat. ed. 3. p. 398. *Suffr.* Stett. Ent. Zeit. v. p. 146, 10, *Redtenb.* Faun. Austr. p. 520. *Bohem.* Mon. Cassid. ii. p. 377, 42.
Cassida viridis, *Payk,* Faun. Suec. ii. p. 46, 2, var. β. *Latr.* Gen. Crust. et Ins. iii. p. 3, 52, var. 2. *Laich.* Tyrol. Ins. i. ii. p. 111, var. γ.
Cassida dorsalis, *Herbst.* Naturg. viii. p. 227, 9? t. cxxix. f. 10.
Patr. Europa tota et Sibiria.

54. Cassida deflorata. B.M.

Illig. Suffr. Stett. Ent. Zeit. v. p. 139, b. *Bohem.* Mon. Cassid. ii. p. 379, 43.
Cassida Scolymi, *Küster* in litt.
Cassida gallica, *Meg.* in. litt.
Cassida herbea, *Latr.*
Patr. Europa meridionalis: Gallia, Sardinia, Sicilia, Africa borealis.

55. Cassida depressa.

v. Heyden Suffr. Stett. Ent. Zeit. v. p. 141, d. *Bohem.* Mon. Cassid. ii. p. 380, 44.
Patr. Europa meridionalis: Hispania, Sardinia, Sicilia.

56. Cassida prasina. B.M.

Fabr. Syst. El. i. p. 388, 4. *Bohem.* Mon. Cassid. ii. p. 381, 45.
Patr. Africa borealis ad Algier. Europa meridionalis: Gallia et Sicilia.

57. Cassida hexastigma.

Kunze Suffr. Stett. Ent. Zeit. v. p. 140, c. *Bohem.* Mon. Cassid. ii. p. 382, 46.
Patr. Gallia meridionalis, Italia.

58. Cassida languida. B.M.

Cornelius, Stett. Ent. Zeit. xii. p. 88. *Bohem.* Mon. Cassid. ii. p. 383, 47.
Patr. Germania (Elberfeldt), Dr. *Cornelius.*

59. Cassida Chloris. B.M.

Suffr. Stett. Ent. Zeit. v. p. 188, 12. *Cornelius,* l. c. viii. p. 361. *Bohem.* Mon. Cassid. ii. p. 384, 48.
Patr. Germania, Austria, Hungaria.

60. Cassida stigmatica.

Illig. Suffr. Stett. Ent. Zeit. v. p. 206, 12 (b.). *Bohem.* Mon. Cassid. ii. p. 385, 49.
Cassida viridana, *Herbst.* Naturs. viii. p. 224, 6, t. cxxix. f. 8? *Schœnh.* Syn. Ins. ii. p. 212, 8?
Patr. Germania, Austria, Hungaria.

61. Cassida rufovirens.

Suffr. Stett. Ent. Zeit. v. p. 144, 9. *Bohem.* Mon. Cassid. ii. p. 386, 50.
Patr. Germania, Austria.

62. Cassida rosea.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 387, 51.
Patr. Styria.

63. Cassida sanguinolenta. B.M.

Fabr. Syst. El. i. p. 389, 8. *Illig.* Káf. Preuss. i. p. 482, 7. *Herbst.* Naturs. viii. p. 229, 11, t. cxxix. f. 12. *Oliv.* Ent. vi. p. 978, 96, 97, t. vi. f. 107. *Latr.* Gen. Crust. et Ins. iii. p. 52. *Payk.* Faun. Suec. ii. p. 48, 3. *Müll.* Zool. Dan. Pr. p. 65, 611. *Gyllenh.* Ins. Suec. iii. p. 441, 6. *Schœnh.* Syn. Ins. ii. p. 212, 7. *Steph.* Illustr. iv. p. 369, 10. *Zetterst.* Ins. Lapp. p. 219, 2. *Suffr.* Stett. Ent. Zeit. v. p. 208, 13. *Redtenb.* Faun. Austr. p. 521. *Bohem.* Mon. Cassid. ii. p. 388, 52.
Cassida cruentata, *Marsh.* Ent. Britann. i. p. 145. *Donov.* Br. Ins. t. lxiii. f. 2, 3.
Patr. Europa media et borealis.

64. Cassida conformis.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 389, 53.
Patr. Brasilia.

65. Cassida spadicea. B.M.

Klug, Preis-Verzeichn. 1829, p. 8, 100. *Dej.* Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 390, 54.
Patr. Brasilia, Buenos Ayres, Monte Video.

66. Cassida lutescens.

Bohem. Mon. Cassid. ii. p. 391, 55.
Patr. Brasilia.

67. Cassida graminea. B.M.

Klug, Preis-Verzeichn. 1829, p. 8, 101. *Dej.* Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 392, 56.
Patr. Brasilia.

68. Cassida sulphurea.

Bohem. Mon. Cassid. ii. p. 393, 57.
Patr. Caput Bonæ Spei.

69. Cassida viridipennis.

Bohem. Mon. Cassid. ii. p. 394, 58.
Patr. Caput Bonæ Spei, Caffraria.

70. Cassida dorsovittata.

Bohem. Mon. Cassid. ii. p. 395, 59.
Patr. Caput Bonæ Spei, Caffraria.

71. Cassida troglodytes.

Bohem. Mon. Cassid. ii. p. 396, 60.
Patr. Sierra Leona.

72. Cassida lacrymosa.

Bohem. Mon. Cassid. ii. p. 397, 61.
Patr. Caffraria.

73. Cassida syrtica.

Rotundata, modice convexa, dilute flavo-testacea, nitida; antennis articulis tribus ultimis corporeque subtus nigris, prosterno abdomineque late flavo-testaceo marginatis; prothorace dorso punctulato, ante basin maculis tribus parvis, transversim positis, media oblonga, lateralibus rotundis, nigris; elytris crebre, mediocriter punctato-striatis, basi impressis, dorso nigris, rugulis flavis variegatis; margine sat late explanato, hyalino, confertim reticulato. Long. 5-5¼, lat. 4½ millim.

Bohem. Mon. Cassid. Append.
Patr. India orientalis, Dom. *Waterhouse*, Mus. Holm.

74. Cassida irregularis.

Bohem. Mon. Cassid. ii. p. 398, 62.
Patr. Caffraria.

75. Cassida rugulosa.

Rotundata, modice convexa, dilute flavescens, nitida; pro-

sterno, pectore abdomineque medio nigris, hoc segmentis apice testaceis; prothorace obsolete, sub-remote punctulato, angulis posticis rotundatis; elytris profunde, sat crebre punctato-striatis, ad scutellum utrinque leviter impressis, dorso nigris, rugulis sat crebris, flavis, parum elevatis, vario modo confluentibus variegatis; margine sat late explanato, oblique parum deflexo, hyalino, confertim reticulato, ante medium impresso. Long. 5, lat. 4 millim.

Bohem. Mon. Cassid. Append.
Var. a. elytrorum dorso parcius flavo-variegato.
Patr. India orientalis, Dom. *Wight.* Mus. Soc. Ind. Orient.

76. CASSIDA FUSCOSIGNATA.

Bohem. Mon. Cassid. ii. p. 399, 62.
Patr. Senegallia.

77. CASSIDA ANCORIFERA. B.M.

Rotundata, modice convexa, dilute flavescens, nitida; antennis articulo ultimo nigro-fusco; prothorace lævi, basi signo ancoræformi, nigro notato; elytris crebre, mediocriter punctato-striatis, basi utrinque ad scutellum impressis, dorso plaga magna communi, nigra, extus medio utrinque leviter emarginata, pone scutellum macula sat magna, rugula instructa et pone medium arcu retrorsum flexo, communibus, flavis; margine late explanato, parum oblique deflexo, confertim reticulato. Long. 4½, lat. 3¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Sierra Leona, Mus. Britann. (Rev. *D. F. Morgan*).

78. CASSIDA ICTERICA.

Bohem. Mon. Cassid. ii. p. 400, 64.
Patr. Almorah Indiæ septentrionalis.

79. CASSIDA TENUICULA.

Rotundata, minus convexa, dilute flavescens, nitida; antennis articulo ultimo fusco; prothorace lævi, plaga mediocri, basali, rotundata, maculis duabus parvis in medio scutelloque nigris; elytris subtiliter striato-punctatis, basi ad scutellum utrinque impressis, dein leviter gibbosis, annulo antico sat lato fasciaque pone medium semilunata, communibus, nigro-fuscis; margine late explanato, hyalino, reticulato; humeris parum prominulis, sub-acutis. Long. 3¾, lat. 3½ millim.

Bohem. Mon. Cassid. Append.
Patr. Guinea, Mus. Dom. *Westermann.*

80. Cassida conspurcata.

Bohem. Mon. Cassid. Append.
Patr. Tranquebaria.

81. Cassida timefacta.

Rotundata, convexa, dilute flavo-testacea, nitida; antennis articulis duobus ultimis nigro-fuscis; prothorace sub-lævi ante basin macula parva ovata, fusca; elytris crebre mediocriter punctato-striatis, basi ad scutellum utrinque impressis, dein ruga transversa instructis, dorso fusco-variegatis; margine sat late explanato, distincte punctato. Long. 5, lat. 4 millim.

Bohem. Mon. Cassid. Append.
Patr. Ceylon, Mus. Dom. *Deyrolle.*

82. Cassida piperita. B.M.

Hope, Proceed. Ent. Soc. 1842, p. 62.
Cassida labilis, *Bohem.* Mon. Cassid. ii. p. 402, 66.
Var. a. supra ut genuina colorata, corpore subtus pulchre cyaneo, nitido.
Patr. China.

83. Cassida innotata.

Bohem. Mon. Cassid. ii. p. 403, 67.
Patr. Caffraria.

84. Cassida granula.

Rotundata, modice convexa, testacea, parum nitida; prothorace obsolete punctulato, angulis posticis late rotundatis; elytris sat profunde, crebre punctato-striatis, ad scutellum profunde impressis, dein rugula transversa, communi, arcuata instructis, dorso plaga sat magna rufo-testacea, extus dilute fusco-marginata ornata; margine modice explanato, parum deflexo, rugoso-punctato; humeris nonnihil prominulis, apice acutis. Long. 5½, lat. 4½ millim.

Bohem. Mon. Cassid. Append.
Patr. Guinea, Mus. Reg. Holm.

85. Cassida bella.

Steven. Falderm. Faun. Ent. Transcauc. P. ii. in Nouv. Mem. de la Soc. de Natur. de Mosc. v. p. 325, 523, t. xii. f. 3.
Bohem. Mon. Cassid. ii. p. 404, 68.
Cassida splendidula, *Godet*, Dej. Cat. ed. 3, p. 399.
Patr. Russia meridionalis: Tauria et Caucasia.

86. Cassida obtusata.

Bohem. Mon. Cassid. ii. p. 405, 69.
Patr. India orientalis et China.

87. Cassida pallida. B.M.

Rotundata, modice convexa, testacea, nitida; antennis articulo ultimo, prosterno, pectore abdomineque nigris, hoc extus flavo-testaceo marginato; prothorace obsolete punctulato, angulis posticis obtusis; elytris mediocriter, ultra medium crebre, dein remotius punctato-striatis, ad scutellum et in dorso antico leviter impressis, antice ruga obsoleta, transversa, communi instructis; margine sat late explanato, hyalino, obsolete punctulato; humeris prominulis, acutis. Long. 5¼, lat. 4¼.

Cassida pallida, *Hope*, Gray Zool. Miscell. 1831, p. 30. *Bohem.* Mon. Cassid. Append.
Patr. Tranquebar, Mus. Britann. (Gen. *Hardwicke*).

88. Cassida pauxilla.

Bohem. Mon. Cassid. ii. p. 406, 70.
Patr. India orientalis.

89. Cassida exilis.

Bohem. Mon. Cassid. ii. p. 407, 71.
Patr. Pondichery, India orientalis.

90. Cassida Delesserti.

Bohem. Mon. Cassid. ii. p. 408, 72.
Patr. India orientalis.

91. Cassida dorsonotata.

Bohem. Mon. Cassid. ii. p. 409, 73.
Patr. India orientalis.

92. Cassida nigriventris.

Bohem. Mon. Cassid. ii. p. 410, 74.
Patr. Thibet.

93. Cassida azurea. B.M.

Fabr. Syst. El. i. p. 389, 7. *Oliv.* Ent. vi. p. 981, 100, 97, t. vi. f. 109. *Schœnh.* Syn. Ins. ii. p. 212, 9. Dej. Cat. ed. 3, p. 374. *Suffr.* Stett. Ent. Zeit. v. p. 213, 15, vii. p. 362. *Redtenb.* Faun. Austr. p. 520. *Bohem.* Mon. Cassid. ii. p. 411, 75.

Cassida ornata, *Creutz*, Ent. Verz. i. p. 118, 10, t. ii. f. 22, a.
Patr. Austria, Bavaria, Dalmatia, Hungaria, Gallia.

94. Cassida lucida. B.M.

Suffr. Stett. Ent. Zeit. v. p. 215, 16, vii. p. 362. *Bohem.* Mon. Cassid. ii. p. 412, 76.
Cassida azurea, *Dej.* Cat. ed. 3, p. 398 var.
Patr. Ems, Gallia meridionalis, Coblentz (Dr. *Cornelius*).

95. Cassida rorida.

Bohem. Mon. Cassid. ii. p. 413, 77.
Patr. Caffraria.

*** *Margo elytrorum oblique deflexus.*

96. Cassida sulphurago.

Bohem. Mon. Cassid. ii. p. 414, 78.
Patr. Caffraria.

97. Cassida obscura. B.M.

Fabr. Syst. El. i. p. 392, 26. Ent. Syst. i. p. 295, 16. *Herbst.* Naturs. viii. 332, 93. *Schœnh.* Syn. Ins. ii. 217, 35. *Bohem.* Mon. Cassid. ii. p. 415, 79.
Cassida corrosa, *Dej.* Cat. ed. 3, p. 398.
Var. Cassida unicolor, *Fabr.* Syst. El. i. p. 391, 18. *Schœnh.* Syn. Ins. ii. p. 216, 26.
Cassida œanthe, Dom. *Buquet* in litt.
Patr. India orientalis, Senegallia.

98. Cassida cruenta.

Fabr. Syst. El. i. p. 389, 9. Ent. Syst. i. p. 293, 5. *Herbst.* Naturs. viii. p. 330, 90. *Schœnh.* Syn. Ins. ii. p. 212, 10. *Bohem.* Mon. Cassid. ii. p. 416, 80.
Patr. India orientalis.

99. Cassida pudibunda.

Ovata, convexa, sub-nitida, sordide flavescens; antennis articulo ultimo apice, pectore abdomineque basi maculis nigro-fuscis; prothorace confertim, rude, disco antico obsolete rugoso-punctato, basi linea brevi scutelloque dilute rubris; elytris mediocriter crebre punctato-striatis, basi non gibbosis, interstitiis convexis, sutura dilute rubra; margine modice explanato, crebre punctato; humeris antrorsum prominulis, sub-angulatis. Long. 6¼, lat. 4½ millim.
Bohem. Mon. Cassid. Append.
Patr. India orientalis, Mus. Reg. Holm.

100. Cassida convexa.

Bohem. Mon. Cassid. ii. p. 418, 81.
Patr. Caffraria.

101. Cassida Tunisiensis.

Bohem. Mon. Cassid. ii. p. 419, 82.
Patr. Tunis.

102. Cassida umbrata.

Bohem. Mon. Cassid. ii. p. 420, 83.
Patr. Caffraria.

103. Cassida convexula.

Bohem. Mon. Cassid. ii. p. 421, 84.
Patr. Caffraria.

104. Cassida oblonga. B.M.

Illig. Káf. Preuss. p. 485, 2. *Suffr.* Stett. Ent. Zeit. v. p. 252, 20. *Redtenb.* Faun. Austr. p. 521. *Bohem.* Mon. Cassid. ii. p. 422, 85.
Cassida nobilis, *Herbst.* Naturs. viii. p. 238, 16, t. cxxx. f. 15.
Cassida Salicorniæ, *Steph.* Illustr. iv. p. 370, 14.
Cassida pulchella, *Dej.* Cat. ed. 3, p. 399.
Patr. Europa media et meridionalis, Africa borealis.

105. Cassida nobilis. B.M.

Linn. Faun. Suec. p. 153, 469. Syst. Nat. i. ii. p. 575, 4. Ed. Gmel. i. iv. p. 1636, 4. *Fabr.* Syst. El. i. p. 396, 47. Ent. Syst. i. p. 297, 24. Mant. Ins. i. p. 63, 21. Spec. Ins. i. p. 110, 17. Syst. Ent. p. 90, 11. *Illig.* Káf. Preuss. i. p. 485, 11. *Payk.* Faun. Suec. ii. p. 49, 7. *Oliv.* Ent. vi. p. 903, 104, 97, t. ii. f. 24. Encycl. method. v. p. 383, 28. *Panz.* Ent. Germ. i. p. 155, 11. *Marsh,* Ent. Brit. i. p. 146, 7. *Cederh.* Faun. Ingr. No. 175. *Scriba,* Journ. p. 180, 134. *Beitr.* ii. p. 87, 18, t. viii. f. 4. *Rossi,* Faun. Etr. i. p. 72, 182. Ed. Hellw. i. p. 78, 182. *Müll.* Zool. Dan. Prodr. p. 65, 609. *Goeze,* Eur. Faun. viii. p. 315, 6. *Donov.* Br. Ins. t. cxxxviii. f. 1, 3. *Schrank,* Faun. Boic. i. p. 521, 610. *Bechst.* Naturg. i. ii. p. 836, 5. *Pontopp.* Naturg. p. 201, 3. *Müll.* Linn. Syst. Nat. v. i. p. 137, 4. *Villers,* Ent. i. p. 92. 4. *Schæff.* Icon. t. 96, f. 6, t. 142, f. 7, a. b. *Harrer,* Beschr, p. 151, ed. 2, No. 103. *Geoffr.* Ins. i. p. 313, 3. *Ray,* Icon. p. 107, 7. *Steph.* Illustr. iv. p. 370, 15. *Schœnh.* Syn. Ins. ii. p. 219, 68. *Gyllenh.* Ins. Suec. iii. p. 445, 9. *Dej.* Cat. ed. 3, p. 399. *Zetterst.* Ins. Lapp. p. 220, 5. *Suffr.* Stett. Ent. Zeit. v.

p. 245, 19, vii. p. 363. *Redtenb.* Faun. Austr. p. 521. *Bohem.* Mon. Cassid. ii. p. 423, 86.
Cassida urticæ, *Brahm.* Ins. Kal. i. p. 109, 307.
Var. Cassida lævis, *Herbst.* Naturs. viii. p. 250, 21, t. cxxxi. f. 6, d.
Cassida pulchella, *Panz.* Faun. Germ. 39, t. xv.
Cassida splendidula, *Marsh.* Ent. Brit. i. p. 147, 8. *Steph.* Illustr. iv. p. 371, 16.
Var. Cassida Spergulæ, *Marsh.* Ent. Brit. i. p. 144, 3.
Cassida margaritacea, *Herbst.* Naturs. viii. p. 225, 7 (ex parte).
Cassida viridula, *Payk.* Faun. Suec. ii. p. 49, 6. *Schœnh.* Syn. Ins. ii. p. 220, 71. *Gyllenh.* Ins. Suec. iii. p. 447, 10. *Dej.* Cat. ed. 3, p. 399. *Suffr.* Stett. Ent. Zeit. v. p. 249, 19 (b).
Patr. Europa tota et Sibiria.

106. Cassida nana.

Bohem. Mon. Cassid. ii. p. 425, 87.
Patr. Caffraria.

107. Cassida pumilio.

Bohem. Mon. Cassid. ii. p. 426, 88.
Patr. Caffraria.

108. Cassida setifera.

Bohem. Mon. Cassid. ii. p. 426, 89.
Patr. Senegallia.

109. Cassida glabella.

Bohem. Mon. Cassid. ii. p. 428, 90.
Patr. Neelgheries, Indiæ orientalis.

110. Cassida parvula.

Bohem. Mon. Cassid. ii. p. 428, 91.
Patr. Mongolia.

111. Cassida navicula.

Bohem. Mon. Cassid. ii. p. 429, 92.
Patr. Ad mare Caspicum.

112. Cassida gibbula.

Bohem. Mon. Cassid. ii. p. 430, 93.
Patr. Turcomania.

113. Cassida minuta.

Bohem. Mon. Cassid. ii. p. 431, 94.
Patr. Caput Bonæ Spei.

114. Cassida margaritacea. B.M.

Fabr. Syst. El. i. p. 397, 51. Ent. Syst. i. p. 297, 25. Mant. Ins. i. p. 63, 22. *Herbst.* Naturs. viii. p. 225, 7 (ex parte), t. cxxx. f. 16. *Oliv.* Ent. vi. p. 284, 105, 97, t. ii. f. 19. Encycl. meth. v. p. 383, 29. *Schall.* Act. Hal. i. p. 259. *Rossi*, Mant. i. p. 27, 66. Ed. Hellw. i. p. 364, 66. *Scriba*, Journ. p. 180, 136. *Linn.* Syst. Nat. Gmel. i. iv. p. 1639, 51. *Panz.* Ent. Germ. i. p. 155, 12. *Gœze*, Eur. Faun. viii. p. 314, 5. *Schrank*, Faun. Boic. i. p. 521, 609. *Schœnh.* Syn. Ins. ii. p. 220, 72. *Redtenb.* Faun. Austr. p. 521. *Dej.* Cat. ed. 3, p. 399. *Suffr.* Stett. Ent. Zeit. v. p. 241, 17, vii. p. 363. *Bohem.* Mon. Cassid. ii. p. 432, 95.
Cassida mutabilis, *Villers*, Ent. t. i. f. 11.
Patr. Europa media et meridionalis, Asia et Africa borealis.

115. Cassida insularis.

Bohem. Mon. Cassid. ii. p. 433, 96.
Patr. Madagascar.

116. Cassida sphærula.

Bohem. Mon. Cassid. ii. p. 434, 97.
Patr. Caput Bonæ Spei.

117. Cassida Hottentotta.

Bohem. Mon. Cassid. ii. p. 435, 98.
Patr. Africa meridionalis.

118. Cassida subreticulata.

Suffr. Stett. Ent. Zeit. v. p. 244, 18. *Redtenb.* Faun. Austr. p. 521.
Bohem. Mon. Cassid. ii. p. 436, 99.
Patr. Austria, Carinthia.

119. Cassida mollis.

Bohem. Mon. Cassid. ii. p. 437, 100.
Patr. Lusitania.

120. Cassida pusilla.

Waltl, Reise durch Tyrol. Ober Ital. und Piemont nach.

Südl. Spanien 1839, p. 82. *Bohem.* Mon. Cassid. ii. p. 438, 101.
Cassida puncticollis, *v. Heyden,* Suffr. Stett. Ent. Zeit. v. p. 257.
Cassida herbacea, *Dej.* Cat. ed. 3, p. 399.
Patr. Lusitania, Sicilia, Gallia meridionalis.

121. Cassida pupillata.

Bohem. Mon. Cassid. ii. p. 439, 102.
Patr. Sicilia.

122. Cassida pulvinata.

Bohem. Mon. Cassid. ii. p. 440, 103.
Patr. Pondichery, India orientalis.

123. Cassida denticulata.

Rotundata, valde convexa, flavo-testacea, sub-nitida; antennis articulis quinque ultimis corporeque nigris, abdomine anguste testaceo-marginato; prothorace sublævi, antice rotundato, postice utrinque modice obliquo, angulis rotundatis; elytris profunde, sat crebre punctato-striatis, interstitiis sub-convexis, singulo apice denticulo instructo; margine modice explanato, oblique deflexo, obsolete punctulato, longe infra medium macula nigro-fusca notato. Long. 4, lat. 3¼ millim.

Bohem. Mon. Cassid. Append.
Patr. Nova Hollandia, Mus. Dom. *Baly.*

124. Cassida semiglobosa.

Bohem. Mon. Cassid. ii. p. 441, 104.
Patr. Caffraria.

125. Cassida bistrilineata.

Bohem. Mon. Cassid. ii. p. 442, 105.
Patr. Montevideo, Buenos Ayres.

126. Cassida tristriata.

Fabr. Syst. El. i. p. 392, 27. Ent. Syst. i. p. 295, 17. *Herbst.* Naturs. viii. p. 332, 94. *Oliv.* Ent. vi. p. 972, 85, 97, t. vi. f. 101. *Schœnh.* Syn. Ins. ii. p. 217, 36. *Bohem.* Mon. Cassid. ii. p. 443, 106.
Coptocycla tristriata, *Dej.* Cat. ed. 3, p. 398.
Patr. Cayenna.

127. Cassida striolata.

Bohem. Mon. Cassid. ii. p. 444, 107.
Coptocycla striolata, *Dej.* Cat. ed. 3, p. 397.
Patr. America borealis.

128. Cassida vittula.

Bohem. Mon. Cassid. ii. p. 445, 108.
Patr. America borealis.

129. Cassida graphica.

Germ. Ins. Spec. i. p. 543, 736. *Bohem.* Mon. Cassid. ii. p. 446, 109.
Coptocycla graphica, *Dej.* Cat. ed. 3, p. 398.
Patr. Buenos Ayres et St. Paul, Brasiliæ.

130. Cassida æmula.

Bohem. Mon. Cassid. ii. p. 447, 110.
Patr. Brasilia.

b. *Elytra punctato-striata, costata.*

131. Cassida Mongolica.

Bohem. Mon. Cassid. ii. p. 449, 111.
Patr. Mongolia.

132. Cassida pallidicollis.

Rotundata, convexa, nigra, sub-opaca; capite, antennis basi, scutello, abdomine extus prothoraceque dilute flavescentibus, hoc confertim, mediocriter punctato; elytris crebre, mediocriter punctato-striatis, singulo dorso bicostato, basi prope scutellum macula parva, sub-triangulari lineolaque brevi infra medium in costa interiore dilute flavescentibus; margine modice explanato, crebre, vage punctato. Long. 6, lat. 5 millim.

Bohem. Mon. Cassid. Append.
Patr. China, Dom. *Buquet.* Mus. Holm.

133. Cassida lineola. B.M.

Creutz. Ent. Verz. i. p. 119, 11, t. ii. f. 23, a. *Schœnh.* Syn. Ins. ii. p. 213, 16. *Dej.* Cat. ed. 3, p. 398. *Suffr.* Stett. Ent. Zeit. v. p. 210, 14. *Redtenb.* Faun. Austr. p. 521. *Bohem.* Mon. Cassid. ii. p. 450, 112.
Cassida Russica, *Herbst.* Naturs. viii. p. 232, 12, t. cxxxi. f. 1.
Var Cassida signata, *Herbst.* Naturs. viii. p. 234, 13, t. cxxxi. f. 2. *Schœnh.* Syn. Ins. ii. p. 213, 15.

Cassida aurea, *Bæber.*
Patr. Austria, Hungaria, et Russia meridionalis.

134. CASSIDA NEBULOSA. B.M.

Linn. Faun. Suec. p. 152, 468. Syst. Nat. i. ii. p. 575, 3. Ed. Gmel. i. iv. p. 1636, 3. *Herbst.* Naturs. viii. p. 236, 15, t. cxxx. f. 9, 11. *Oliv.* Ent. vi. p. 979, 97, t. ii. f. 31. Encycl. meth. v. p. 379, 8. *Scriba,* Journ. p. 178, 130. *Brahm.* Ins. Kal. i. p. 56, 181. *Scriba,* Beitr. ii. p. 83, 16, t. viii. f. 2. *Petagn.* Ins. Calabr. p. 8, 38. *Marsh,* Ent. Brit. i. p. 145, 6. *Villers,* Ent. i. p. 91, 3. *Müll.* Zool. Dan. Pr. p. 65, 610. *Schrank.* Enum. Ins. p. 49, 91. Faun. Boic. i. p. 520, 603. *Gœze,* Eur. Faun. viii. p. 310, 2. *Geoffr.* Ins. i. p. 313. 2. *Schæff.* Icon. t. xxvii. f. 4. *Müll.* Linn. Syst. Nat. v. i. p. 137, 3. *Gœd.* Met. i. p. 96, t. 44. *List.* Gœd. p. 287, 117. *Illig.* Káf. Preuss. i. p. 483, 9. *Payk.* Faun. Suec. ii. p. 47, 4. *Pontopp.* Nachtr. p. 201, 2. *Schœnh.* Syn. Ins. ii. p. 214, 21. *Gyllenh.* Ins. Suec. iii. p. 442, 7. *Zetterst.* Ins. Lapp. p. 220, 3. *Suffr.* Stett. Ent. Zeit. v. p. 277, 24. *Redtenb.* Faun. Austr. p. 522. *Bohem.* Mon. Cassid. ii. p. 451, 113. *Wollaston,* Insecta Maderensia, p. 439.

Var. Cassida affinis, *Fabr.* Syst. El. i. p. 388, 5. Ent. Syst. i. p. 293, 3. Mant. Ins. i. p. 62, 4. Spec. Ins. i. p. 108, 3. Syst. Ent. p. 88, 3. *Oliv.* Ent. vi. p. 980, 98, 97, t. iii. f. 7. Encycl. meth. v. p. 379, 6. *Panz.* Ent. Germ. i. p. 153, 3. *Cederh.* Faun. Ingr. No. 173. *Thunb.* Ins. Suec. viii. p. 104. *Herbst.* Arch. p. 50, 2. *Linn.* Syst. Nat. Gmel. i. iv. p. 1637, 45. *Laich,* Tyrol. Ins. i. ii. p. 110, 2. *Villers,* Ent. p. 93, 8. *Dej.* Cat. ed. 3, p. 398.

Cassida tigrina, *De Geer,* Mem. v. p. 168, 1, 5, f. 15, 16. *Oliv.* Encycl. meth. v. p. 378, 3. *Rossi,* Mant. ii. Append. 89, 35. *Scriba,* Journ. p. 178, 131. *Panz.* Ent. Germ. i p. 155, 14. *Frisch.* Ins. iv. p. 30, 15, t. xv.

Patr. Europa tota, Sibiria. Madera, prope Funchal (Coll. *Wollaston*).

135. CASSIDA FERRUGINEA. B.M.

Fabr. Syst. El. i. p. 391, 16. Ent. Syst. i. p. 294, 10. Mant. Ins. i. p. 62, 9. Spec. Ins. i. p. 102, 7. *Illig.* Káf. Preuss. i. p. 482, 8. *Herbst.* Naturs. viii. p. 245, 19, t. cxxx. f. 14. Arch. p. 50, 6, t. xxii. f. 29. *Oliv.* Ent. vi. p. 981, 101, 97, t. i. f. 2. Encycl. meth. v. p. 380, 10. *Panz.* Ent. Germ. i. p. 154, 9. *Scriba,* Journ. p. 179, 133. *Thunb.* Ins. Suec. viii. p. 104. *Rossi,* Faun. Etr. i. p. 73, 184. Ed. Hellw. i. p. 78, 184. *Petagna,* Ins. Calabr. p. 9, 39. *Goeze,* Eur. Faun. viii. p. 313, 4. *Linn.* Syst. Nat. Gmel. i. iv. p. 1637, 44. *Brahm.* Ins. Kal. i. p. 239, 280. *Schœnh.* Syn. Ins. ii. p.

214, 19. *Steph.* Illustr. iv. p. 368, 8. *Dej.* Cat. ed. 3, p. 398. *Suffr.* Stett. Ent. Zeit. v. p. 274, 23. *Redtenb.* Faun. Austr. p. 521. *Cornelius,* Stett. Ent. Zeit. viii. p. 365, x. p. 22. *Bohem.* Mon. Cassid. ii. p. 453, 114.

Cassida fusca, *Laich,* Tyrol. Ins. i. ii. p. 112, 4. *Oliv.* Encycl. meth. v. p. 380, 13. *Herbst.* Arch. p. 51, 9. *Linn.* Syst. Nat. Gmel. i. iv. p. 1639, 53.

Cassida subferruginea, *Schrank,* Beitr. p. 62, 3. Enum. Ins. Austr. p. 49, 90. *Schæff.* Icon. t. 40, f. 5, 6. *Harrer,* Beschr. v. 149.

Patr. Europa media et meridionalis, Sibiria.

136. CASSIDA MERIDIONALIS. B.M.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 454.

Patr. Gallia meridionalis.

137. CASSIDA COSTATA.

Ovata, convexa, griseo-flavescens, nitida; antennis articulis quatuor ultimis dilute fuscescentibus; pectore postice plaga transversa nigro-picea; prothorace obsolete punctulato, postice utrinque oblique truncato, angulis obtusis; elytris sat profunde, minus regulariter punctato-striatis, basi utrinque impressis, obsolete gibbosis; singulo carinis tribus, interiore reliquis magis elevato; margine modice explanato, oblique deflexo, rugoso-punctato. Long. 5¾, lat. 3¾ millim.

Bohem. Mon. Cassid. Append.

Patr. Madras, Mus. Soc. Ind. orient.

138. CASSIDA OBSOLETA. B.M.

Illig. Käf. Preuss. i. p. 424, 10. *Herbst.* Naturs. viii. p. 248, 20, t. cxxxi. f. 5. *Schœnh.* Syn. Ins. ii. p. 215, 22. *Gyllenh.* Ins. Suec. iii. p. 444, 8. *Steph.* Illustr. iv. p. 368, 9. *Dej.* Cat. ed. 3, p. 399. *Zetterst.* Ins. Lapp. p. 220, 4. *Suffr.* Stett. Ent. Zeit. v. p. 272, 22. *Redtenb.* Faun. Austr. p. 521. *Bohem.* Mon. Cassid. ii. p. 455, 116.

Cassida nebulosa, *Fabr.* Syst. El. i. p. 390, 11. Ent. Syst. i. p. 293, 7. Mant. Ins. i. p. 62, 6. Spec. Ins. i. p. 108, 5. Syst. Ent. p. 89, 5. *Rossi,* Faun. Etr. p. 73, 183. Ed. Hellw. i. p. 78, 183? *Cederh.* Faun. Ingr. No. 174. *Panz.* Ent. Germ. i. p. 153, 6.

Cassida pallida, *Payk,* Faun. Suec. ii. p. 50, 8.

Cassida flaveola, *Thunb.* Ins. Suec. viii. p. 103.

Cassida ferruginea, *Marsh,* Ent. Brit. i. p. 148, 11.

Cassida exsculpta, *Charp.* Horæ Entom. p. 231.

Patr. Europa tota.

139. Cassida pallidula.

Bohem. Mon. Cassid. ii. p. 457, 117.
Patr. Mexico ad Oaxaca.

140. Cassida Daurica.

Bohem. Mon. Cassid. ii. p. 457, 118.
Patr. Dauria.

141. Cassida Berolinensis.

Dej. Cat. ed. 3, p. 398. *Suffr.* Stett. Ent. Zeit. v. p. 270, 21.
Bohem. Mon. Cassid. ii. p. 458, 119.
Patr. Borussia, prope Berolinum et Sedinum.

c. *Elytra seriatim punctata.*

142. Cassida heteropunctata.

Bohem. Mon. Cassid. ii. p. 459, 120.
Patr. Columbia.

143. Cassida quadriplagiata.

Bohem. Mon. Cassid. ii. p. 460, 121.
Coptocycla quadriplagiata, *Dej.* Cat. ed. 3, p. 397.
Patr. Senegallia.

144. Cassida opposita. B.M.

Bohem. Mon. Cassid. ii. p. 461, 122.
Patr. Senegallia. Sierra Leona (Rev. *D. F. Morgan*).

145. Cassida octoguttata.

Bohem. Mon. Cassid. ii. p. 462, 123.
Coptocycla octoguttata, *Dej.* Cat. ed. 3, p. 397.
Patr. Senegallia.

146. Cassida quadriguttata.

Bohem. Mon. Cassid. ii. p. 464, 124.
Coptocycla quadriguttata, *Dej.* Cat. ed. 3, p. 397.
Patr. Senegallia.

147. Cassida Suffriani.

Bohem. Mon. Cassid. ii. p. 465, 125.
Patr. Caffraria.

148. Cassida unimaculata.

Bohem. Mon. Cassid. ii. p. 466, 126.
Patr. Caffraria.

149. Cassida quinquemaculata. B.M.

Bohem. Mon. Cassid. ii. p. 467, 127.
Patr. Insul. Philipp. (Coll. *Cuming*).

d. *Elytra rugosa vel tuberculata.*

150. Cassida coagulata.

Bohem. Mon. Cassid. ii. p. 468, 122.
Patr. Caffraria.

151. Cassida testudinea. B.M.

Bohem. Mon. Cassid. ii. p. 469, 129.
Patr. Caput Bonæ Spei.

152. Cassida atrata. B.M.

Fabr. Syst. El. i. p. 390. 13. Ent. Syst. i. p. 294, 8. Mant. Ins. i. p. 62, 7. *Herbst.* Naturs. viii. p. 257, 26, t. cxxxi. f. 11, f. *Oliv.* Ent. vi. p. 982, 102, 97, t. vi. f. 110. Encycl. meth. v. p. 380, 9. *Panz.* Faun. Germ. p. 88, t. 1. Ent. Germ. i. p. 154, 7, *Linn.* Syst. Nat. Gmel. i. iv. p. 1636, 32. *Schœnh.* Syn. Ins. ii. p. 212, 13. *Dej.* Cat. ed. 3, p. 398. *Suffr.* Stett. Ent. Zeit. v. p. 282, 25. *Redtenb.* Faun. Austr. p. 519. *Bohem.* Mon. Cassid. ii. p. 470, 130.
Patr. Austria, Styria, Bavaria, et Hungaria.

153. Cassida fossulata. B.M.

Ovata, convexa, sordide dilute flavescens, nitida, abdomine basi infuscato; prothorace inæquali, lævi, angulis posticis valde rotundatis; elytris subtiliter, minus regulariter punctato-striatis, dilute fuscis, basi utrinque impressis, dein leviter gibbosis, dorso irregulariter, valde elevato-reticulatis; margine modice explanato, ante medium macula mediocri intus rotundata et limbo postico albidis. Long. 7½. lat 5¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Calabar, Mus. Dom. *Baly.*

154. Cassida callosa.

Bohem. Mon. Cassid. ii. p. 471, 131.
Patr. America borealis.

155. Cassida rugosa. B.M.

Bohem. Mon. Cassid. ii. p. 472, 132.
Deloyala rugosa, *Dej.* Cat. ed. 3, p. 395.
Patr. Mexico ad Yucatan. Honduras (Coll. *Dyson*).

156. Cassida rimosa.

Aspidomorpha rimosa. *Bohem.* Mon. Cassid. ii. p. 290, 45.
Patr. Madagascar.

157. Cassida tædiosa.

Sub-triangularis, convexa, nitida, dilute flavescens; antennis articulis quatuor ultimis nigris; prothorace obsolete, crebre punctato, dorso longitudinaliter aciculato, postice plaga magna triangulari scutelloque nigris; elytris nigris, profunde, crebre punctato-striatis, basi ad scutellum utrinque profunde impressis, dorso parce rugosis; humeris antice, limbo medio anguste, macula irregulari intramarginali apiceque flavis; margine sat late explanato oblique deflexo, crebre rugoso-punctato. Long. 5½, lat. 4½ millim.

Bohem. Mon. Cassid. Append.
Patr. Madagascar, Mus. Dom. *Deyrolle.*

158. Cassida frontalis.

Sub-rotundata, convexa, nigra, sub-nitida; fronte, antennis, prothorace antice tibiis tarsisque testaceis; prothorace paullo convexo, dorso crebre aciculato, antice lateribusque rugoso-punctato; elytris crebre, sat profunde punctato-striatis, rugulis valde elevatis, remotis, anastomosantibus instructis, basi ad scutellum impressis. Long. 4½, lat. 3¾ millim.

Bohem. Mon. Cassid. Append.
Patr. Madagascar, Dom. Com. *Mannerheim*, Mus. Holm.

159. Cassida fuscosparsa.

Bohem. Mon. Cassid. ii. p. 473, 133.
Patr. Assam, India orientalis.

e. *Elytra vage punctata.*

160. Cassida equestris. B.M.

Fabr. Syst. El. i. p. 388, 3. Ent. Syst. i. p. 292, 2. Mant. Ins. i. p. 62, 2. *Payk.* Faun. Suec. ii. p. 45, 1. *Oliv.* Ent. vi. p. 976, 92, 97, t. i. f. 3. Encycl. method. v. p. 378, 2. *Latr.* Gen. Crust. et Ins. iii. p. 51, 1. Hist. Nat. des Crust. et Ins. xii. p. 28. *Herbst.* Naturs. viii. p. 216, 2, t. cxxx. f. 10. *Panz.* Faun. Germ. 96, f. 5. Ent. Germ. i. p. 153, 2. *Thunb.* Ins. Suec. viii. p. 103. *Linn.* Syst. Nat. Gmel. i. iv. p. 1638, 46. *Gyllenh.* Ins. Suec. iii. p. 436, 2. *Schœnh.* Syn. Ins. ii. p. 210, 3. *Dej.* Cat. ed. 3, p. 398. *Suffr.* Stett. Ent.

Zeit. v. p. 89, 1. *Redtenb.* Faun. Austr. p. 519. *Cornelius,* Stett. Ent. Zeit. viii. p. 363. *Bohem.* Mon. Cassid. ii. p. 474, 134.

Cassida viridis, *Linn.* Faun. Suec. p. 152, 467. Syst. Nat. i. ii. p. 571. 1. It. Oel. p. 53. *Illig.* Káf. Preuss. i. p. 480, 5. *Oliv.* Ent. vi. p. 975, 91, 97, t. ii. f. 29. Encycl. meth. v. p. 378, 1. *Scriba,* Journ. p. 176, 128, var. *Scop,* Ent. Carn. p. 37, 117. *Herbst.* Naturs. viii. p. 209, 1, t. cxxx. f. 8. *Marsh,* Ent. Br. i. p. 143, 1. *Gœze.* Eur. Faun. viii. p. 304, 1. *Linn.* Syst. Nat. Gmel. i. iv. p. 1635, 1. *Müll.* Zool. Dan. Pr. p. 65, 608.

Cassida Cardui, *De Geer,* Mem. v. p. 174, 2. *Schæff.* Elem. t. xxxv. Icon. t. xxvii. f. 5. *Harrer,* Beschr. No. 147. Ed. 2, No. 97. *Frisch.* Ins. xiii. 35, t. xxix. *Gœd.* Met. i. p. 94, t. xliii. *Blank,* Belg. p. 89, t. xi. f. F. *Merian,* Europ. t. cxv. *Volkam,* Hesperid. i. p. 166, f. 4. *Ray,* Ins. p. 107, 5. Act. Upsal, 1736, p. 17, 1.

Patr. Europa tota.

161. Cassida nigriceps.

Fairm. Ann. Soc. Ent. Fr. ii. Ser. x. p. 92, 25. *Bohem.* Mon. Cassid. ii. p. 476, 135.

Patr. Lusitania, Hispania.

162. Cassida consimilis.

Dej. Cat. ed. 3, p. 399. *Bohem.* Mon. Cassid. ii. p. 477, 136.

Patr. Caput Bonæ Spei.

163. Cassida scutellaris.

Klug, Ermanns Reise Ins. p. 48, 182. Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 478, 137.

Patr. Senegallia.

164. Cassida limbiventris.

Bohem. Mon. Cassid. ii. p. 479, 138.

Patr. Senegallia.

165. Cassida decolorata.

Rotundata, modice convexa, supra dilute virescens sub-opaca, subtus flavo-testacea nitida; antennis articulo penultimo apice, ultimo toto nigris; prothorace crebre, subtiliter rugoso-punctato, angulis posticis rotundatis; elytris dorso crebre, vage, mediocriter punctulatis, ad scutellum subretusis, dein leviter gibbosis; margine late explanato, mo-

dice oblique deflexo, profundius, crebrius punctato. Long. 6, lat. 5 millim.
Bohem. Mon. Cassid. Append.
Patr. Anjonama, Dom. *Sommer.* Mus. Holm.

166. CASSIDA MELANOPHTHALMA.

Bohem. Mon. Cassid. ii. p. 480, 139.
Patr. Caput Bonæ Spei.

167. CASSIDA LITIGIOSA.

Dej. Cat. ed. 3, p. 399. *Bohem.* Mon. Cassid. ii. p. 481, 140.
Patr. Caput Bonæ Spei, Senegallia.

168. CASSIDA PUDENS.

Bohem. Mon. Cassid. ii. p. 482, 141.
Patr. Caput Bonæ Spei.

169. CASSIDA CORDULA.

Bohem. Mon. Cassid. ii. p. 482, 142.
Patr. Caput Bonæ Spei.

170. CASSIDA TENERA.

Bohem. Mon. Cassid. ii. p. 483, 143.
Patr. Caput Bonæ Spei

171. CASSIDA HEMISPHÆRICA. B.M.

Herbst. Naturs. viii. p. 226, 8, t. cxxix. f. 9 g. *Schœnh.* Syn. Ins. ii. p. 220, 73. *Steph.* Illustr. iv. p. 319, 12. *Gyllenh.* Ins. Suec. iv. p. 645, 11. *Dej.* Cat. ed. 3, p. 399. *Suffr.* Stett. Ent. Zeit. v. p. 92, 2, vii. p. 361. *Redtenb.* Faun. Austr. p. 519. *Bohem.* Mon. Cassid. ii. p. 484, 144. *Wollaston,* Insecta Maderensia, p. 439.
Cassida margaritacea, *Thunb.* Ins. Suec. viii. p. 103. *Brahm.* Ins. Kal. i. p. 73, 240. *Steph.* Illustr. iv. p. 317, 17. *Payk.* Faun. Suec. ii. p. 49, 6. not. sub C. viridula.
Cassida concinna, *Steph.* Illustr. iv. p. 317, 18.
Patr. Europa tota et Africa borealis. Madera (Coll. *Wollaston*).

B. *Elytra pone basin distincte gibbosa.*

172. CASSIDA COMBUSTA. B.M.

Bohem. Mon. Cassid. ii. p. 485, 145.
Patr. Cayenna. Amazon (*R. Tapayos*), (Coll. *H. W. Bates*).

173. Cassida tosta.

Klug, Ermanns Reise Ins. p. 47, 176. *Bohem.* Mon. Cassid. p. 486, 146.
Patr. Guinea, Isle de Prince.

174. Cassida farinosa.

Sub-triangularis, convexa, supra obscure, subtus dilutius ferruginea, nitida; antennis articulis quatuor ultimis nigro-fuscis; prothorace dorso remote, parum profunde, margine nonnihil crebrius et obsoletius punctato; elytris sat profunde, minus regulariter striato-punctatis, basi retusis, profunde impressis, dein gibbosis; margine modice explanato, oblique deflexo, dilutiori, crebre rugoso-punctato; humeris antrorsum prominulis, apice rotundatis. Long. 6¾, lat. 5½ millim.

Bohem. Mon. Cassid. Append.
Patr. Ceylon, Mus. Dom. *Dohrn.* Insectum vivum pulvere niveo, mucoris instar adspersum, secundum observat. collectoris.

175. Cassida gibbipennis. B.M.

Dej. Cat. ed. 3, p. 398. *Bohem.* Mon. Cassid. ii. p. 488, 147, t. vi. E.
Patr. Caput Bonæ Spei (*Earl of Derby*).

176. Cassida roseomarginata.

Bohem. Mon. Cassid. ii. p. 489, 148.
Patr. Senegallia.

177. Cassida pæminosa..

Oblongo-subtriangularis, flavo-testacea, sub-nitida; prothorace mediocriter, crebre punctato; elytris sat profunde, vage punctatis, dorso nigris, crebre, irregulariter flavo-testaceo-rugosis, basi ad scutellum retusis, dein leviter gibbosis; margine modice explanato, oblique deflexo, vage, minus profunde punctato. Long. 6¼, lat. 4 millim.

Bohem. Mon. Cassid. Append.
Patr. Guinea, Dom. *Buquet.* Mus. Holm.

178. Cassida Hablitziæ.

Steven. Victor Bull. de Moscou, 1838, p. 182, t. iii. f. e. *Bohem.* Mon. Cassid. ii. p. 490, 149.
Cassida Testudo, *Friwaldsky*, Suffr. Stett. Ent. Zeit. 1844, p. 284.
Patr. Armenia et Caucasus.

179. Cassida seraphina.

Menetriès, Mem. de l'Acad. de Sciences de St. Petersbourg N. Ser. Scienc. Nat. t. v. p. 49, 235. *Küster*, Käf. Europ. vii. p. 84. *Bohem.* Mon. Cassid. ii. p. 491, 150.
Cassida Testudo, var. *Suffr.* Stett. Ent. Zeit. 1846, p. 364.
Patr. Græcia, Turcia, Brussa Asiæ minoris.

Species mihi ignotæ.

180. Cassida desertorum.

Gebler, Bull. de Moscou, 1833, vi. p. 305, 1848, No. 3, p. 8. *Bohem.* Mon. Cassid. ii. p. 492, 151.
Patr. Sibiria et Kirguisia.

181. Cassida Sibirica.

Gebler, Bull. de Moscou, 1833, vi. p. 306, 1848, No. 3, p. 9. *Bohem.* Mon. Cassid. ii. p. 493, 152.
Patr. Utkamengorsk, Sibiriæ.

182. Cassida undecimnotata.

Gebler, Bull. de Moscou, 1833, vi. p. 306, 1831, p. 615, 13, 1848, No. 3, p. 10. *Bohem.* Mon. Cassid. ii. p. 493, 153.
Patr. Loktewsk, Sibiriæ.

183. Cassida mera.

Germ. Linn. Entom. iii. p. 246, 183. *Bohem.* Mon. Cassid. ii. p. 494, 154.
Patr. Adelaide, Novæ Hollandiæ.

184. Cassida inquinata.

Brullé, Expédit. Scient. de Morée, iii. Part. i. p. 266, 523, t. 44, f. 3. *Bohem.* Mon. Cassid. ii. p. 494, 155.
Patr. Morea.

Genus 33. ASTERIZA.

Chevrolat, Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. ii. p. 495.
Cassida, *Oliv.* Ent. vi. p. 922.

1. Asteriza flavicornis. B.M.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. ii. p. 496, 1, t. vi. F.
Cassida flavicornis, *Oliv.* Ent. vi. p. 257, 55, 97, t. i. f. 5.
Patr. St. Domingo, Guyana.

Genus 34. HYBOSA.

Chevrolat, Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 1.

1. Hybosa gibbera. B.M.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 2, 1.
Patr. Brasilia (*Ent. Club*).

2. Hybosa indecens.

Bohem. Mon. Cassid. iii. p. 3, 2.
Patr. Brasilia.

3. Hybosa fornicata.

Bohem. Mon. Cassid. iii. p. 4, 3.
Patr. Bolivia.

4. Hybosa Cynthia. B.M.?

Bohem. Mon. Cassid. iii. p. 5, 4, t. vii. A.
Patr. Brasilia.

5. Hybosa galbanata.

Bohem. Mon. Cassid. iii. p. 6, 5.
Patr. Columbia.

Genus 35. CHARIDOTIS.

Bohem. Mon. Cassid. iii. p. 7.
Cassida, *Fabr.* Syst. El. i. p. 387. *Guér.* Mag. de Zool. 1831.
Coptocycla, *Dej.* Cat. ed. 3, p. 396.

I. *Elytra flavescentia, roseo-marginata.*

1. Charidotis miniata. B.M.

Bohem. Mon. Cassid. iii. p. 8, 1.
Coptocycla miniata, *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia (*Ent. Club*).

II. *Elytra dorso antico sanguineo, postico nigro, fascia transversa flava interjacente.*

2. Charidotis aurulenta. B.M.

Fabr. Syst. El. i. p 393, 31. *Schœnh.* Syn. Ins. ii. p. 217, 39.
Bohem. Mon. Cassid. iii. p. 9, 2.
Patr. Cayenna. Amazon (*Tapayos*), *H. W. Bates.*

3. Charidotis tricolor.

Guér. Mag. de Zool. 1831, Ins. pl. 31. *Bohem.* Mon. Cassid. iii. p. 10, 3.
Patr. Cayenna.

III. *Elytra flavescentia, singulo vitta antica lata sanguinea lateri parallela, postice arcu communi nigro.*

4. Charidotis abrupta. B.M.

Bohem. Mon. Cassid. iii. p. 11, 4, t. vii. B.
Patr. Para, Brasiliæ (Coll. *H. W. Bates*).

IV. *Elytra flavescentia, dorso annulo nigro.*

5. Charidotis luteola. B.M.

Bohem. Mon. Cassid. iii. p. 12, 5.
Patr. Columbia, Para (Coll. *H. W. Bates*).

6. Charidotis Dahlbomi.

Bohem. Mon. Cassid. iii. p. 13, 6.
Patr. Brasilia.

7. Charidotis orbifera.

Bohem. Mon. Cassid. iii. p. 14, 7.
Patr. Brasilia ad flumen Amazonum.

8. Charidotis soror.

Bohem. Mon. Cassid. iii. p. 15, 8.
Patr. Brasilia.

9. Charidotis ambigua.

Bohem. Mon. Cassid. iii. p. 17, 9.
Coptocycla ambigua, *Dej.* Cat. ed. 3, p. 396.
Patr. Cayenna.

10. Charidotis nucleata.

Bohem. Mon. Cassid. iii. p. 18, 10.
Patr. Brasilia.

V. *Elytra dorso sanguinea aut rufescentia.*

11. Charidotis coccinea.

Bohem. Mon. Cassid. iii. p. 19, 11.
Coptocycla coccinea, *Dej.* Cat. ed. 3, p. 397.
Patr. Cayenna.

12. Charidotis cognata.

Bohem. Mon. Cassid. iii. p. 20, 12.
Coptocycla cognata, *Dej.* Cat. ed. 3, p. 396.
Patr. Cayenna.

13. Charidotis cinctella.

Bohem. Mon. Cassid. iii. p. 21, 13.
Coptocycla cinctella, *Dej.* Cat. ed. 3, p. 397.
Patr. Brasilia, Rio Janeiro.

14. Charidotis spreta.

Bohem. Mon. Cassid. iii. p. 22, 14.
Patr. Brasilia.

VI. *Prothorax et elytra nigra, ferrugineo- aut flavo-marginata.*

15. Charidotis rubrocincta.

Bohem. Mon. Cassid. iii. p. 23, 15.
Patr. Brasilia.

16. Charidotis circumscripta. B.M.

Bohem. Mon. Cassid. iii. p. 24, 16.
Coptocycla circumscripta, *Dej.* Cat. ed. 3, p. 397.
Patr. Brasilia, Rio Janeiro.

17. Charidotis marginalis.

Bohem. Mon. Cassid. iii. p. 25, 17.
Patr. Brasilia.

VII. *Prothorax et elytra nigra, illo antice, hæc margine basali flava.*

18. Charidotis clypeolata.

Bohem. Mon. Cassid. iii. p. 26, 18.
Patr. Brasilia.

19. Charidotis seminulum.

Bohem. Mon. Cassid. iii. p. 27, 19.
Patr. Mexico.

VIII. *Prothorax et elytra nigra, illo antice flavo-marginato.*

20. Charidotis atramentosa.

Bohem. Mon. Cassid. iii. p. 28, 20.
Patr. Brasilia, Chiquitos.

21. Charidotis furva.

Bohem. Mon. Cassid. iii. p. 29, 21.
Patr. Brasilia, Cayenna.

IX. *Elytra dorso nigra, sanguinea vel ferruginea, flavo-maculata.*

22. Charidotis ocularis.

Bohem. Mon. Cassid. iii. p. 30, 22.
Coptocycla ocularis, *Dej.* Cat. ed. 3, p. 397.
Patr. Cassapava et St. Leopoldo Brasiliæ.

23. Charidotis gemellata.

Bohem. Mon. Cassid. iii. p. 31, 23.
Coptocycla gemellata, *Dej.* Cat. ed. 3. p. 397.
Patr. St. Leopoldo Brasiliæ, Brasilia meridionalis.

24. Charidotis centromaculata.

Bohem. Mon. Cassid. iii. p. 32, 24.
Coptocycla centromaculata, *Dej.* Cat. ed. 3, p. 397.
Patr. Brasilia.

25. Charidotis fenestralis.

Bohem. Mon. Cassid. iii. p. 33, 25.
Patr. Brasilia.

26. Charidotis admirabilis.

Bohem. Mon. Cassid. iii. p. 34, 26.
Patr. Brasilia.

27. Charidotis bellula.

Bohem. Mon. Cassid. iii. p. 35, 27.
Patr. Brasilia, Rio Janeiro.

28. Charidotis tessellata.

Bohem. Mon. Cassid. iii. p. 37, 28.
Coptocycla tessellata, *Dej.* Cat. ed. 3, p. 397.
Patr. Brasilia.

29. Charidotis scabriuscula.

Bohem. Mon. Cassid. iii. p. 38, 29.
Patr. Columbia.

30. CHARIDOTIS PLICATULA.

Bohem. Mon. Cassid. iii. p. 39, 30.
Patr. Brasilia.

31. CHARIDOTIS AUROGUTTATA.

Bohem. Mon. Cassid. iii. p. 40, 31.
Patr. Bolivia, Brasilia.

X. *Elytra flavescentia, sutura vittaque disci exterioris apicem versus arcuata ibique suturam attingente, nigris.*

32. CHARIDOTIS NIGROCINCTA. B.M.

Bohem. Mon. Cassid. iii. p. 41, 32.
Coptocycla nigrocincta, *Dej.* Cat. ed. 3, p. 397.
Patr. Brasilia.

33. CHARIDOTIS BICINCTA.

Bohem. Mon. Cassid. iii. p. 42, 33.
Patr. Bahia.

34. CHARIDOTIS BIMARGINATA.

Bohem. Mon. Cassid. iii. p. 43, 34.
Patr. Brasilia.

35. CHARIDOTIS EXIGUA.

Bohem. Mon. Cassid. iii. p. 44, 35.
Patr. Columbia.

36. CHARIDOTIS TANTILLA.

Bohem. Mon. Cassid. iii. p. 45, 36.
Patr. Bahia, Brasiliæ.

XI. *Elytra flavescentia, prothorax basi plaga triangulari, nigra.*

37. CHARIDOTIS DISCICOLLIS.

Bohem. Mon. Cassid. iii. p. 46, 37.
Coptocycla discicollis, *Dej.* Cat. ed. 3, p. 397.
Patr. Nova Granada, Cayenna.

XII. *Elytra sordide dilute virescentia, pone medium macula communi ferruginea.*

38. CHARIDOTIS GESTATRIX.

Bohem. Mon. Cassid. iii. p. 47, 38.
Coptocycla gestatrix, *Dej.* Cat. ed. 3, p. 396.
Patr. Cayenna, Para.

XIII. *Prothorax et elytra flavescentia vel virescentia, immaculata.*

39. CHARIDOTIS FLAVICANS.

Bohem. Mon. Cassid. iii. p. 48, 39.
Coptocycla flavicans, *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia.

40. CHARIDOTIS PUNCTATOSTRIATA.

Bohem. Mon. Cassid. iii. p. 49, 40.
Patr. Bolivia, Chiquitos.

41. CHARIDOTIS MARGINELLA. B.M.

Bohem. Mon. Cassid. iii. p. 50, 41.
Cassida marginella, *Fabr.* Syst. El. i. p. 392, 20. Ent. Syst. i. p. 294, 12. Mant. Ins. i. p. 63, 11. Spec. Ins. i. p. 108, 8. Syst. Entom. p. 89, 6. *Linn.* Syst. Nat. Gmel. i. iv. p. 1637, 42. *Herbst.* Naturs. viii. p. 330, 91. *Oliv.* Encycl. meth. v. p. 380, 15. *Schœnh.* Syn. Ins. ii. p. 216, 28.
Coptocycla thalassina, *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia.

42. CHARIDOTIS HERBIDA.

Bohem. Mon. Cassid. iii. p. 51, 42.
Patr. Saltogrande, Brasiliæ.

43. CHARIDOTIS GLOMEROSA.

Bohem. Mon. Cassid. iii. p. 52, 43.
Patr. Para Brasiliæ.

44. CHARIDOTIS ROTUNDATA.

Bohem. Mon. Cassid. iii. p. 53, 44.
Patr. Mexico.

45. CHARIDOTIS PYGMÆA.

Bohem. Mon. Cassid. iii. p. 54, 45.
Cassida pygmæa, *Klug*, Preis-Verzeichn. 1829, p. 8, 103.
Coptocycla pygmæa, *Dej.* Cat. ed. 3, p. 397.
Patr. Brasilia.

Genus 36. LACCOPTERA.

Bohem. Mon. Cassid. iii. p. 55.
Cassida, *Fabr.* Syst. El. i. p. 387. Dej. Cat. ed. 3, p. 398.
Deloyala, Chevrolat. *Dej.* Cat. ed. 3, p. 395.

I. *Prothorax et elytra ferruginea immaculata.*

1. Laccoptera excavata. B.M.

Bohem. Mon. Cassid. iii. p. 56, 1.
Cassida excavata, *Dej.* Cat. ed. 3, p. 398.
Patr. Caput Bonæ Spei, Caffraria. Port Natal (Coll. *Gueinzius*).

2. Laccoptera corrugata.

Bohem. Mon. Cassid. iii. p. 57, 2.
Cassida corrugata, *Sahlb.* Spec. Ins. p. 60, 40, t. iii. f. 6. *Dej.* Cat. ed. 3, p. 398.
Patr. Sierra Leona.

3. Laccoptera picea. B.M.?

Bohem. Mon. Cassid. iii. p. 58, 3.
Patr. Africa meridionalis.

4. Laccoptera intricata. B.M.

Bohem. Mon. Cassid. iii. p. 59, 4.
Cassida intricata, *Klug*, Ermanns Reise Ins. p. 47, 177.
Patr. Caput Bonæ Spei, Fernando Po (Old Calabar), Coll. *Murray*.

5. Laccoptera cancellata.

Bohem. Mon. Cassid. iii. p. 60, 5.
Patr. Caffraria, Caput Bonæ Spei.

6. Laccoptera ruginosa. B.M.

Bohem. Mon. Cassid. iii. p. 61, 6.
Patr. Caffraria (*Ent. Club*).

II. *Prothorax et elytra nigra vel nigro-picea, illo antice pallido-marginato.*

7. Laccoptera cicatricosa.

Bohem. Mon. Cassid. iii. p. 62, 7.
Cassida cicatricosa, *Dej.* Cat. ed. 3, p. 398.
Patr. Senegallia.

III. *Prothorax et elytra rufo-testacea, hæc vitta disci inæquali, nigro-cærulea.*

8. Laccoptera spectrum.

Bohem. Mon. Cassid. iii. p. 63, 8.
Patr. Madagascar.

IV. *Prothorax et elytra rufo-testacea, nigro-maculata.*

9. Laccoptera quatuordecimnotata. B.M.

Bohem. Mon. Cassid. iii. p. 64, 9.
Patr. India orientalis (*Walter Elliott*, Esq.).

10. Laccoptera vigintisexnotata.

Bohem. Mon. Cassid. iii. p. 66, 10.
Patr. Assam, Indiæ orientalis.

11. Laccoptera novemdecimnotata. B.M.

Bohem. Mon. Cassid. iii. p. 67, 11, t. vii. C.
Deloyala novemdecimnotata, *Buquet*, Dej. Cat. ed. 3, p. 395.
Patr. Java, Assam, Indiæ orientalis.

12. Laccoptera hospita. B.M.

Bohem. Mon. Cassid. iii. p. 68, 12.
Patr. China (Hong Kong), (Lieut.-Col. *Champion*).

13. Laccoptera multinotata. B.M.

Bohem. Mon. Cassid. iii. p. 70, 13.
Patr. Java.

14. Laccoptera Chinensis. B.M.

Bohem. Mon. Cassid. iii. p. 71, 14.
Cassida Chinensis, *Fabr.* Syst. El. i. p. 402, 82. Suppl. p. 84, 42, 43. *Herbst.* Naturs. viii. p. 353, 126. *Schœnh.* Syn. Ins. ii. p. 224, 109.
Patr. China.

15. Laccoptera impressa.

Bohem. Mon. Cassid. iii. p. 72, 15.
Aspidomorpha impressa, *Hombr.* et *Jaquin.* Voy. d'Urville, t. xviii. f. 15.
Patr. Baie Triton, Novæ Hollandiæ.

16. Laccoptera tredecimpunctata. B.M.

Bohem. Mon. Cassid. iii. p. 73, 16.
Cassida tredecimpunctata, *Fabr.* Syst. El. i. p. 398, 59. *Schœnh.* Syn. Ins. ii. p. 221, 80.
Cassida miliaris, *Herbst.* Naturs. viii. p. 312, 76, t. 135, f. 8?
Patr. Java, Assam, Indiæ orientalis. China (Hong Kong).

17. Laccoptera quadrimaculata.

Bohem. Mon. Cassid. iii. p. 75, 17.
Cassida quadrimaculata, *Thunb.* Nov. Ins. Spec. p. 86, t. v. f. 94. *Herbst.* Naturs. viii. p. 263, 33, t. cxxxii. f. 5. *Schœnh.* Syn. Ins. ii. p. 221, 92.
Patr. China.

18. Laccoptera Nepalensis.

Bohem. Mon. Cassid. iii. p. 76, 18.
Patr. Nepal, Indiæ orientalis.

19. Laccoptera discreta.

Bohem. Mon. Cassid. iii. p. 77, 19.
Patr. Java.

20. Laccoptera sculpturata.

Bohem. Mon. Cassid. iii. p. 78, 20.
Patr. Celebes.

21. Laccoptera Philippinensis. B.M.

Bohem. Mon. Cassid. iii. p. 79, 21.
Aspidomorpha Philippinensis, *Hombr.* et *Jaquin.* Voy. d'Urville, t. xviii. f. 14.
Patr. Manilla, Bombay, Indiæ orientalis. Sumatra.

Genus 37. PSALIDONOTA.

Bohem. Mon. Cassid. iii. p. 81.
Cassida, *Klug*, Preis-Verzeichn. 1829, p. 14.
Coptocycla, *Dej.* Cat. ed. 3, p. 396.
Deloyala, *Dej.* Cat. ed. 3, p. 395.

1. Psalidonota contempta. B.M.

Bohem. Mon. Cassid. iii. p. 82, 1.
Deloyala contempta, *Dej.* Cat. ed. 3, p. 395.
Patr. Cuba, Brasilia, Chiquitos.

2. Psalidonota ærata.

Bohem. Mon. Cassid. iii. p. 83, 2.
Patr. Brasilia.

3. Psalidonota atriceps.

Bohem. Mon. Cassid. iii. p. 84, 3.
Patr. Brasilia.

4. PSALIDONOTA CANDENS.

Bohem. Mon. Cassid. iii. p. 85, 4.
Patr. Brasilia.

5. PSALIDONOTA LEPROSA. B.M.?

Bohem. Mon. Cassid. iii. p. 86, 5.
Coptocycla dorsopunctata, var. *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia.

6. PSALIDONOTA DORSOPUNCTATA. B.M.

Bohem. Mon. Cassid. iii. p. 87, 6, t. vii. D.
Cassida dorsopunctata, *Klug*, Preis-Verzeichn. 1829, p. 14, 202.
Coptocycla dorsopunctata, *Dej.* Cat. ed. 3, p. 396.
Patr. Insula Cuba. Mexico (*Ent. Club*).

7. PSALIDONOTA OBTUSA.

Bohem. Mon. Cassid. iii. p. 88, 7.
Patr. Brasilia.

8. PSALIDONOTA CIRCUMSPICUA.

Bohem. Mon. Cassid. iii. p. 89, 8.
Patr. Brasilia.

Genus 38. COPTOCYCLA.

Chevrolat, Dej. Cat. ed. 3. p. 396. *Bohem.* Mon. Cassid. iii. p. 90.
Cassida, *Linn.* Syst. Nat. i. ii. p. 574. *Fabr.* Syst. El. i. p. 387. *De Geer*, Ins. v. p. 178. *Herbst.* Naturs. viii. p. 204. *Oliv.* Ent. vi. p. 922. *Schœnh.* Syn. Ins. ii. p. 209.
Deloyala, *Dej.* Cat. ed. 3, p. 395 (ex parte).

I. Unguiculi mutici vel basi obsolete unidentati.

A. *Elytra pone basin non gibbosa.*

1. *Angulus humeralis antrorsum parum prominulus.*

† *Elytra parum convexa (Majores, varicolores).*

1. COPTOCYCLA COMPLANATA. B.M.

Bohem. Mon. Cassid. iii. p. 92. 1.
Deloyala complanata, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia.

2. COPTOCYCLA JUSTINI.

Bohem. Mon. Cassid. iii. p. 93, 2.
Patr. Bogota.

3. Coptocycla picta. B.M.

Bohem. Mon. Cassid. iii. p. 94, 3.
Deloyala picta, *Dej.* Cat. ed. 3. p. 395.
Patr. Brasilia, Rio Grande (*J. C. Bowring*, Esq.).

4. Coptocycla postica.

Bohem. Mon. Cassid. iii. p. 95, 4.
Deloyala picta, *Dej.* Cat. ed. 3, p. 395, *part.*
Patr. Brasilia.

5. Coptocycla bifenestra. B.M.

Bohem. Mon. Cassid. iii. p. 97, 5.
Deloyala bifenestra, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia, (Rio Doce) *J. C. Bowring*, Esq.

6. Coptocycla Maura. B.M.

Bohem. Mon. Cassid. iii. p. 98. 6.
Deloyala Maura, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia.

7. Coptocycla varicolor.

Bohem. Mon. Cassid. iii. p. 99, 7.
Aspidomorpha varians, *Guér.-Ménéville*, Icon. du Règne Anim. Ins. texte ii. p. 291.
Patr. Bolivia, Chiquitos.

8. Coptocycla Blanchardi.

Bohem. Mon. Cassid. iii. p. 101, 8.
Patr. Provincia Cusco Peruviæ.

9. Coptocycla Lucasi.

Bohem. Mon. Cassid. iii. p. 102, 9.
Patr. Brasilia.

10. Coptocycla nubilosa.

Bohem. Mon. Cassid. iii. p. 103, 10.
Patr. Columbia.

†† *Elytra supra modice vel valde convexa.*

a. *Elytra flavescentia, nigro-maculata.*

11. Coptocycla quadrinotata.

Bohem. Mon. Cassid. iii. p. 104, 11.
Patr. Bahia, Brasiliæ.

12. COPTOCYCLA BICOLON. B.M.

Bohem. Mon. Cassid. iii. p. 105, 12.
Cassida bicolon, *Germ.* Ins. Spec. i. p. 542, 733.
Coptocycla quadra, *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia.

13. COPTOCYCLA ROSEOCINCTA. B.M.

Bohem. Mon. Cassid. iii. p. 106, 13, t. vii. E.
Cassida roseocincta, *Vigors*, Zool. Journ. ii. p. 241, t. ix. f. 10.
Patr. Brasilia, Bahia.

14. COPTOCYCLA BISBIPUSTULATA.

Bohem. Mon. Cassid. iii. p. 107, 14.
Patr. Bahia, Brasiliæ.

15. COPTOCYCLA UNDECIMPUNCTATA. B.M.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 108, 15.
Cassida undecimpunctata, *Fabr.* Syst. El. i. p. 397, 55. Ent. Syst. i. p. 298, 28. Mant. Ins. i. p. 64, 25. Spec. Ins. i. p. 110, 20. *Linn.* Syst. Nat. Gmel. i. iv. p. 1639, 55. *Voet.* Col. ii. t. xliii. f. 16. *Herbst.* Naturs. viii. p. 335, 98. *Oliv.* Encycl. meth. v. p. 384, 33. Entomol. vi. p. 994, 34, 97, t. iv. f. 67. a, b. *Schœnh.* Syn. Ins. ii. p. 221, 79.
Cassida arcuata, *Oliv.* Encycl. meth. v. p. 393, 82. Entom. vi. p. 945, 35, 97, t. ii. f. 26. *Fabr.* Syst. El. i. p. 398, 61. Ent. Syst. i. p. 298. *Herbst.* Naturs. viii. p. 336, 100. *Schh.* Ins. ii. p. 221, 82.
Patr. Cayenna, Guyana.

16. COPTOCYCLA STIGMA. B.M.

Bohem. Mon. Cassid. iii. p. 110, 16.
Cassida stigma, *Germ.* Ins. Spec. i. p. 542, 734.
Coptocycla trimaculata, *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia, Rio Janeiro.

17. COPTOCYCLA VENTRALIS. B.M.

Bohem. Mon. Cassid. iii. p. 111, 17.
Patr. India orientalis ad Neelgheries. Madras (*Walter Elliott*, Esq.).

18. COPTOCYCLA BISTRIMACULATA.

Bohem. Mon. Cassid. iii. p. 112, 18.
Patr. Tranquebar.

19. Coptocycla bistrinotata.

Bohem. Mon. Cassid. iii. p. 113, 19.
Patr. Bengalia.

20. Coptocycla sexmaculata. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 114, 20.
Patr. Java, Madras, India orientalis.

21. Coptocycla sexnotata. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 115, 21.
Cassida sexnotata, *Fabr.* Syst. El. i. p. 394, 35. Suppl. p. 82, 18, 19. *Oliv.* Ent. vi. p. 962, 66, 97, t. v. f. 84? *Schœnh.* Syn. Ins. ii. p. 217, 45.
Var. Cassida bipunctata, *Fabr.* Syst. El. i. p. 408, 119. Ent. Syst. i. p. 307, 69. Mant. Ins. i. p. 66, 58. Spec. Ins. i. p. 115, 49. Syst. Ent. p. 93, 33. *Linn.* Syst. Nat. i. ii. p. 578, 29. Ed. Gmel. i. iv. p. 1643, 29. *Herbst.* Naturs. viii. p. 271, 43, t. cxxxiii. f. 2? *Schœnh.* Syn. Ins. ii. p. 229, 151.
Patr. India orientalis, Coromandel, Ceylon (*Walter Elliott*, Esq.).

22. Coptocycla undecimnotata.

Bohem. Mon. Cassid. iii. p. 116, 22.
Patr. India orientalis.

23. Coptocycla septemdecimpunctata.

Bohem. Mon. Cassid. iii. p. 117, 23.
Patr. India orientalis.

24. Coptocycla bipunctipennis.

Bohem. Mon. Cassid. iii. p. 118, 24.
Patr. Travancour.

25. Coptocycla bisignata.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 119, 25.
Patr. America borealis.

26. Coptocycla subnotata.

Bohem. Mon. Cassid. iii. p. 120, 26.
Patr. Bolivia.

27. Coptocycla polita.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 121, 27.
Cassida polita, *Klug*, Preis-Verzeichn. 1829, p. 8, 99.
Patr. Brasilia.

b. *Elytra flavescentia, nigro-maculata, sutura nigra.*

* *Prothorax immaculatus.*

28. Coptocycla dolosa.

Bohem. Mon. Cassid. iii. p. 122, 28.
Patr. Bolivia.

29. Coptocycla Bowringii. B.M.

Bohem. Mon. Cassid. iii. p. 123, 29.
Patr. China ad Hong Kong (*J. C. Bowring*, Esq.).

30. Coptocycla scalaris. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 124, 30.
Cassida scalaris, *Fabr.* Syst. El. i. p. 391, 19. *Weber.* Observ. Ent. i. p. 51. 4. *Oliv.* Entom. vi. p. 967, 76, 97, t. vi. f. 94. *Schœnh.* Syn. Ins. ii. p. 216, 27.
Patr. Insula Java, Sumatra.

31. Coptocycla Javanica.

Bohem. Mon. Cassid. iii. p. 125, 31.
Patr. Java.

** *Prothorax nigro-maculatus.*

32. Coptocycla adamantina. B.M.

Bohem. Mon. Cassid. iii. p. 126, 32.
Cassida adamantina, *Germ.* Ins. Spec. i. p. 543, 735.
Coptocycla decemnotata, *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia, Venezuela (Coll. *Dyson*).

33. Coptocycla tredecimsignata. B.M.

Bohem. Mon. Cassid. iii. p. 128, 33.
Patr. Ceylon, China.

34. Coptocycla tredecimnotata. B.M.

Bohem. Mon. Cassid. iii. p. 129, 34.
Patr. Ceylon.

35. Coptocycla promiscua.

Bohem. Mon. Cassid. iii. p. 130, 35.
Cassida sexnotata, *Herbst.* Col. viii. p. 271, 42, t. cxxxiii. f. 1.
Patr. India orientalis.

36. Coptocycla decempunctata.

Bohem. Mon. Cassid. iii. p. 131, 36.
Patr. Bolivia, Para.

c. *Elytra flavescentia, nigro-maculata, sutura ferruginea.*

37. Coptocycla fastidiosa.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 132, 37.
Patr. Brasilia.

38. Coptocycla septemnotata. B.M.

Bohem. Mon. Cassid. iii. p. 133, 38.
Patr. India orientalis? Venezuela (B.M.).

39. Coptocycla ornata. B.M.

Bohem. Mon. Cassid. iii. p. 134, 39.
Cassida ornata, *Fabr.* Syst. El. i. p. 392, 23. Suppl. p. 81, 13, 14. *Herbst.* Naturs. viii. p. 350, 121. *Schœnh.* Syn. Ins. ii. p. 218, 31.
Patr. India orientalis, Coromandel, Ceylon.

d. *Elytra flavescentia vel testacea, maculis minoribus fuscis vel olivaceis ornata.*

40. Coptocycla bifossulata. B.M.?

Bohem. Mon. Cassid. iii. p. 135, 40.
Patr. Mexico.

41. Coptocycla sexpunctata. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 136, 41.
Cassida sexpunctata, *Fabr.* Syst. El. i. p. 395, 39. Ent. Syst. i. p. 296, 19. Mant. Ins. i. p. 63, 16. Spec. Ins. i. p. 109, 13. *Linn*, Syst. Nat. Gmel. i. iv. p. 1636, 37. *Schœnh.* Syn. Ins. ii. p. 218, 55.
Cassida bistripustulata, *Herbst.* Naturs. viii. p. 256, 25, t. cxxxi. f. 10.
Patr. Brasilia, Rio Janeiro.

42. Coptocycla trisignata.

Bohem. Mon. Cassid. iii. p. 138, 42.
Patr. Mexico ad Yucatan.

43. Coptocycla ternata.

Bohem. Mon. Cassid. iii. p. 139, 43.
Patr. Brasilia.

44. Coptocycla biimpressa. B.M.

Bohem. Mon. Cassid. iii. p. 140, 44.
Patr. Mexico ad Alvarado.

45. Coptocycla bistripunctata. B.M.

Bohem. Mon. Cassid. iii. p. 141, 45.
Cassida sexpunctata, *Oliv.* Encycl. meth. v. p. 382, 22. Entom. vi. p. 961, 65, 97, t. iii. f. 42.
Cassida bistripunctata, *Herbst.* Naturs. viii. p. 275, 47? t. cxxxiii. f. 6. *Schœnh.* Syn. Ins. ii. p. 218, 56.
Patr. Cayenna, Mexico, Brasilia.

46. Coptocycla aurichalcea. B.M.

Bohem. Mon. Cassid. iii. p. 142, 46.
Cassida aurichalcea. *Fabr.* Syst. El. i. p. 397, 53. *Illig.* Mag. iii. p. 157, 58. *Oliv.* Entom. vi. p. 965, 71, 97, t. v. f. 89. *Schœnh.* Syn. Ins. ii. p. 220, 76.
Var. Cassida bicolor, *Fabr.* Syst. El. i. p. 396, 48. Suppl. p. 83, 24, 25. *Herbst.* Naturs. viii. p. 352, 125.
Patr. America borealis. "St. John's Bluff, E. Florida," (*E. Doubleday*, Esq.).

47. Coptocycla bivulnerata.

Bohem. Mon. Cassid. iii. p. 143, 47.
Patr. Mexico.

48. Coptocycla bisbinotata.

Bohem. Mon. Cassid. iii. p. 144, 48.
Patr. Insula Cuba.

e. *Elytra flavescentia, plagis vel vittis olivaceis aut brunneis ornata.*

49. Coptocycla auricoma.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 145, 49.
Patr. Brasilia.

50. Coptocycla læta.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 146, 50.
Patr. Brasilia.

51. Coptocycla subplagiata.

Bohem. Mon. Cassid. iii. p. 147, 51.
Patr. Brasilia.

52. Coptocycla bifasciata.

Bohem. Mon. Cassid. iii. p. 148, 52.
Cassida bifasciata, *Linn.* Syst. Nat. i. ii. p. 576, 10. Ed. Gmel.

i. iv. p. 1639, 10. *Fabr.* Syst. El. i. p. 394, 36. *Schœnh.* Syn. Ins. ii. p. 217, 46.
Cassida obliterata, *Herbst.* Naturs. viii. p. 260, 29, t. cxxxii. f. 1.
Patr. Brasilia.

53. Coptocycla nigromaculata.

Bohem. Mon. Cassid. iii. p. 149, 53.
Cassida nigromaculata, *De Geer*, Mem. Ins. v. p. 189, 15, t. xx. f. 18.
Coptocycla mendax, *Dej.* Cat. ed. 3, p. 397.
Patr. Cayenna, Surinam.

54. Coptocycla quadrisignata.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 150, 54.
Var. Coptocycla livida, *Chevrol.* in litter.
Patr. Cuba, St. Domingo.

55. Coptocycla latevittata.

Bohem. Mon. Cassid. iii. p. 151, 55.
Patr. Insula, St. Thomas.

56. Coptocycla præusta.

Bohem. Mon. Cassid. iii. p. 152, 56.
Patr. St. Domingo.

57. Coptocycla glaucovittata.

Bohem. Mon. Cassid. iii. p. 153, 57.
Cassida glaucovittata, *Erichs.* Archiv. f. Naturg. xiii. p. 154, 3.
Patr. Peruviæ montes.

58. Coptocycla submaculata.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 154, 58.
Patr. Brasilia.

59. Coptocycla inconstans.

Bohem. Mon. Cassid. iii. p. 155, 59.
Coptocycla Zona var. *Dej.* Cat. ed. 3. p. 397.
Patr. Cayenna.

60. Coptocycla fumosa.

Bohem. Mon. Cassid. iii. p. 156, 60.
Patr. Cayenna.

61. Coptocycla liquida.

Bohem. Mon. Cassid. iii. p. 157, 61.
Cassida liquida, *Erichs.* Arch. f. Naturg. xiii. p. 154, 4.
Patr. Peruvia.

62. Coptocycla Hogbergi.

Bohem. Mon. Cassid. iii. p. 158, 62.
Patr. Mexico ad Vera Cruz.

63. Coptocycla vinula.

Bohem. Mon. Cassid. iii. p. 159, 63.
Patr. Columbia.

f. *Elytra dorso olivacea, annulo flavescente.*

64. Coptocycla Zona. B.M.

Dej. Cat. ed. 3, p 397. *Bohem.* Mon. Cassid. iii. p. 160, 64.
Cassida Zona, *Fabr.* Syst. El. i. p. 393, 28.
Cassida annulus, *Oliv.* Encycl v. p. 381, 19. Entom. vi. p. 969, 80, 97, t. i. f. 14. *Herbst.* Naturs. viii. p. 272, 44, t. cxxxiii. f. 3, c. *Illig.* Mag. i. p. 406, 34. *Schœnh.* Syn. Ins. ii. p. 217, 44, ex parte.
Patr. Cayenna, Venezuela (Coll. *Dyson*).

g. *Elytra flavescentia, dorso annulo nigro vel rubro ultra medium continuato.*

65. Coptocycla circumducta. B.M.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 161, 65.
Patr. Brasilia, Rio Grande (*J. C. Bowring,* Esq.).

66. Coptocycla procincta.

Bohem. Mon. Cassid. iii. p. 162, 66.
Patr. Brasilia.

67. Coptocycla aurifera. B.M.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 163, 67.
Patr. Brasilia.

68. Coptocycla speculum.

Bohem. Mon. Cassid. iii. p. 164, 68.
Patr. Bahia.

69. Coptocycla neglecta. B.M.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 165, 69.
Patr. Brasilia.

70. Coptocycla oculata.

Bohem. Mon. Cassid. iii. p. 166, 70.
Patr. Brasilia.

71. Coptocycla surda.

Bohem. Mon. Cassid. iii. p. 167, 71.
Patr. Bahia, Brasiliæ.

72. Coptocycla circulus.

Mannerh. Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 168, 72.
Patr. Brasilia.

73. Coptocycla Cyclops.

Bohem. Mon. Cassid. iii. p. 169, 73.
Cassida Cyclops, *Fabr.* Syst. El. i. p. 394, 33. *Schœnh.* Syn. Ins. ii. p. 217, 43.
Patr. Brasilia.

74. Coptocycla circinata.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 171, 74.
Patr. Brasilia.

75. Coptocycla consentanea. B.M.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 172, 75.
Patr. Brasilia.

76. Coptocycla connexa.

Bohem. Mon. Cassid. iii. p. 173, 76.
Patr. Bahia.

77. Coptocycla mansueta.

Bohem. Mon. Cassid. iii. p. 174, 77.
Patr. Brasilia.

78. Coptocycla ancilla.

Bohem. Mon. Cassid. iii. p. 175, 78.
Patr. Chiquitos et St. Cruz, Bolivia.

79. Coptocycla rubrodiscoidalis.

Bohem. Mon. Cassid. iii. p. 176, 79.
Patr. Bolivia.

80. Coptocycla similata.

Bohem. Mon. Cassid. iii. p. 177, 80.
Patr. Brasilia interior.

81. Coptocycla sellata.

Bohem. Mon. Cassid. iii. p. 178, 81.
Patr. St. Paul, Brasiliæ.

82. Coptocycla purpurea.

Bohem. Mon. Cassid. iii. p. 179, 82.
Cassida purpurea, *Linn.* Syst. Nat. i. ii. p. 576, 12. Ed. Gmel. i. iv. p. 1639, 12. *Fabr.* Syst. El. i. p. 393, 32. *De Geer*, Ins. v. p. 190, 16, t. 15, f. 19. *Thunb.* Nov. Ins. Spec. p. 87, t. v. f. 98. *Herbst.* Naturs. viii. p. 266, t. cxxxii. f. 9. *Oliv.* Encycl. meth. Ins. v. p. 381, 20. Entomol. vi. p. 966, 73, 97, t. vi. f. 91. *Schœnh.* Syn. Ins. ii. p. 217, 40, 41.
Coptocycla concentrica, *Dej.* Cat. ed. 3, p. 397.
Patr. Cayenna.

83. Coptocycla myops.

Bohem. Mon. Cassid. iii. p. 180, 83.
Patr. Columbia, Guyana Britann.

84. Coptocycla puella.

Bohem. Mon. Cassid. iii. p. 181, 84.
Patr. Mexico ad Teapa.

85. Coptocycla repanda.

Bohem. Mon. Cassid. iii. p. 182, 85.
Patr. Chiquitos, Brasiliæ interioris.

86. Coptocycla discoidalis.

Bohem. Mon. Cassid. iii. p. 183, 86.
Patr. Brasilia.

87. Coptocycla cyclographa.

Bohem. Mon. Cassid. iii. p. 184, 87.
Patr. Cayenna.

88. Coptocycla consobrina.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 185, 88.
Patr. Columbia ad Carthagena.

h. *Elytra flavescentia, dorso annulo nigro vel rubro ad medium vix continuato.*

89. Coptocycla semiannulata.

Bohem. Mon. Cassid. iii. p. 186, 89.
Patr. St. Martha, Brasiliæ.

90. Coptocycla serena.

Bohem. Mon. Cassid. iii. p. 187, 90.
Patr. Brasilia.

91. Coptocycla involuta.

Bohem. Mon. Cassid. iii. p. 188, 91.
Patr. Confinia fluminis Amazonum.

i. *Elytra flavescentia, dorso annulo nigro vel rubro, postice ad suturam continuato.*

92. Coptocycla amœna.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 189, 92.
Patr. Guatemala.

93. Coptocycla opulenta.

Bohem. Mon. Cassid. iii. p. 191, 93.
Patr. Mexico.

94. Coptocycla incerta. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 191, 94.
Patr. Brasilia.

k. *Elytra flavescentia, dorso plaga nigra, macula flava litteram X fere referente notata.*

95. Coptocycla aucta.

Bohem. Mon. Cassid. iii. p. 193, 95.
Patr. Columbia.

l. *Elytra flavescentia, singulo vitta nigra vel rubra, postice inflexa, suturam attingente.*

* *Sutura infuscata.*

96. Coptocycla semivittata. B.M.

Bohem. Mon. Cassid. iii. p. 194, 96.
Patr. Old Calabar (" Celebes," Bohem. *err.*).

97. Coptocycla Aubei. B.M.

Bohem. Mon. Cassid. iii. p. 195, 97.
Patr. Old Calabar.

98. Coptocycla æquinoctialis. B.M.

Bohem. Mon. Cassid. iii. p. 196, 98. *Dej.* Cat. ed. 3, p. 397. Cassida æquinoctialis, *Oliv.* Entom. iv. p. 966, 74, 97, t. vi. f. 92.
Patr. Senegallia, Oware et Benin (Old Calabar, Coll. *A. Murray*).

** *Sutura non infuscata.*

99. Coptocycla gyrata.

Bohem. Mon. Cassid. iii. p. 197, 99.
Patr. Brasilia.

100. Coptocycla forcipata.

Bohem. Mon. Cassid. iii. p. 198, 100.
Patr. Brasilia.

101. Coptocycla amœnula.

Bohem. Mon. Cassid. iii. p. 199, 101.
Patr. Brasilia.

102. Coptocycla proxima.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 200, 102.
Patr. Mexico.

103. Coptocycla erratica.

Bohem. Mon. Cassid. iii. p. 201, 103.
Patr. Mexico, Columbia.

m. *Elytra dorso nigra vel ferruginea, annulo communi flavo.*

104. Coptocycla Wesmæli.

Bohem. Mon. Cassid. iii. p. 202, 104.
Patr. Mexico.

105. Coptocycla pupillata.

Bohem. Mon. Cassid. iii. p. 203, 105.
Patr. Brasilia.

106. Coptocycla circumflexa.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 204, 106.
Patr. Brasilia.

107. Coptocycla concentrica.

Bohem. Mon. Cassid. iii. p. 205, 107.
Patr. Brasilia.

108. Coptocycla cincticula.

Bohem. Mon. Cassid. iii. p. 206, 108.
Patr. Brasilia.

109. Coptocycla annularis.

Bohem. Mon. Cassid. iii. p. 207, 109.
Patr. Brasilia.

110. Coptocycla annulifera.

Bohem. Mon. Cassid. iii. p. 208, 110.
Patr. Rio Janeiro Brasiliæ.

111. Coptocycla consimilis.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 209, 111.
Patr. Cayenna.

112. Coptocycla Signoreti.

Bohem. Mon. Cassid. iii. p. 210, 112.
Patr. Brasilia ad Rio Janeiro.

n. *Elytra dorso nigra, castanea vel sanguinea, margine flavescente.*

113. Coptocycla circulata.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 211, 113.
Patr. Cayenna.

114. Coptocycla bicingulata.

Bohem. Mon. Cassid. iii. p. 212, 114.
Patr. Chiquitos.

115. Coptocycla Diomma.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 213, 115.
Cassida Diomma, *Boisd.* Faune de l'Océan. p. 540, 7.
Patr. Nova Guinea.

116. Coptocycla ephippiata. B.M.?

Bohem. Mon. Cassid. iii. p. 214, 116.
Patr. Mexico.

117. Coptocycla clitelligera.
Bohem. Mon. Cassid. iii. p. 215, 117.
Patr. Brasilia.

118. Coptocycla furunculus. B.M.
Bohem. Mon. Cassid. iii. p. 216, 118.
Patr. Brasilia.

119. Coptocycla villica.
Bohem. Mon. Cassid. iii. p. 217, 119.
Patr. Brasilia.

120. Coptocycla coccinelloides.
Bohem. Mon. Cassid. iii. p. 218, 120.
Patr. Brasilia ad flumen Amazonum.

121. Coptocycla profana.
Bohem. Mon. Cassid. iii. p. 219, 121.
Patr. Brasilia.

122. Coptocycla posticata.
Bohem. Mon. Cassid. iii. p. 220, 122.
Patr. Brasilia.

123. Coptocycla conclusa.
Bohem. Mon. Cassid. iii. p. 221, 123.
Patr. Brasilia.

124. Coptocycla crispata.
Bohem. Mon. Cassid. iii. p. 222, 124.
Patr. Brasilia.

o. *Elytra dorso nigra vel rubra, fascia transversa flava.*

125. Coptocycla Leprieuri. B.M.
Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 223, 125.
Patr. Cayenna, Honduras (Coll. *Dyson*). Ega, on Amazon (Coll. *Bates*).

126. Coptocycla aurofasciata. B.M.
Bohem. Mon. Cassid. iii. p. 224, 126.
Cassida aurofasciata, *Erichs.* Archiv. f. Naturg. xiii. p. 154, 9.
Patr. Cayenna, Brasilia, Peruvia.

127. Coptocycla bipartita.

Bohem. Mon. Cassid. iii. p. 225, 127.
Patr. Cayenna, Nova Granada.

128. Coptocycla succincta. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 226, 128.
Patr. Brasilia.

129. Coptocycla Jousselini.

Bohem. Mon. Cassid. iii. p. 227, 129.
Patr. Nova Granada.

130. Coptocycla semifasciata.

Bohem. Mon. Cassid. iii. p. 228, 130.
Patr. Columbia.

p. *Elytra flavescentia, dorso plagis magnis nigris vel sanguineis ornata.*

131. Coptocycla Fahræi.

Bohem. Mon. Cassid. iii. p. 229, 131.
Patr. Honduras.

132. Coptocycla Waterhousei.

Bohem. Mon. Cassid. iii. p. 230, 132.
Patr. Bahia.

133. Coptocycla formosa.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 231, 133.
Patr. Brasilia.

134. Coptocycla virgata.

Bohem. Mon. Cassid. iii. p. 232, 134.
Patr. Cassapava.

135. Coptocycla roseopicta.

Bohem. Mon. Cassid. iii. p. 233, 135.
Patr. Brasilia.

q. *Elytra dorso nigra, annulo antico et arcu postico flavis.*

136. Coptocycla arcuata. B.M.

Bohem. Mon. Cassid. iii. p. 234, 136.
Cassida arcuata, *Swederus* Acta Holm. 1787, p. 193, 12. *Linn.*

Syst. Nat. Gmel. i. iv. p. 1644, 41. *Schœnh.* Syn. Ins. ii. p. 230, 159.
Coptocycla lunifera, var. *Dej.* Cat. ed. 3, p. 396.
Patr. Novum Friburgum Brasiliæ, St. Paul.

137. Coptocycla lunifera.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 235, 137.
Patr. Brasilia.

138. Coptocycla annulus. B.M.

Bohem. Mon. Cassid. iii. p. 236, 138.
Cassida annulus, *Fabr.* Syst. El. i. p. 394, 34. Ent. Syst. i. p. 296, 18. Mant. Ins. i. p. 63, 15. Spec. Ins. i. p. 109, 12. *Linn.* Syst. Nat. Gmel. i. iv. p. 1637, 38.
Cassida circularis, *Oliv.* Ent. vi. p. 970, 81, 97, t. vi. f. 98.
Coptocycla circularis, *Dej.* Cat. ed. 3, p. 396.
Patr. Cayenna, Bolivia, Columbia Para, Guatemala (Coll. *Scherzer*).

139. Coptocycla liturata.

Bohem. Mon. Cassid. iii. p. 237, 139.
Patr. Rio Janeiro, Brasilia.

140. Coptocycla voluptaria.

Bohem. Mon. Cassid. iii. p. 238, 140.
Patr. Bolivia.

141. Coptocycla contexta.

Bohem. Mon. Cassid. iii. p. 239, 141.
Patr. Brasilia.

r. *Elytra dorso nigra vel ferruginea, flavo- vel testaceo-maculata.*

142. Coptocycla sexguttata.

Bohem. Mon. Cassid. iii. p. 240, 142. *Dej.* Cat. ed. 3, p. 397.
Cassida sexguttata, *Boisd.* Faune Ent. de l'Océan. ii. p. 539, 6.
Patr. Nova Guinea.

143. Coptocycla Nympha.

Bohem. Mon. Cassid. iii. p. 241, 143.
Patr. Brasilia interior.

144. Coptocycla separata.

Bohem. Mon. Cassid. iii. p. 242, 144.
Patr. Brasilia.

145. Coptocycla Bremei.

Bohem. Mon. Cassid. iii. p. 243, 145.
Patr. Columbia, Bogota.

146. Coptocycla quadrioculata.

Bohem. Mon. Cassid. iii. p. 244, 146.
Patr. Bolivia.

147. Coptocycla constricta.

Bohem. Mon. Cassid. iii. p. 245, 147.
Patr. Brasilia.

148. Coptocycla mamillata.

Bohem. Mon. Cassid. iii. p. 246, 148.
Patr. Brasilia.

149. Coptocycla Zelleri.

Bohem. Mon. Cassid. iii. p. 247, 149.
Patr. Columbia.

150. Coptocycla ambigena.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 248, 150.
Patr. Brasilia.

151. Coptocycla glyphica.

Bohem. Mon. Cassid. iii. p. 250, 151.
Patr. Havanna.

152. Coptocycla stillata.

Bohem. Mon. Cassid. iii. p. 251, 152.
Patr. Brasilia ad Rio Janeiro.

153. Coptocycla dodonea.

Bohem. Mon. Cassid. iii. p. 252, 153.
Patr. Brasilia.

154. Coptocycla congregata.

Bohem. Mon. Cassid. iii. p. 253, 154.
Patr. Brasilia.

155. Coptocycla punctaria. B.M.

Bohem. Mon. Cassid. iii. p. 254, 155.

Cassida punctaria, *Fabr.* Syst. El. i. p. 392, 21. *Oliv.* Ent. vi. p. 965, 72, 97, t. vi. f. 90. *Schœnh.* Syn. Ins. ii. p. 216, 29.
Cassida punctata, *Weber*, Observ. Ent. i. p. 51, 3.
Coptocycla pardalina, *Dej.* Cat. ed. 3, p. 397.
Patr. Java (Dr. *Horsfield*), Sumatra (Coll. *Raffles*), Tennaserim (*J. D. C. V. Packman*, Esq.).

156. Coptocycla leopardina.

Bohem. Mon. Cassid. iii. p. 255, 156.
Patr. Madagascar.

157. Coptocycla vesicularis.

Bohem. Mon. Cassid. iii. p. 256, 157.
Cassida vesicularis, *Thunb.* Nov. Insect. Spec. p. 88. *Herbst.* Naturs. viii. p. 266, 38. *Schœnh.* Syn. Ins. ii. p. 217, 42.
Patr. Japonia.

158. Coptocycla Australica.

Bohem. Mon. Cassid. iii. p. 257, 158.
Patr. Nova Hollandia, ad Swan River.

159. Coptocycla pustulata.

Bohem. Mon. Cassid. iii. p. 258, 159.
Patr. Cayenna, Para Brasiliæ.

160. Coptocycla subirrorata.

Bohem. Mon. Cassid. iii. p. 259, 160.
Patr. Columbia.

161. Coptocycla extensa.

Bohem. Mon. Cassid. iii. p. 261, 161.
Patr. Mexico.

162. Coptocycla catenata. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 262, 162.
Patr. Java, Bengalia, Ceylon.

163. Coptocycla viridinotata.

Bohem. Mon. Cassid. iii. p. 263, 163.
Patr. Java.

164. Coptocycla Goudoti.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 264, 164.
Patr. Madagascar.

165. Coptocycla verrucata.

Bohem. Mon. Cassid. iii. p. 265, 165.
Patr. Madagascar.

166. Coptocycla physodes.

Bohem. Mon. Cassid. iii. p. 266, 166.
Patr. Sumatra.

167. Coptocycla Ceylonica. B.M.

Bohem. Mon. Cassid. iii. p. 267, 167.
Patr. Ceylon.

168. Coptocycla microcera.

Bohem. Mon. Cassid. iii. p. 268, 168.
Patr. Bahia.

169. Coptocycla plagiata.

Bohem. Mon. Cassid. iii. p. 269, 169.
Patr. Bahia.

s. *Elytra flavescentia, dorso annulo antico et arcu vel macula postica nigris aut sanguineis.*

170. Coptocycla elegans.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 270, 170.
Patr. Brasilia.

171. Coptocycla excelsa.

Bohem. Mon. Cassid. iii. p. 271, 171.
Patr. Brasilia.

172. Coptocycla vana.

Bohem. Mon. Cassid. iii. p. 272, 172.
Patr. Brasilia.

173. Coptocycla virginalis.

Bohem. Mon. Cassid. iii. p. 273, 173.
Patr. Bahia.

174. Coptocycla biarcuata.

Bohem. Mon. Cassid. iii. p. 274, 174.
Patr. Brasilia.

175. Coptocycla rubropicta.

Bohem. Mon. Cassid. iii. p. 275, 175.
Patr. Rio Grande Brasiliæ.

176. Coptocycla roseosignata. B.M.

Bohem. Mon. Cassid. iii. p. 276, 176.
Patr. Brasilia.

t. *Elytra flavescentia, singulo linea angusta sanguinea vel nigra, medio introrsum flexa.*

177. Coptocycla conspicillata. B.M.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 276, 177.
Patr. Brasilia.

178. Coptocycla cayennensis.

Bohem. Mon. Cassid. iii. p. 278, 178.
Patr. Cayenna.

u. *Elytra flavescentia, dorso nigro-vel rubro-vittata.*

179. Coptocycla circumdata. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 279, 179.
Cassida circumdata, *Herbst.* Naturs. viii. p. 268, 40, t. cxxxii. f. 11. *Oliv.* Ent. vi. p. 967, 75, 97, t. vi. f. 93. *Schœnh.* Syn. Ins. ii. p. 218, 50. *Boisd.* Faune Entom. de l'Océan. ii. p. 536, 1.
Patr. India orientalis, China, Nova Guinea.

180. Coptocycla trivittata. B.M.

Bohem. Mon. Cassid. iii. p. 280, 180.
Cassida trivittata, *Fabr.* Syst. El. i. p. 397, 50. *Oliv.* Ent. vi. p. 973, 87, 97, t. vi. f. 103. *Schœnh.* Syn. Ins. ii. p. 220, 70. *Boisd.* Faune Ent. de l'Océan. ii. p. 544, 5.
Aspidomorpha trivittata, *Hombr.* et *Jaquin.* Voy. d'Urville, t. xviii. f. 16.
Patr. Insula Java, Manilla (Ins. Phillipp. Coll. *Cuming*). Nova Cambria, N. China (Coll. *Fortune*).

181. Coptocycla fuscovittata.

Bohem. Mon. Cassid. iii. p. 281, 181.
Patr. Maldonado.

182. Coptocycla vittipennis.

Bohem. Mon. Cassid. iii. p. 282, 182.
Patr. Bolivia.

183. COPTOCYCLA RUBROVITTATA.

Bohem. Mon. Cassid. iii. p. 283, 183.
Patr. Porto Allegro Brasiliæ.

184. COPTOCYCLA VARIANS.

Bohem. Mon. Cassid. iii. p. 284, 184.
Cassida varians, *Herbst.* Naturs. viii. p. 269, 41, t. cxxxii. f. 12, e.
Schœnh. Syn. Ins. ii. p. 218, 51.
Patr. India orientalis.

v. *Elytra dorso nigra vel ferruginea, flavo-vittata.*

185. COPTOCYCLA QUADRIVITTATA.

Bohem. Mon. Cassid. iii. p. 285, 185.
Patr. Brasilia.

186. COPTOCYCLA FLAVOVITTATA.

Bohem. Mon. Cassid. iii. p. 286, 186.
Patr. Minas Geræs Brasiliæ.

187. COPTOCYCLA SUBVITTATA.

Bohem. Mon. Cassid. iii. p. 287, 187.
Patr. Bolivia.

188. COPTOCYCLA FLAVOLINEATA.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 288, 188.
Cassida flavolineata, *Latr.* in Humboldt et Bonpland Observ.
de Zool. et d'Anatom. ii. p. 34, t. ii. f. 7.
Patr. Columbia.

189. COPTOCYCLA PROPINQUA. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 289, 189.
Patr. Columbia (Magdeleine), Coll. *Goudot.*

190. COPTOCYCLA COMPUNCTA.

Bohem. Mon. Cassid. iii. p. 290, 190.
Patr. Nova Hollandia.

x. *Elytra testacea, singulo vitta lata marginali, nigra.*

191. COPTOCYCLA VALIDA.

Bohem. Mon. Cassid. iii. p. 291, 191.
Patr. Brasilia.

192. Coptocycla Drewseni.

Bohem. Mon. Cassid. iii. p. 292, 192.
Patr. Rio de la Plata.

y. *Elytra dorso fusco-flavoque variegata.*

193. Coptocycla judaica. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 293, 193.
Cassida judaica, *Fabr.* Syst. El. i. p. 392, 25. Ent. Syst. i. p. 295, 15. Mant. Ins. i. p. 63, 14. Spec. Ins. i. p. 109, 11. *Linn.* Syst. Nat. Gmel. i. iv. p. 1637, 39. *Herbst.* Naturs. viii. p. 280, 52, t. cxxxiii. f. 11, k. *Oliv.* Encycl. v. p. 381, 18. Entomol. vi. p. 968, 78, 97, t. vi. f. 96. *Schœnh.* Syn. Ins. ii. p. 216, 34.
Var. Coptocycla operosa, *Dej.* Cat. olim.
Patr. Brasilia, Santarem (*H. W. Bates*), Cayenna, Guyana, Columbia (Coll. *Goudot*).

194. Coptocycla conflagrata.

Bohem. Mon. Cassid. iii. p. 294, 194.
Patr. Cuba.

195. Coptocycla stupidula.

Bohem. Mon. Cassid. iii. p. 295, 195.
Patr. Brasilia.

z. *Elytra testacea, rugulis nonnullis flavescentibus instructa.*

196. Coptocycla flavorugosa.

Bohem. Mon. Cassid. iii. p. 296, 196.
Patr. Rio Janeiro Brasiliæ.

aa. *Elytra sanguinea vel testacea, macula magna, oblonga, laterali, flavescente-hyalina.*

197. Coptocycla incurva.

Bohem. Mon. Cassid. iii. p. 297, 197.
Patr. Brasilia.

198. Coptocycla emarginata. B.M.

Bohem. Mon. Cassid. iii. p. 298, 198.
Deloyala emarginata, *Dej.* Cat. ed. 3, p. 396.
Patr. Mexico, Guatemala (Coll. *Scherzer*).

199. Coptocycla annexa.

Bohem. Mon. Cassid. iii. p. 299, 199.
Patr. Mexico.

200. Coptocycla purpurata.

Bohem. Mon. Cassid. iii. p. 300, 200.
Deloyala purpurata, *Dej.* Cat. ed. 3, p. 396.
Patr. Missouri Americæ borealis.

201. Coptocycla bicurvata.

Bohem. Mon. Cassid. iii. p. 301, 201.
Patr. Cayenna.

202. Coptocycla connectens.

Bohem. Mon. Cassid. iii. p. 302, 202.
Patr. Rio Grande Brasiliæ meridionalis.

bb. *Elytra dorso plaga magna nigra vel ferruginea, sæpe flavo-rugosa, utrinque ramulos duos in marginem emittente.*

203. Coptocycla Zetterstedti.

Bohem. Mon. Cassid. iii. p. 303, 203.
Patr. Mexico.

204. Coptocycla decussata.

Bohem. Mon. Cassid. iii. p. 305, 204.
Patr. Mexico.

205. Coptocycla usta.

Bohem. Mon. Cassid. iii. p. 306, 205.
Patr. Cayenna.

206. Coptocycla crucipennis.

Bohem. Mon. Cassid. iii. p. 307, 206.
Patr. Mexico.

207. Coptocycla insubida.

Bohem. Mon. Cassid. iii. p. 308, 207.
Patr. Columbia.

208. Coptocycla munda.

Bohem. Mon. Cassid. iii. p. 309, 208.
Patr. Nova Valencia.

209. Coptocycla lateraria.

Bohem. Mon. Cassid. iii. p. 310, 209.
Patr. Brasilia ad fluvium Amazonum.

210. COPTOCYCLA GRANARIA.

Bohem. Mon. Cassid. iii. p. 311, 210.
Patr. Brasilia ad fluvium Amazonum.

211. COPTOCYCLA EFFUSA.

Aspidomorpha effusa, *Bohem.* Mon. Cassid. ii. p. 320, 74.
Patr. Java.

212. COPTOCYCLA PAGANA.

Bohem. Mon. Cassid. iii. p. 312, 211.
Patr. Java.

213. COPTOCYCLA BAHIANA.

Bohem. Mon. Cassid. iii. p. 313, 212.
Patr. Bahia.

cc. *Elytra dorso plaga nigra vel ferruginea, interdum flavo-variegata, utrinque basi ramulum in marginem emittente.*

214. COPTOCYCLA GUTTATA. B.M.

Bohem. Mon. Cassid. iii. p. 314, 213.
Cassida guttata, *Oliv.* Encycl. v. p. 383, 30. Entomol. vi. p. 955, 53, 97, t. i. f. 13, a. b. fig. mala.
Cassida cruciata, *Fabr.* Syst. El. i. p. 397, 52. Ent. Syst. i. p. 297, 26. Mant. Ins. i. p. 64, 23. Spec. Ins. i. p. 110, 18. Syst. Ent. p. 90, 12. *Herbst.* Naturs. viii. p. 354, 129. *Schœnh.* Syn. Ins. ii. p. 220, 75.
Var. Cassida signifer, *Herbst.* Naturs. viii. p. 313, 77, t. cxxxv. f. 9, c. *Schœnh.* Syn. Ins. ii. p. 221, 78.
Cassida cruciata, *De Geer*, Mem. des Ins. v. p. 187, 12 var.
Deloyala vicina, *Dej.* Cat. ed. 3, p. 395.
Patr. Jamaica (Coll. *Gosse*), Cuba, America borealis, Yucatan, Honduras (Coll. *Dyson*), ("Warm Springs, N. Carolina," *E. Doubleday*, Esq.), New Orleans, Delaware (*Ent. Club*), Canada (W.).

215. COPTOCYCLA HAMULATA.

Bohem. Mon. Cassid. iii. p. 316, 214.
Patr. St. Domingo.

216. COPTOCYCLA LUGUBRINA.

Bohem. Mon. Cassid. iii. p. 317, 215.
Patr. Mexico.

217. Coptocycla immunita.

Bohem. Mon. Cassid. iii. p. 318, 216.
Patr. Yucatan.

218. Coptocycla trabeata.

Bohem. Mon. Cassid. iii. p. 319, 217.
Deloyala trabeata, *Dej.* Cat. ed. 3, p. 395.
Patr. Mexico, America borealis.

219. Coptocycla immunda.

Bohem. Mon. Cassid. iii. p. 320, 218.
Patr. Mexico.

220. Coptocycla lucidula.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii p. 321, 219.
Patr. America borealis.

221. Coptocycla dorsalis.

Bohem. Mon. Cassid. iii. p. 322, 220.
Patr. Java.

dd. *Elytra testacea, dorso nigro-variegata, pone medium macula marginali nigra.*

222. Coptocycla Dohrni.

Bohem. Mon. Cassid. iii. p. 323, 221.
Patr. Mexico.

223. Coptocycla distorta.

Bohem. Mon. Cassid. iii. p. 324, 222.
Patr. Mexico.

ee. *Elytra testacea, dorso parce fusco-variegata.*

224. Coptocycla sordida.

Bohem. Mon. Cassid. iii. p. 326, 223.
Patr. Mexico.

ff. *Elytra flavescentia, sutura sanguinea.*

225. Coptocycla virgulata.

Bohem. Mon. Cassid. iii. p. 327, 224.
Patr. Mexico.

gg. *Elytra flavescentia, annulo antico obsoleto, dilute sanguineo.*

226. COPTOCYCLA PUDICA.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 328, 225.
Patr. Brasilia.

227. COPTOCYCLA VIRGO.

Bohem. Mon. Cassid. iii. p. 329, 226.
Patr. Salto grande, Brasiliæ.

hh. *Elytra virescentia, flavescentia vel rufo-testacea, immaculata.*

228. COPTOCYCLA UNIFORMIS.

Bohem. Mon. Cassid. iii. p. 330, 227.
Patr. Cayenna.

229. COPTOCYCLA CONCOLOR.

Bohem. Mon. Cassid. iii. p. 330, 228.
Patr. Para, Brasiliæ.

230. COPTOCYCLA SUBRUGOSA.

Bohem. Mon. Cassid. iii. p. 331, 229.
Patr. Brasilia.

231. COPTOCYCLA INCULTA.

Bohem. Mon. Cassid. iii. p. 332, 230.
Patr. Brasilia.

232. COPTOCYCLA GLAUCINA.

Bohem. Mon. Cassid. iii. p. 333, 231.
Patr. St. Jean, Portorico.

233. COPTOCYCLA LIMPIDA. B.M.

Bohem. Mon. Cassid. iii. p. 334, 232.
Patr. Brasilia, Mexico.

234. COPTOCYCLA CALVA.

Bohem. Mon. Cassid. iii. p. 335, 233.
Patr. Brasilia.

235. COPTOCYCLA MARCULENTA.

Bohem. Mon. Cassid. iii. p. 336, 234.
Patr. Mexico.

236. Coptocycla incorrupta.

Bohem. Mon. Cassid. iii. p. 337, 235.
Patr. Brasilia.

237. Coptocycla striatopunctata.

Bohem. Mon. Cassid. iii. p. 338, 236.
Patr. Bahia.

238. Coptocycla pallescens.

Bohem. Mon. Cassid. iii. p. 339, 237.
Patr. Rio Janeiro, Brasilia.

239. Coptocycla insculpta.

Bohem. Mon. Cassid. iii. p. 340, 238.
Patr. Yucatan.

240. Coptocycla exsanguis.

Bohem. Mon. Cassid. iii. p. 341, 239.
Patr. Brasilia.

241. Coptocycla glabricula.

Bohem. Mon. Cassid. iii. p. 341, 240.
Patr. St. Domingo.

242. Coptocycla emarcida.

Bohem. Mon. Cassid. iii. p. 342, 241.
Patr. Brasilia.

243. Coptocycla pellucida.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 343, 242.
Patr. Brasilia.

244. Coptocycla immaculata. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 344, 243.
Cassida immaculata, *Oliv.* Encycl. Ins. v. p. 382, 20. Entomol. vi. p. 969, 79, 97, t. vi. f. 97. *Voet.* Col. t. ii. f. 18.
Patr. Cayenna, Guyana et Brasilia.

245. Coptocycla sejuncta.

Bohem. Mon. Cassid. iii. p. 345, 244.
Patr. Brasilia.

246. Coptocycla immerita.

Bohem. Mon. Cassid. iii. p. 346, 245.
Patr. Columbia.

247. Coptocycla fulgida.

Bohem. Mon. Cassid. iii. p. 347, 246.
Patr. Java.

248. Coptocycla varicornis.

Bohem. Mon. Cassid. iii. p. 348, 247.
Patr. Mexico.

249. Coptocycla aurisplendens.

Mannerh. Bull. de Moscou, xxi. p. 285. *Bohem.* Mon. Cassid. iii. p. 349, 248.
Coptocycla sexpunctata, var. *Dej.* Cat. ed. 3, p. 397.
Patr. California, Mexico.

250. Coptocycla succinea.

Bohem. Mon. Cassid. iii. p. 350, 249.
Patr. Mexico.

251. Coptocycla meticulosa.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 351, 250.
Patr. Mexico.

252. Coptocycla crenata.

Bohem. Mon. Cassid. iii. p. 352, 251.
Patr. Brasilia.

253. Coptocycla herbea. B.M.

Bohem. Mon. Cassid. iii. p. 353, 252.
Patr. Brasilia.

254. Coptocycla tenella.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 354, 253.
Cassida tenella. *Klug,* Preis-Verz. 1829, p. 8, 102.
Patr. Brasilia.

255. Coptocycla gibberosa.

Bohem. Mon. Cassid. iii. p. 355, 254.
Patr. Salto grande, Brasilia.

256. Coptocycla inedita.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 356, 255.
Patr. Brasilia.

257. Coptocycla vilis.

Bohem. Mon. Cassid. iii. p. 357, 256.
Patr. Yucatan.

258. Coptocycla punctipennis.

Bohem. Mon. Cassid. iii. p. 358, 257.
Patr. Brasilia.

259. Coptocycla globosa.

Bohem. Mon. Cassid. iii. p. 358, 258.
Patr. Guyana Britannica.

260. Coptocycla pullula.

Bohem. Mon. Cassid. iii. p. 359, 259.
Patr. Brasilia.

2. *Angulus humeralis antrorsum valde productus.*
a. *Elytra flavescentia, annulo nigro notata.*

261. Coptocycla præcincta. B.M.

Bohem. Mon. Cassid. iii. p. 360, 260.
Patr. Brasilia.

262. Coptocycla designata.

Bohem. Mon. Cassid. iii. p. 361, 261.
Patr. Brasilia.

263. Coptocycla Deyrollei. B.M.

Bohem. Mon. Cassid. iii. p. 363, 262.
Patr. St. Catharina, Brasiliæ, Columbia.

b. *Elytra dorso nigra, maculis duabus magnis longitudinalibus, flavis.*

264. Coptocycla centralis.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 364, 263.
Patr. Brasilia.

265. Coptocycla amplexa.

Bohem. Mon. Cassid. iii. p. 365, 264.
Cassida amplexa, *Erichs.* Arch. f. Naturg. xiii. p. 154, 10.
Patr. Peruvia.

c. *Elytra dorso nigra vel castanea.*

266. Coptocycla Diana.

Bohem. Mon. Cassid. iii. p. 366, 265.
Patr. Mexico.

267. Coptocycla mœsta.

Bohem. Mon. Cassid. iii. p. 367, 266.
Patr. Nova Granada.

d. *Elytra dorso nigra vel ferruginea, annulo flavo.*

268. Coptocycla Phœbe.

Bohem. Mon. Cassid. iii. p. 367, 267.
Patr. Cayenna.

269. Coptocycla inscripta.

Bohem. Mon. Cassid. iii. p. 368, 268.
Patr. Brasilia.

e. *Elytra dorso nigra, flavo-vel testaceo-maculata.*

270. Coptocycla incita.

Bohem. Mon. Cassid. iii. p. 369, 269.
Patr. Columbia.

271. Coptocycla clandestina.

Bohem. Mon. Cassid. iii. p. 370, 270.
Patr. Brasilia?

272. Coptocycla Buqueti. B.M.

Bohem. Mon. Cassid. iii. p. 371, 271.
Patr. Columbia (Coll. *Goudot*).

273. Coptocycla bis-triguttata.

Bohem. Mon. Cassid. iii. p. 372, 272.
Patr. Nova Valencia, Venezuela, Caraccas.

274. Coptocycla bis-bimaculata.

Bohem. Mon. Cassid. iii. p. 373, 273.
Patr. Columbia.

275. Coptocycla ludicra.

Bohem. Mon. Cassid. iii. p. 374, 274.
Patr. Brasilia.

276. Coptocycla Sahlbergi.

Bohem. Mon. Cassid. iii. p. 375, 275.
Patr. Brasilia.

277. Coptocycla scenica.

Bohem. Mon. Cassid. iii. p. 377, 276.
Patr. Brasilia.

278. Coptocycla dorsosignata.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 378, 277.
Patr. Brasilia interior.

279. Coptocycla multisignata. B.M.

Bohem. Mon. Cassid. iii. p. 379, 278.
Patr. Brasilia.

f. *Elytra dilute flavo-virescentia, immaculata.*

280. Coptocycla flavescens. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 380, 279.
Coptocycla porracea, *Dej.* Cat. ed. 3, p. 397.
Patr. Brasilia.

B. *Elytra pone basin gibbosa.*

1. *Angulus humeralis antrorsum modice prominulus.*

a. *Elytra plaga magna dorsali saturatiore, utrinque ramulos duos in marginem emittente.*

281. Coptocycla Andreæ. B.M.

Bohem. Mon. Cassid. iii. p. 381, 280.
Deloyala Crux, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia.

282. Coptocycla bisinuata.

Bohem. Mon. Cassid. iii. p. 382, 281.
Patr. Chiquitos.

283. Coptocycla nitida.

Bohem. Mon. Cassid. iii. p. 383, 282.
Deloyala nitida, *Dej.* Cat. ed. 3, p. 396.
Patr. Mexico.

284. Coptocycla tuberculata. B.M.

Bohem. Mon. Cassid. iii. p. 385, 283.
Cassida tuberculata, *Fabr.* Syst. El. i. p. 399, 67. Ent. Syst. i.

p. 299, 33. Mant. Ins. i. p. 64, 26. Syst. Ent. p. 96, 13. *Linn.* Syst. Nat. Gmel. i. iv. p. 1640, 14. *Herbst.* Naturs. viii. p. 338, 103. *Schœnh.* Syn. Ins. ii. p. 221, 91.
Deloyala acuticornis, *Chevrol.* Dej. Cat. ed. 3, p. 396.
Patr. Mexico, Venezuela (Coll. *Dyson*).

285. Coptocycla lobata. B.M.

Bohem. Mon. Cassid. iii. p. 386, 284.
Patr. Mexico, Honduras (Coll. *Dyson*).

286. Coptocycla rutila.

Bohem. Mon. Cassid. iii. p. 387, 285.
Cassida rutila, *Erichs.* Archiv. f. Naturg. xiii. p. 153, 1.
Patr. Peruvia.

287. Coptocycla rubicunda. B.M.

Bohem. Mon. Cassid. iii. p. 388, 286.
Deloyala rubicunda, *Dej.* Cat. ed. 3, p. 396.
Aspidomorpha rubicunda, *Guér.* Icon. de Règne Anim. Ins. texte ii. p. 291.
Patr. Brasilia.

288. Coptocycla radiata.

Bohem. Mon. Cassid. iii. p. 389, 287.
Patr. Nova Guinea.

289. Coptocycla testudinaria.

Bohem. Mon. Cassid. iii. p. 390, 288.
Patr. Mexico.

290. Coptocycla clavata. B.M.

Bohem. Mon. Cassid. iii. p. 391, 289.
Cassida clavata. *Fabr.* Syst. El. i. p. 398, 62. Suppl. p. 83, 30, 31. *Herbst.* Naturs. viii. p. 312, 75, t. cxxxv. f. 7, C. *Oliv.* Ent. vi. p. 963, 69, 97. tab. v. f. 87. *Schœnh.* Syn. Ins. ii. p. 221, 83.
Deloyala clavata, *Dej.* Cat. ed. 3, p. 395.
Patr. America borealis, Carolina (Coll. *Shephard*), *E. Doubleday*, Esq.

291. Coptocycla ramosa.

Bohem. Mon. Cassid. iii. p. 393, 290.
Patr. Columbia.

b. *Elytra plaga dorsali magna, saturatiore, utrinque basi ramulum in marginem emittente.*

292. Coptocycla ventricosa.

Bohem. Mon. Cassid. iii. p. 394, 291.
Patr. Mexico.

293. Coptocycla asperata.

Bohem. Mon. Cassid. iii. p. 395, 292.
Patr. Mexico.

294. Coptocycla cruciata. B.M.

Bohem. Mon. Cassid. iii. p. 396, 293.
Cassida cruciata, *Linn.* Syst. Nat. ed. xii. i. ii. p. 576, 9. Ed. Gmel. i. iv. p. 1639, 9. *De Geer.* Mem. des Ins. v. p. 187, 12, t. xv. f. 15. *Oliv.* Encycl. v. p. 383, 30. Entomol. vi. p. 955, 52, 97, t. v. f. 77.
Cassida crux, *Fabr.* Syst. El. i. p. 397, 54. Ent. Syst. i. p. 298, 27. Mant. Ins. i. p. 64, 24. Spec. Ins. i. p. 110, 19. *Illig.* Mag. i. p. 407, 54. *Herbst.* Naturs. viii. p. 308, 73, t. cxxxv. f. 5. *Linn.* Syst. Nat. ed. Gmel. i. iv. p. 1639, 54. *Oliv.* Encycl. v. p. 384, 32. *Voet.* Col. ii. t. 44, f. 27. *Schœnh.* Syn. Ins. ii. p. 220, 77.
Deloyala fuliginosa, *Dej.* Cat. ed. 3, p. 395.
Var. Cassida fuliginosa, *Oliv.* Ent. vi. p. 971, 83, 97, t. i. f. 8?
Patr. Cayenna, Bolivia, Columbia, Bahia, Surinam, Venezuela (Coll. *Dyson*), Brasilia (*R. Napo*, Coll. *H. W. Bates*).

295. Coptocycla hamata.

Bohem. Mon. Cassid. iii. p. 398, 294.
Patr. Brasilia.

c. *Elytra rubra, macula intramarginali pellucida.*

296. Coptocycla carnulenta.

Bohem. Mon. Cassid. iii. p. 399, 295.
Cassida carnulenta, *Erichs.* Archiv. f. Naturg. xiii. p. 154, 2.
Patr. Chiquitos, Nova Granada, Peruvia.

d. *Elytra testacea, interdum nigro-variegata, macula marginali albida.*

297. Coptocycla egregia. B.M.

Bohem. Mon. Cassid. iii. p. 400, 296.
Patr. Mexico.

298. Coptocycla vitreata. B.M.

Bohem. Mon. Cassid. iii. p. 401, 297.
Cassida vitreata, *Perty*, Delect. Anim. p. 103, t. xx. f. 13.
Deloyala vetusta, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia in montibus Prov. Minarum.

e. *Elytra testacea, singulo extus arcu nigro, antice et pone medium in marginem excurrente.*

299. Coptocycla flexuosa.

Bohem. Mon. Cassid. iii. p. 402, 298.
Patr. Bahia, Brasiliæ.

f. *Elytra flavescentia, nigro-maculata.*

300. Coptocycla Balyi. B.M.

Bohem. Mon. Cassid. iii. p. 403, 299.
Patr. Ceylon.

g. *Elytra dorso testacea, parce fusco-variegata.*

301. Coptocycla cribrosa.

Bohem. Mon. Cassid. iii. p. 404, 300.
Patr. Assam, Indiæ orientalis.

302. Coptocycla rugipennis.

Bohem. Mon. Cassid. iii. p. 405, 301.
Patr. Madagascar.

h. *Elytra dorso plaga magna castanea, pone medium ramulum in marginem emittente.*

303. Coptocycla Horsfieldi.

Bohem. Mon. Cassid. iii. p. 406, 302.
Patr. Java.

304. Coptocycla biramosa.

Bohem. Mon. Cassid. iii. p. 408, 303.
Patr. Pulo Penang.

305. Coptocycla luctifera.

Bohem. Mon. Cassid. iii. p. 408, 304.
Patr. St. Catharina, Brasiliæ.

i. *Elytra ferruginea, profunde, vage, nigro-punctata.*

306. COPTOCYCLA INTUMIDA.

Bohem. Mon. Cassid. iii. p. 410, 305.
Patr. Brasilia.

k. *Elytra flavescentia, singulo vitta lata nigra, pone medium arcuatim ad suturam ducta.*

307. COPTOCYCLA HILARIS.

Bohem. Mon. Cassid. iii. p. 411, 306.
Patr. Pulo Penang, Java.

l. *Elytra dorso ferruginea vel brunnea, flavo-vel rubro-maculata vel rugosa.*

308. COPTOCYCLA DISCOLOR.

Bohem. Mon. Cassid. iii. p. 412, 307.
Patr. Madagascar.

309. COPTOCYCLA RUBROORNATA.

Bohem. Mon. Cassid. iii. p. 413, 308.
Patr. Madagascar.

310. COPTOCYCLA VERSICOLOR.

Bohem. Mon. Cassid. iii. p. 414, 309.
Patr. China.

m. *Elytra flavescentia, annulo communi nigro vel sanguineo.*

311. COPTOCYCLA PLACIDA.

Bohem. Mon. Cassid. iii. p. 415, 310.
Patr. Cayenna.

312. COPTOCYCLA IRIS.

Bohem. Mon. Cassid. iii. p. 416, 311.
Cassida Iris, *Perty*, Del. Anim. Brasil. p. 102, t. xx. f. 12.
Patr. Bahia, Sebastianopolis.

313. COPTOCYCLA SANGUINOLENTA. B.M.

Bohem. Mon. Cassid. iii. p. 417, 312.
Cassida sanguinolenta, *Swederus*, Acta Holm. 1787, p. 193, 11. *Linn.* Syst. Nat. Gmel. i. iv. p. 1644, 41. *Schœnh.* Syn. Ins. ii. p. 230, 158.
Patr. Brasilia.

n. *Elytra flavescentia, annulo antico angusto lineaque disci exterioris pone medium introrsum valde flexa, nigris.*

314. COPTOCYCLA CONSTELLATA. B.M.

Bohem. Mon. Cassid. iii. p. 418, 313.
Patr. St. Cruz, Bolivia (Coll. *Bridges*).

o. *Elytra flavescentia, virescentia vel rufo-testacea, immaculata.*

315. COPTOCYCLA ELATIOR. B.M.

Bohem. Mon. Cassid. iii. p. 419, 314.
Cassida elatior, *Klug*, Preis-Verz. 1829, p. 8, 104.
Deloyala straminea, *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia (Maj.-Gen. *Hardwicke*).

316. COPTOCYCLA PALACEA.

Bohem. Mon. Cassid. iii. p. 420, 315.
Patr. Brasilia.

317. COPTOCYCLA PUNCTATISSIMA.

Bohem. Mon. Cassid. iii. p. 421, 316.
Patr. Brasilia.

318. COPTOCYCLA SUSPECTA.

Bohem. Mon. Cassid. iii. p. 422, 317.
Patr. Brasilia.

319. COPTOCYCLA RUBROTESTACEA.

Bohem. Mon. Cassid. iii. p. 423, 318.
Patr. Columbia.

320. COPTOCYCLA STULTA.

Bohem. Mon. Cassid. iii. p. 424, 319.
Patr. Brasilia.

321. COPTOCYCLA FALLAX. B.M.

Bohem. Mon. Cassid. iii. p. 425, 320.
Deloyala fallax, *Dej.* Cat. ed. 3, p. 396.
Cassida dilatata, *Billberg*, Mus. Childr.
Var. Deloyala perplexa, *Dej.* Cat. ed. 3, p. 396.
Patr. Brasilia (Coll. *Children*).

322. COPTOCYCLA OBESA.

Bohem. Mon. Cassid. iii. p. 426, 323.
Patr. Brasilia.

323. Coptocycla guadeloupensis.

Bohem. Mon. Cassid. iii. p. 427, 324.
Patr. Guadeloupe.

324. Coptocycla simplex.

Bohem. Mon. Cassid. iii. p. 428, 323.
Patr. Brasilia.

2. *Angulus humeralis antrorsum valde productus.*

325. Coptocycla vigens.

Bohem. Mon. Cassid. iii. p. 429, 324.
Patr. Rio Grande Brasiliæ meridionalis, Minas Geraes.

II. Unguiculi basi pectinati.

A. *Elytra basi ad scutellum non impressa.*

a. *Elytra flavescentia, parce nigro-vel fusco-maculata.*

326. Coptocycla glareosa.

Bohem. Mon. Cassid. iii. p. 430, 325.
Patr. Brasilia.

327. Coptocycla Cynaræ.

Bohem. Mon. Cassid. iii. p. 431, 326.
Patr. Montevideo.

328. Coptocycla subtincta.

Bohem. Mon. Cassid. iii. p. 432, 327.
Patr. Minas Geraes, Brasiliæ.

329. Coptocycla vitellina.

Bohem. Mon. Cassid. iii. p. 433, 328.
Patr. Brasilia.

b. *Elytra flavescentia, dorso nigro-vel ferrugineo-vittata.*

330. Coptocycla vittigera.

Bohem. Mon. Cassid. iii. p. 434, 329.
Patr. Brasilia.

331. Coptocycla rustica.

Bohem. Mon. Cassid. iii. p. 435, 330.
Patr. Porto Allegro, Brasiliæ.

332. Coptocycla rubrolineata.

Bohem. Mon. Cassid. iii. p. 436, 331.
Patr. Brasilia.

c. *Elytra flavescentia, dorso annulo communi, nigro vel rubro.*

333. Coptocycla nobilitata.

Bohem. Mon. Cassid. iii. p. 437, 332.
Patr. Brasilia.

334. Coptocycla magica.

Bohem. Mon. Cassid. iii. p. 438, 333.
Patr. Brasilia.

335. Coptocycla rotalis.

Bohem. Mon. Cassid. iii. p. 439, 334.
Patr. Brasilia.

336. Coptocycla nigroannulata.

Bohem. Mon. Cassid. iii. p. 440, 335.
Patr. Brasilia interior.

337. Coptocycla zonata.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 441, 336.
Patr. Brasilia.

338. Coptocycla porosa.

Bohem. Mon. Cassid. iii. p. 442, 337.
Patr. Brasilia.

339. Coptocycla centropunctata.

Bohem. Mon. Cassid. iii. p. 444, 338.
Patr. Columbia.

340. Coptocycla Servula.

Bohem. Mon. Cassid. iii. p. 445, 339.
Patr. Bolivia.

d. *Elytra dorso nigra, annulo communi flavo.*

341. Coptocycla mystica.

Bohem. Mon. Cassid. iii. p. 446, 340.
Patr. Cayenna.

342. Coptocycla circinaria.

Bohem. Mon. Cassid. iii. p. 447, 341.
Cassida circinaria, *Erichs.* Arch. f. Naturg. xiii. p. 154, 5.
Patr. Peruvia.

343. Coptocycla diffinis.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 448, 342.
Patr. Cayenna.

344. Coptocycla circumcincta.

Dej. Cat. ed. 3, p. 372. *Bohem.* Mon. Cassid. iii. p. 449, 343.
Patr. Bahia, Brasiliæ.

345. Coptocycla media.

Bohem. Mon. Cassid. iii. p. 450, 344.
Patr. Bahia.

e. *Elytra dorso nigra vel castanea, margine flavescenti.*

346. Coptocycla melanota.

Bohem. Mon. Cassid. iii. p. 451, 345.
Patr. Teapa in Mexico, Nova Granada.

347. Coptocycla flavocincta.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 452, 346.
Patr. Brasilia.

f. *Elytra dorso nigra, annulo antico et arcu postico flavis.*

348. Coptocycla Fairmairei.

Bohem. Mon. Cassid. iii. p. 453, 347.
Patr. Peruvia, Columbia.

349. Coptocycla signatifera.

Dej. Cat. ed. 3, p. 396. *Bohem.* Mon. Cassid. iii. p. 454, 348.
Coptocycla circularis, *Dej.* Cat. ed. 2, p. 116.
Patr. Brasilia.

350. Coptocycla gemina. B.M.

Bohem. Mon. Cassid. iii. p. 455, 349.
Patr. Para (Coll. *H. W. Bates*).

g. *Elytra dorso nigra, flavo-maculata.*

351. COPTOCYCLA AMICTA.

Bohem. Mon. Cassid. iii. p. 456, 350.
Patr. Cayenna.

352. COPTOCYCLA FLAVONOTATA.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 457, 351.
Patr. Columbia, Surinam.

353. COPTOCYCLA GUTTIGERA.

Bohem. Mon. Cassid. iii. p. 459, 352.
Patr. Brasilia.

h. *Elytra flavescentia, singulo arcu laterali nigro.*

354. COPTOCYCLA BLANDIDICA.

Bohem. Mon. Cassid. iii. p. 460, 353.
Patr. Guarayos.

i. *Elytra flavescentia vel virescentia, immaculata.*

355. COPTOCYCLA IMPURA.

Bohem. Mon. Cassid. iii. p. 460, 354.
Patr. Brasilia.

356. COPTOCYCLA VIRESCENS.

Bohem. Mon. Cassid. iii. p. 461, 355.
Patr. Brasilia.

357. COPTOCYCLA INDECORATA.

Bohem. Mon. Cassid. iii. p. 462, 356.
Patr. Rio Janeiro Brasiliæ.

358. COPTOCYCLA PALLIDULA.

Bohem. Mon. Cassid. iii. p. 463, 357.
Patr. Celebes.

359. COPTOCYCLA INGENUA.

Bohem. Mon. Cassid. iii. p. 464, 358.
Patr. Cassapa, Brasiliæ.

B. *Elytra basi ad scutellum impressa, sæpe gibbosa.*

a. *Elytra plaga magna dorsali communi, obscuriore, ramulos duos in marginem emittente.*

360. Coptocycla aciculata. B.M.

Bohem. Mon. Cassid. iii. p. 465, 359.
Deloyala aciculata, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia.

361. Coptocycla difficilis.

Bohem. Mon. Cassid. iii. p. 466, 360.
Deloyala difficilis, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia.

362. Coptocycla patruelis.

Bohem. Mon. Cassid. iii. p. 467, 361.
Deloyala patruelis, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia.

363. Coptocycla glaberata.

Bohem. Mon. Cassid. iii. p. 469, 362.
Patr. Brasilia.

364. Coptocycla optata.

Bohem. Mon. Cassid. iii. p. 470, 363.
Patr. Brasilia.

365. Coptocycla hectica. B.M.

Bohem. Mon. Cassid. iii. p. 471, 364.
Patr. Mexico.

366. Coptocycla maculosa.

Bohem. Mon. Cassid. iii. p. 472, 365.
Patr. St. Catharina, Brasiliæ.

367. Coptocycla fraterna.

Bohem. Mon. Cassid. iii. p. 473, 366.
Deloyala fraterna, *Dej.* Cat. ed. 3, p. 395.
Var. Coptocycla modesta, *Dej.* Cat. ed. 3, p. 397.
Patr. Columbia.

368. Coptocycla signaticollis.

Bohem. Mon. Cassid. iii. p. 475, 367.
Patr. Caraccas.

369. COPTOCYCLA STIGMATICA.

Bohem. Mon. Cassid. iii. p. 476, 368.
Patr. Bahia.

370. COPTOCYCLA ANXIA.

Bohem. Mon. Cassid. iii. p. 477, 369.
Patr. St. Catharina, Brasiliæ.

371. COPTOCYCLA LUGUBRIS.

Bohem. Mon. Cassid. iii. p. 478, 370.
Patr. Columbia.

b. *Elytra dorso obscuriora, singulo ante medium macula intramarginali pellucida.*

372. COPTOCYCLA DEROSA. B.M.

Bohem. Mon. Cassid. iii. p. 479, 371.
Deloyala derosa, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia.

373. COPTOCYCLA BIFENESTRATA.

Bohem. Mon. Cassid. iii. p. 480, 372.
Patr. Caraccas.

374. COPTOCYCLA BIPELLUCIDA.

Bohem. Mon. Cassid. iii. p. 481, 373.
Patr. Cayenna.

c. *Elytra dorso plaga magna communi, obscuriore, ramulum basalem in marginem emittente.*

375. COPTOCYCLA FOLIATA.

Bohem. Mon. Cassid. iii. p. 482, 374.
Deloyala foliata, *Dej.* Cat. ed. 3, 395.
Patr. Cayenna.

376. COPTOCYCLA HYPOCRITA.

Bohem. Mon. Cassid. iii. p. 483, 375.
Deloyala hypocrita, *Dej.* Cat. ed. 3, p. 395.
Patr. Brasilia.

377. COPTOCYCLA SEVERA.

Bohem. Mon. Cassid. iii. p. 485, 376.
Patr. Bahia.

d. *Elytra dorso plaga magna communi obscuriore, pone medium ramulum in marginem emittente.*

378. Coptocycla bilobata.

Bohem. Mon. Cassid. iii. p. 486, 377.
Patr. Mexico.

379. Coptocycla palmata.

Bohem. Mon. Cassid. iii. p. 487, 378.
Patr. Mexico.

380. Coptocycla dissimilis.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 488, 379.
Patr. Mexico.

e. *Elytra nigro-picea, angulo humerali margineque postico flavis.*

381. Coptocycla dolorosa.

Bohem. Mon. Cassid. iii. p. 489, 380.
Patr. Brasilia.

f. *Elytra dorso plaga magna communi obscuriore, margine flavescenti.*

382. Coptocycla divulsa.

Bohem. Mon. Cassid. iii. p. 490, 381.
Patr. Cayenna.

383. Coptocycla sepulchralis.

Bohem. Mon. Cassid. iii. p. 491, 382.
Patr. Bogota.

384. Coptocycla peltata.

Bohem. Mon. Cassid. iii. p. 492, 383.
Patr. Peruvia.

385. Coptocycla plicata.

Bohem. Mon. Cassid. iii. p. 493, 384.
Patr. America borealis, Mexico.

386. Coptocycla punicea.

Bohem. Mon. Cassid. iii. p. 494, 385.
Patr. Mexico.

387. Coptocycla scabra.

Bohem. Mon. Cassid. iii. p. 495, 386.
Patr. Brasilia.

g. *Elytra nigro-picea vel sanguinea, margine parum dilutiore.*

388. Coptocycla Morio.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 496, 387.
Cassida Morio, *Fabr.* Syst. El. i. p. 393, 29. *Schœnh.* Syn. Ins. ii. p. 217, 37.
Patr. America meridionalis.

389. Coptocycla Carnifex.

Bohem. Mon. Cassid. iii. p. 497, 388.
Patr. Mexico.

390. Coptocycla rubrocincta.

Bohem. Mon. Cassid. iii. p. 498, 389.
Patr. Caraccas, Mexico, St. Martha Brasiliæ.

h. *Elytra dorso nigra, flavo-rugosa.*

391. Coptocycla conscripta.

Bohem. Mon. Cassid. iii. p. 499, 390.
Patr. Cayenna.

392. Coptocycla papulosa.

Bohem. Mon. Cassid. iii. p. 500, 391.
Patr. Surinam, Cayenna.

393. Coptocycla intermedia. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 502, 392.
Patr. Cayenna, Columbia (Coll. *Goudot*).

394. Coptocycla hieroglyphica.

Bohem. Mon. Cassid. iii. p. 503, 393.
Patr. Mexico, Veragua.

395. Coptocycla hebræa. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 504, 394.
Cassida hebræa, *Fabr.* Syst. El. i. p. 392, 24. Ent. Syst. i. p. 295, 14. Mant. Ins. i. p. 63, 13. Spec. Ins. i. p. 109, 10. *Herbst.* Naturs. viii. p. 273, 45, t. cxxxiii. f. 4, d. *Oliv.* Ent.

Ins. v. p. 381, 17. Entom. vi. p. 968, 77, 97, t. i. f. 8. *Schœnh.* Syn. Ins. ii. p. 216, 32.
Cassida reticularis, *De Geer*, Mem. v. p. 188, 14, t. xv. f. 17.
Patr. Cayenna, Brasilia, Venezuela (Coll. *Dyson*), Mexico.

396. Coptocycla plecta.

Bohem. Mon. Cassid. iii. p. 505, 395.
Cassida plecta, *Erichs.* Archiv. f. Naturg. xiii. p. 154, 7.
Patr. Peruvia.

397. Coptocycla retifera.

Bohem. Mon. Cassid. iii. p. 506, 396.
Cassida reticulata? *Thunb.* Nov. Ins. Spec. v. p. 87, t. v. f. 97. *Herbst.* Naturs. viii. p. 265, 36, t. cxxxii. f. 8?
Patr. Nova Granada.

398. Coptocycla reticulata.

Bohem. Mon. Cassid. iii. p. 507, 397.
Cassida reticulata, *Fabr.* Syst. El. i. p. 395, 38. *Schœnh.* Syn. Ins. ii. p. 218, 52.
Patr. America meridionalis.

399. Coptocycla cumulata.

Bohem. Mon. Cassid. iii. p. 508, 398.
Patr. Nova Valencia, Cabello.

400. Coptocycla confusa.

Bohem. Mon. Cassid. iii. p. 509, 399.
Patr. Brasilia.

i. *Elytra flavescentia, singulo in disco exteriore vitta obscura.*

401. Coptocycla sertata.

Bohem. Mon. Cassid. iii. p. 510, 400.
Cassida sertata, *Erichs.* Archiv. f. Naturg. xiii. p. 154, 6.
Patr. Peruvia, Columbia.

402. Coptocycla tabida.

Bohem. Mon. Cassid. iii. p. 511, 401.
Patr. Mexico.

403. Coptocycla quadrata. B.M.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 512, 402.
Cassida quadrata, *Fabr.* Syst. El. i. p. 395, 37. *De Geer*,

Mem. v. p. 188, 13, t. xv. f. 16. *Schœnh.* Syn. Ins. ii. p. 218, 49.
Patr. Brasilia, Bahia, Bolivia, Columbia, Cayenna, Surinam.

404. Coptocycla aprica.

Bohem. Mon. Cassid. iii. p. 514, 403.
Patr. Cayenna.

405. Coptocycla coronata.

Bohem. Mon. Cassid. iii. p. 515, 404.
Patr. Mexico.

406. Coptocycla plebeja.

Dej. Cat. ed. 3, p. 397. *Bohem.* Mon. Cassid. iii. p. 516, 405.
Patr. Mexico.

k. *Elytra flavescentia, annulo communi nigro ornata.*

407. Coptocycla vivida.

Bohem. Mon. Cassid. iii. p. 517, 406.
Patr. Mexico.

408. Coptocycla trepida.

Bohem. Mon. Cassid. iii. p. 518, 407.
Patr. Bahia, Columbia.

Cassidæ apud Auctores descriptæ, mihi ignotæ.

Cassida binotata.

Fabr. Syst. El. i. p. 398, 63. Ent. Syst. i. p. 299, 31. *Herbst.* Naturs. viii. p. 337, 101. *Schœnh.* Syn. Ins. ii. p. 221, 85. *Bohem.* Mon. Cassid. iii. p. 520.
Patr. Cayenna.

Cassida bituberculata.

Fabr. Syst. El. i. p. 399, 66. Ent. Syst. i. p. 299, 32. Mant. Ins. App. p. 379, 21, 22. *Herbst.* Naturs. viii. p. 338, 102. *Linn.* Syst. Nat. Gmel. i. iv. p. 1643, 32. *Schœnh.* Syn. Ins. ii. p. 221, 89. *Bohem.* Mon. Cassid. iii. p. 520.
Patr. Cayenna.

Cassida leucophæa.

Linn. Syst. Nat. i. ii. p. 576, 13. Ed. Gmel. i. iv. p. 1640, 13. Amœn. Acad. vi. p. 393, 10. Cent. Ins. p. 10, 10. *Fabr.*

Syst. El. i. p. 403, 92. Ent. Syst. i. p. 303, 52. Mant. Ins. i. p. 65, 43. Spec. Ins. i. p. 113, 36. Syst. Ent. p. 92, 22. *Herbst.* Naturs. viii. p. 346, 115. *Schœnh.* Syn. Ins. ii. p. 225, 119. *Bohem.* Mon. Cassid. iii. p. 520.
Patr. America meridionalis.

CASSIDA SUPPOSITA.

Lederm. Miscrop. t. xxviii. f. F. g. et t. xliv. f. b. c. d. *Linn.* Syst. Nat. i. ii. p. 578, 25. Ed. Gmel. i. iv. p. 1642, 25. *Schœnh.* Syn. Ins. ii. p. 229, 153. *Bohem.* Mon. Cassid. iii. p. 520.
Patr. America calidioris.

CASSIDA CRUENTATA.

Oliv. Entom. vi. p. 929, 7, 97, t. iv. f. 53. *Bohem.* Mon. Cassid. iii. p. 521.
Patr. — — —

CASSIDA QUATUORDECIMPUNCTATA.

Oliv. Entom. vi. p. 943, 32, 97, t. iv. f. 66. *Bohem.* Mon. Cassid. iii. p. 521.
Patr. America meridionalis.

CASSIDA MORBILLOSA.

Oliv. Entom. vi. p. 949, 43, 97, t. v. f. 72. *Bohem.* Mon. Cassid. iii. p. 521.
Patr. America meridionalis.

CASSIDA PALLIDA.

Oliv. Entom. vi. p. 952, 47, 97, t. iii. f. 43. *Bohem.* Mon. Cassid. iii. p. 521.
Patr. Guyana.

CASSIDA MICANS.

Oliv. Entom. vi. p. 960, 63, 97, t. v. f. 83. *Bohem.* Mon. Cassid. iii. p. 522.
Patr. Java.

CASSIDA CASSIDATA.

Herbst. Naturs. viii. p. 255, 14, t. cxxxi. f. 3. *Schœnh.* Syn. Ins. ii. p. 216, 24. *Bohem.* Mon. Cassid. iii. p. 522.
Patr. — — —

CASSIDA LIVIDA.

Herbst. Naturs. viii. p. 261, 30, t. cxxxii. f. 2. *Schœnh.* Syn. Ins. ii. p. 217, 47. *Bohem.* Mon. Cassid. iii. p. 522.
Patr. India.

CASSIDA PALLIDA.

Herbst. Naturs. viii. p. 262, 31, t. cxxxii. f. 3. *Schœnh.* Syn. Ins. ii. p. 218, 48. *Bohem.* Mon. Cassid. iii. p. 523.
Patr. Carolina.

CASSIDA DISPAR.

Herbst. Naturs. viii. p. 267, 39, t. cxxxii. f. 10. *Schœnh.* Syn. Ins. ii. p. 218, 57. *Bohem.* Mon. Cassid. iii. p. 523.
Patr. India orientalis.

CASSIDA MARYLANDICA.

Herbst. Naturs. viii. p. 274, 46, t. cxxxiii. f. 5. *Schœnh.* Syn. Ins. ii. p. 216, 33. *Bohem.* Mon. Cassid. iii. p. 524.
Patr. Maryland.

CASSIDA PUMILA.

Herbst. Naturs. viii. p. 276, 48, t. cxxxiii. f. 7. *Schœnh.* Syn. Ins. ii. p. 218, 58. *Bohem.* Mon. Cassid. iii. p. 524.
Patr. Sierra Leona.

CASSIDA NIGROMACULATA.

Herbst. Naturs. viii. p. 277, 49, t. cxxxiii. f. 8. *Schœnh.* Syn. Ins. ii. p. 218, 54. *Bohem.* Mon. Cassid. iii. p. 525.
Patr. Sierra Leona.

CASSIDA ATERRIMA.

Herbst. Naturs. viii. p. 294, 62, t. cxxxiv. f. 7. *Schœnh.* Syn. Ins. ii. p. 230, 163. *Bohem.* Mon. Cassid. iii. p. 525.
Patr. America borealis.

CASSIDA TESTACEA.

Herbst. Naturs. viii. p. 300, 67, t. cxxxiv. f. 11. *Schœnh.* Syn. Ins. ii. p. 230, 165. *Bohem.* Mon. Cassid. iii. p. 526.
Patr. India.

CASSIDA TRICOLOR.

Herbst. Naturs. viii. p. 304, 70, t. cxxxv. f. 2. *Schœnh.* Syn. Ins. ii. p. 228, 145. *Bohem.* Mon. Cassid. iii. p. 526.
Patr. India.

CASSIDA LIRIOPHORA. B.M.

Kirby, Act. Soc. Linn. iii. p. 8, 1. *Marsh.* Entom. Britann. i. p. 145, 5. *Schœnh.* Syn. Ins. ii. p. 212, 6. *Bohem.* Mon. Cassid. iii. p. 527. (C. vibex, *L.* sec. *Stephens*, Syst. Cat. I. p. 229.
Patr. Britannia in Serratula arvensi.

CASSIDA PALLIDA.

Thunb. Ins. Suec. viii. p. 103. *Schœnh.* Syn. Ins. ii. p. 220, 74. *Bohem.* Mon. Cassid. iii. p. 528.
Patr. Upsalia.

CASSIDA GUERINI.

Boisd. Faun, Entom. de l'Océan. ii. p. 538, 4. *Bohem.* Mon. Cassid. iii. p. 528.
Patr. Nova Guinea.

CASSIDA NIGROPUNCTATA.

Klug, Erman. Reise Ins. p. 47, 175. *Bohem.* Mon. Cassid. iii. p. 528.
Patr. Africa occidentalis.

CASSIDA TESTUDINARIA.

Montrouzier. Ann. Sc. Phys. Lyon. 1855, vii. 67.
Patr. Woodlark I.

CASSIDA AURATA.

Montrouzier. Ann. Sc. Phys. Lyon. 1855, vii. 67.
Patr. Woodlark I.

CASSIDA STRIGULA.

Montrouzier. Ann. Sc. Phys. Lyon. 1855, vii. 67.
Patr. Woodlark I.

INDEX.

abrupta (Char.), 149.
Abyssinica (Cass.), 117.
aciculata (Copt.), 198.
Acromis, 85.
aculeata (Bat.), 95.
—— (Cass.), 19.
acuminata (Mes.), 48.
acuta (Mes.), 50.
acuticornis (Deloy.), 189.
adamantina (Copt.), 161.
adhærens (Asp.), 108.
admirabilis (Char.), 151.
adnata (Chel.), 83.
adspersa (Cass.), 121.
—— (Mes.), 61.
advena (Chel.), 80.
ægrota (Pœcil.), 67.
æmula (Cass.), 138.
ænea (Cyrt.), 50.
—— (Dol.), 28.
—— (Mes.), 47.
æneo-maculata (Chel.), 74.
æquinoctialis (Copt.), 169.
ærata (Psal.), 156.
ærea (Dem.), 23.
æruginosa (Taur.), 20.
Æthiopica (Cass.), 125.
affinis (Cass.), 139.
Afzelii (Aspid.), 109.
albicolle (Im.), 13.
albicollis (Cass.), 13.
albida (Mes.), 64.
albiventris (Om.), 90.
albo-fasciculata (Mes.), 39.
alternans (Chel.), 76.
alutacea (Phys.), 98.
amabilis (Del.), 115.
ambigena (Copt.), 174.
ambigua (Char.), 149.
amicta (Copt.), 197.
Ammon (Taur.), 20.
amæna (Copt.), 168.
amænula (Copt.) 169.
amplexa (Copt.), 186.
ampliata (Mes.), 39.
amplissima (Asp.), 106.
Ampulla (El.), 72.
anceps (Mes.), 43.
anchoralis (Taur.), 21.
Ancilla (Copt.), 166.
ancorifera (Cass.), 130.
Andreæ (Copt.), 188.
angulata (Pœcil.), 69.
angulifera (Cass.), 118.
angustata (C.), 86.
annexa (Copt.), 179.
annularis (Copt.), 170.
annulata (Cass.), 56.
annulifera (Copt.), 170.
annulum (Im.), 12.
annulus (Cass.), 165.
—— (Copt.), 173.
anthracina (Pœcil.), 71.
antiqua (Mes.), 50.
—— (Taur.), 22.
anxia (Copt.), 199.
apiata (Chel.), 76.
apicalis (Asp.), 106.

apiculata (Bat.), 95.
aprica (Copt.), 203.
arcuata (Cass.), 159.
—— (Copt.), 172.
areata (Cass.), 124.
—— (Chel.), 84.
areolata (Mes.), 43.
Argus (Chel.), 78.
armata (Dor.), 94.
armigera (Hopl.), 3.
asperata (Copt.), 190.
ASPIDOMORPHA, 104.
assimilis (Aspid.), 114.
—— (Mes.), 44.
ASTERIZA, 147.
Astrolabiana (Aspid.), 113.
aterrima (Cass.), 205.
atomaria (Chel.), 80.
—— (Phys.), 100.
atra (C.), 93.
atramentosa (Char.), 150.
atrata (Cass.), 142.
atriceps (Psal.), 156.
atrofasciata (Mes.), 60.
atroguttata (Mes.), 60.
atro-rubra (Mes.), 60.
attenuata (Phys.), 102.
Aubei (Copt.), 169.
aucta (Copt.), 168.
Augur (Mes.), 61.
augusta (Om.), 92.
aulica (Om.), 86.
aurata (Cass.), 206.
aurea (Cass.), 139.
aureicornis (Taur.), 20.
aureomarginata (Mes.), 58.
aurichalcea (Copt.), 163.
auricoma (Copt.), 163.
aurifera (Copt.), 165.
aurisplendens (Copt.), 185.
aurofasciata (Copt.), 171.
auroguttata (Char.), 152.
aurolimbata (Desm.), 24.
aurulenta (Char.), 148.
Australasiæ (Aspid.), 113.
Australica (Copt.), 175.
Austriaca (Cass.), 122.
axillaris (Om.), 89.
azurea (Cass.), 132.
azureicornis (Taur.), 20.
Bacca (El.), 72.
badia (Hopl.), 7.
Bahiana (Copt.), 181.
bajula (Aspid.), 111.
Ballista (Bat.), 94.
Balyi (Copt.), 191.
basalis (C.), 89.
—— (Ech.), 87.
basilica (Om.), 92.
BASIPRIONOTA, 8.
BASIPTA, 97.
Batesi (Dol.), 32.
BATONOTA, 93.
bella (Cass.), 131.
bellicosa (Bat.), 94.
bellula (Char.), 151.
Berolinensis (Cass.), 141.
Beskei (Porph.), 17.
biaculeata (Hopl.), 3.
biannularis (Chel.), 82.
biarcuata (Copt.), 176.
bicincta (Char.), 152.
bicingulata (Copt.), 170.
bicolon (Copt.), 159.
bicolor (Call.), 15.
—— (Cass.), 163.
bicolorata (Mes.), 63.
bicornis (Taur.), 19.
bicornuta (Desm.), 25.
bicostata (Phys.), 99.
bicruciata (Pœcil.), 67.
bicurvata (Copt.), 180.
bidens (Bat.), 93.
bifasciata (Copt.), 163.
bifenestra (Copt.), 158.
bifenestrata (Copt.), 199.
—— (Mes.), 53.
bifossulata (Copt.), 162.
biguttata (Asp.), 108.
bihamata (Dol.), 31.
biimpressa (Copt.), 162.
bilineata (Mes.), 62.
bilobata (Aspid.), 111.
—— (Copt.), 200.
bimacula (Cass.), 9.
bimaculata (Asp.), 108.

bimaculata (Cass.), 49.
—— (Pri.), 9.
bimarginata (Char.), 152.
binotata (Cass.), 203.
—— (Dol.), 32.
—— (Pri.), 10.
bioculata (Mes.), 54.
bipartita (Copt.), 172.
bipellucida (Copt.), 199.
biplagiata (Dol.), 31.
biplagosa (Om.), 91.
bipunctata (Cass.), 160.
bipuncticollis (Cass.), 118.
—— (Pri.), 9.
bipunctipennis (Copt.), 160.
bipustulata (C.), 35.
—— (Mes.), 53.
biradiata (Aspid.), 112.
biramosa (Copt.), 191.
biremis (Aspid.), 115.
bisignata (Copt.), 160.
bisinuata (Copt.), 188.
bis-bimaculata (Copt.), 187.
bis-binotata (Copt.), 163.
bis-biplagiata (Dol.), 31.
bis-bipustulata (Copt.), 159.
bispinosa (Desm.), 24.
—— (Hopl.), 3.
bistigma (Mes.), 52.
bis-triguttata (Copt.), 187.
bis-trilineata (Cass.), 137.
bis-trimaculata (Copt.), 159.
bis-trinotata (Copt.), 159.
bis-tripunctata (Copt.), 163.
bis-tripustulata (Cass.), 162.
bituberculata (Cass.), 203.
—— (Mes.), 37.
bivittipennis (Bat.) 96.
bivulnerata (Copt.), 163.
Blanchardi (Copt.), 158.
blanda (Cyrt.), 43.
—— (Mes.), 48.
blandidica (Copt.), 197.
Boisduvali (Aspid.), 110.
Boliviana (Chel.), 82.
—— (Taur.), 20.
Bonariensis (Pœcil.), 67.
Bonfilsii (Om.), 91.
Botanochara, 65.
Bowringii (Copt.), 161.
brachiata (Mes.), 63.
Bremei (Copt.), 174.
brevicollis (Mes.), 59.
brevicornis (Cal.), 17.
brevilineata (Eur.), 103.
breviuscula (Phys.), 99.
brunnea (Chel.), 73.
brunneosignata (Om.), 89.
brunnicollis (Cyrt.), 63.
bullata (Chel.), 74.
Buqueti (Copt.), 187.

cædemadens (C.), 36.
calamitosa (Can.), 27.
Calaspidea, 34.
calcarata (Om.), 90.
—— (Phys.), 102.
Calliaspis, 14.
calligera (Aspid.), 112.
callosa (Cass.), 142.
Calopepla, 2.
calva (Chel.), 76.
—— (Copt.), 183.
Calyptocephala, 17.
canaliculata (Cass.), 122.
cancellata (Lac.), 154.
—— (Pœcil.), 69.
candens (Psalid.), 157.
candida (Phys.), 101.
Canistra, 26.
Capensis (Cass.), 11.
carbonaria (Can.), 27.
Cardui (Cass.), 144, var. 126.
carnifex (Copt.), 201.
carnulenta (Copt.), 190.
cariosa (Desm.), 23.
Cassida, 116.
cassidata (Cass.), 204.
cassidoides (Call.), 14.
castigata (Mes.), 45.
catenata (Copt.), 175.
catenulata (Chel.), 77.
Catharinensis (Chel.), 75.
caudata (Mes.), 46.
—— (Phys.), 101.
cavata (Chel.), 73.

Cayennensis (Copt.), 177.
centralis (Copt.), 186.
centromaculata (Char.), 151.
centropunctata (Copt.), 195.
Cephaloleia, 14.
Cerastes (C.), 86.
cerea (Phys.), 103.
Ceylonica (Copt.), 176.
chalybea (Mes.), 42.
chalybeum (Him.), 13.
Charidotis, 148.
chalcoptera (Mes.), 45.
Chelymorpha, 73.
Chevrolati (Mes.), 49.
Chinensis (Lac.), 155.
chlorina (Asp.), 106.
Chloris (Cass.), 127.
—— (Dol.), 34.
chlorosoma (Dol.), 31.
chlorotica (Asp.), 104.
cicatricosa (Lac.), 154.
cincta (Asp.), 105.
—— (C.), 73.
cinctella (Char.), 150.
cincticula (Copt.), 170.
cinctipennis (Chel.), 85.
cinctum (Him.), 13.
cinctus (Sphær.), 1.
cinerea (C.), 87.
cingulata (Chel.), 78.
cinnabarina (Call.), 14.
cinnabaris (Call.), 14.
circinaria (Copt.), 196.
circinata (Copt.), 166.
circularis (Cass.), 173.
—— (Copt.), 196.
circulata (Copt.), 170.
circulus (Copt.), 166.
circumcincta (Copt.), 196.
circumdata (Copt.), 177.
—— (Porph.), 16.
circumducta (Copt.), 165.
circumflexa (Copt.), 169.
circumpuncta (Chel.), 82.
circumspicua (Psalid.), 157.
citrina (Phys.), 103.
citrinella (Phys.), 101.
clandestina (Copt.), 187.
clathrata (Cass.), 116.
—— (Om.), 91.
clavata (Copt.), 189.
clitelligera (Copt.), 171.
clivosa (Chel.), 74.
clypeata (C.), 90.
—— (Dol.), 28.
clypeolata (Char.), 150.
coagulata (Cass.), 142.
cobaltina (Chel.), 75.
coccinea (Char.) 149.
—— (Porph.), 16.
coccinelloides (Copt.) 171.
cœca (Chel.), 85.
cœruleata (Dol.), 30.
cœruleomaculata (Meg.), 8.
cœruleonotata (Mes.), 64.
cœrulescens (Mes.), 58.
cognata (Char.), 150.
collaris (Cass.), 13.
collata (Om.), 88.
Colliculus (Dol.), 29.
collina (Cass.), 118.
Columbiana (Cal.), 35.
comata (Chel.), 73.
combusta (Cass.), 145.
commutabilis (Chel.), 80.
complanata (Copt.), 157.
comptum (Im.), 11.
compuncta (Copt.), 178.
concentrica (Copt.), 167, 170.
concinna (Bot.), 67.
—— (Cass.), 145.
conclusa (Copt.), 171.
concolor (Copt.), 183.
confinis (Asp.), 106.
conflagrata (Copt.), 179.
conformis (Cass.), 128.
confusa (Copt.), 202.
—— (Mes.), 47.
congener (Mes.), 38.
congregata (Copt.), 174.
conjugata (Chel.), 78.
conjuncta (Pœcil.), 68.
connectens (Copt.), 180.
connexa (Copt.), 165.
consanguinea (Del.), 105.
conscripta (Copt.), 201.

consentanea (Copt.), 166.
consimilis (Cass.), 144.
—— (Copt.), 170.
consobrina (Copt.), 167.
—— (Mes.), 43.
conspersa (Mes.), 42.
conspicillata (Copt.), 177.
conspurcata (Cass.), 131.
constellata (Chel.), 74.
—— (Copt.), 193.
constricta (Copt.), 174.
contaminata (Can.), 27.
contempta (Psal.), 156.
contexta (Copt.), 173.
contubernalis (Mes.), 48.
convexa (Cass.), 134.
convexula (Cass.), 134.
Coptocycla, 157.
coracina (Taur.), 21.
corallina (Hemi.), 15.
Cordula (Cass.), 145.
cornuta (Taur.), 21.
cornigera (Bat.), 94.
coronata (Copt.), 203.
coronula (Cass.), 124.
corrosa (Cass.), 133.
corrugata (Lac.), 154.
corticina (Pœcil.), 67.
Cossyphoides (Del.), 1.
costata (Cass.), 140.
costipennis (Porph.), 16.
Craspedonta, 2.
creberrima (Taur.), 19.
crenata (Copt.), 185.
crenulata (Desm.), 23.
cribraria (C.), 74.
—— (C.), 78.
—— (Chel.), 80.
cribripennis (Desm.), 25.
cribrosa (Copt.), 191.
cribrum (Mes.), 61.
crispata (Copt.), 171.
cruciata (Cass.), 181.
—— (Copt.), 190.
crucifera (Chel.), 84.
crucipennis (Copt.), 180.
cruenta (Cass.), 133.
cruentata (Cass.), 128.
cruentata (Cass.), 204.
—— (Mes.), 46.
Crux (Cass.), 190.
—— (Deloy.), 188.
Cumingii (Phys.), 99.
cumulata (Copt.), 202.
cuprea (Cass.), 53.
cyanea (C.), 35.
—— (Mes.), 58.
—— (Porph.), 16.
cyanescens (Cyrt.), 58.
cyaneum (Him.), 16.
cyanipenne (Him.), 13.
cyclographa (Copt.), 167.
Cyclops (Copt.) 166.
Cynaræ (Copt.), 194.
Cynthia (Hyb.), 148.
Cyphomorpha, 73.
cyrtodes (Phys.), 98.
Cyrtonota, 22, 73.

Dahlbomi (Char.), 149.
Daurica (Cass.), 141.
decem-guttata (Mes.), 57.
decem-maculata (Pri.), 11.
decemnotata (Copt.), 161.
—— (Pœcil.), 71.
—— (Pri.), 10.
decempunctata (Copt.), 161.
decempustulata (Pœcil.), 70.
—— (Pri.), 9.
decemsignata (Pri.), 11.
decemstillata (Chel.), 82.
—— (Pri.). 9.
decemverrucata (Pœcil.), 70.
decipiens (Om.), 88.
decolorata (Cass.), 144.
decora (Desm.), 24.
decorata (Cyrt.), 42.
decorum (Him.), 24.
decussata (C.), 35.
—— (Copt.), 180.
—— (Mes.), 65.
deflorata (Cass.), 127.
Dejeanii (Bat.), 96.
delecta (Mes.), 62.
Delesserti (Cass.), 132.
deleta (Mes.), 62.

Delocrania, 1.
Deloyala, 97, 104.
demissa (Mes.), 47.
denticollis (Cass.), 125.
denticulata (Cass.), 137.
—— (Desm.), 23.
dentipennis (Desm.), 24.
denudata (Mes.), 39.
depilata (Dol.), 30.
deplanata (Pœcil.), 68.
deplorabunda (Mes.), 41.
depressa (Cass.), 127.
derosa (Copt.), 199.
deruta (Plat.), 104.
desertorum (Cass.), 147.
designata (Copt.), 186.
Desmaresti (Cal.), 36.
Desmonota, 22.
deusta (Cass.), 118.
Deyrollei (Copt.), 186.
Diana (Copt.), 187.
diaphana (Aspid.), 113.
dichroa (Om.), 87.
didyma (C.), 65.
difficilis (Copt.), 198.
diffinis (Copt.), 196.
difformis (Mes.), 37.
dilatata (Cass.), 193.
—— (Mes.), 63.
dilecta (Aspid.), 114.
dimidiata (Mes.) 53.
Diomma (Copt.), 170.
discicollis (Char.), 152.
discipennis (Isch.), 115.
discoidalis (Copt.), 167.
discoides (Mes.), 54.
discoideum (Him.), 13.
discolor (Copt.), 192.
Discomorpha, 26.
disconotata (Chel.), 79.
discophora (Call.), 15.
discors (Cal.), 35.
—— (Cass.), 53.
discreta (Lac.), 156.
disjuncta (Phys), 100.
dispar (Cass.), 205.
dissecta (Mes.), 56.
dissimilis (Copt.), 200.
distigma (Dol.), 33.
distincta (Om.), 91.
distinguenda (Chel.), 74.
distorta (Copt.), 182.
diversa (Mes.), 41.
divisa (Aspid.), 110.
divulsa (Copt.), 200.
Dodonea (Copt.), 174.
Dohrni (Copt.), 182.
Dolichotoma, 28.
dolorosa (Copt.), 200.
dolosa (Copt.), 161.
dorsalis (Cass.), 127.
—— (Copt.), 182.
dorsata (Aspid.), 112.
—— (Cass.), 114.
dorsonotata (Cass.), 132.
dorsopunctata (Psalid.), 157.
dorsosignata (Copt.), 188.
dorsovittata (Cass.), 129.
Dorynota, 93.
Douei (Aspid.), 110.
Dregei (Cass.), 124.
Drewseni (Copt.), 179.
Dromedarius (Dol.), 29.
dubia (Phys.), 103.
ducalis (Dol.), 33.
—— (Pœcil.), 67.
duodecimmaculata (Pœcil.), 69.
duodecimnotata (Pœcil.), 71.
duodecimpustulata (Cass.), 53.
duodecimsignata (Chel.), 78.
duodecimverrucata (Pœcil.), 71.
duplicata (Pœcil.), 72.
Duponti (Desm.), 24.

echinata (Hopl.), 3.
Echoma, 86, 97.
effusa (Copt.), 181.
egena (Aspid.), 115.
egregia (Copt.), 190.
elatior (Copt.), 193.
elegans (Copt.), 176.
—— (Mes.), 49.
elevata (Aspid.), 111.
elocata (Mes.), 56.
Elytrogona, 72.
emarcida (Copt.), 184.

emarginata (Copt.), 179.
encausta (Pœcil.), 71.
ensifera (Bat.), 95.
ephippiata (Copt.), 170.
ephippium (Mes.), 55.
Epistictia, 2.
equestris (Cass.), 143.
Erichsoni (Pœcil.), 68.
erratica (Copt.), 169.
erythrocata (Hopl.), 6.
erythrocera (Cass.), 16.
Eugenysa, 34.
Eurypepla, 103.
evanida (Mes.), 65.
exaltata (Bat.), 96.
exarata (Phys.), 101.
excavata (Lac.), 154.
excelsa (Copt.), 176.
excisa (Mes.), 55.
Exclamationis (Mes.), 64.
exigua (Char.), 152.
exilis (Cass.), 132.
eximia (Meg.), 8.
—— (Taur.), 20.
exsanguis (Copt.), 184.
exsculpta (Cass.), 140.
extensa (Copt.), 175.
extricata (Mes.), 59.

Fahrœi (Copt.), 172.
Fairmairei (Copt.), 196.
falcifera (Mes.), 64.
fallax (Copt.), 193.
farinosa (Cass.), 146.
fasciata (Cass.), 11.
—— (Chel.), 77.
fasciatum (Him.), 11.
fasciculata (Mes.), 40.
fastidiosa (Copt.), 162.
fastidita (Pœcil.), 71.
fastuosa (Cass.), 123.
—— (Mes.), 48.
fenestralis (Char.), 151.
fenestrata (Asp.), 108.
ferrugata (Disc.), 27.
ferruginea (Cass.), 52.
—— (Cass.), 139.
festiva (Mes.), 58.
finitima (Pœcil.), 68.
flava (Cass.), 110.
—— (Om.), 87.
flavago (Phys.), 100.
flavata (Om.), 89.
flaveola (Cass.), 140.
—— (Phys.), 102.
flavescens (Copt.), 188.
flavicans (Char.), 153.
flavicollis (Chel.), 75.
flavicornis (Ast.), 147.
flavipennis (Mes.), 62.
flavo-cincta (Copt.), 196.
flavo-guttata (Mes.), 50.
flavo-lineata (Copt.), 178.
flavo-maculata (Mes.), 55.
flavo-notata (Copt.), 197.
—— (Mes.), 50.
flavo-reticulata (Mes.), 45.
flavo-rugosa (Copt.), 179.
flavo-signata (Mes.), 60.
flavo-sparsa (Mes.), 42.
flavo-vittata (Copt.), 178.
flexuosa (Copt.), 191.
floccosa (Mes.), 46.
foliata (Copt.), 199.
forcipata (Copt.), 169.
formosa (Copt.), 172.
fornicata (C.), 72.
—— (Hyb.), 148.
fossulata (Cass.), 142.
—— (Mes.), 41.
foveolata (Cass.), 116.
fraterna (Copt.), 198.
frenata (Asp.), 108.
frontalis (Cass.), 143.
fulgida (Copt.), 185.
fuliginosa (Deloy.), 190.
fulvum (Him.), 14.
fumosa (Copt.), 164.
funebris (Mes.), 54.
funeraria (Call.), 15.
funerea (Aspid.), 109.
funesta (Bat.), 95.
furcata (Cass.), 114.
furunculus (Copt.), 171.
furva (Char.), 151.
fusca (Cass.), 140.

fuscata (Phys.), 100.
fusco-notata (Aspid.), 109.
fusco-punctata (Aspid.), 112.
fusco-signata (Cass.), 130.
fusco-sparsa (Cass.), 143.
fusco-vittata (Copt.), 177.

Galamensis (Asp.), 108.
galbanata (Hyb.), 148.
Gallica (Cass.), 127.
gemellata (Char.), 151.
gemina (Copt.), 196.
gemmata (Hopl.), 7.
generosa (Om.), 87.
geniculata (Chel.), 78.
gentilis (Mes.), 59.
gestatrix (Char.), 152.
gibba (Chel.), 74.
gibbera (Hyb.), 148.
—— (Taur.), 22.
gibberosa (Copt.), 185.
gibbipennis (Cass.), 146.
gibbosa (Mes.), 38.
gibbula (Cass.), 135.
gigantea (Phys.), 99.
glabella (Cass.), 135.
glabrata (Copt.), 198.
glabricollis (Mes.), 40.
glabricula (Copt.), 184.
Gladiator (Bat.), 94.
glareosa (Copt.), 194.
glaucina (Copt.), 183.
glauco-vittata (Copt.), 164.
globosa (Copt.), 186.
glomerosa (Char.), 153.
glyphica (Copt.), 174.
Godetii (Mes.), 41.
Goryi (Mes.), 51.
Goudoti (Copt.), 175.
graminea (Cass.), 129.
granaria (Copt.), 181.
grandis (Eng.), 35.
granula (Cass.), 131.
graphica (Cass.), 138.
graphiptera (Chel.), 84.
gravida (Desm.), 24.
Grayi (Dol.), 32.
gregaria (Bat.), 95.
grossa (Cal.), 34.
Guadeloupensis (Copt.), 194.
Guerini (Cass.), 206.
Guianiensis (Cal.), 36.
guttata (Copt.), 181.
guttifera (Chel.), 84.
guttigera (Copt.), 197.
guttula (Chel.), 85.
gyrata (Copt.), 169.

Hablitziæ (Cass.), 146.
hæmatidea (Om.), 89.
hæmatina (Mes.), 59.
hæmatites (C.), 63.
hæmatodes (Pœcil.), 67.
hæmatura (Chel.), 81.
hamata (Copt.), 190.
hamulata (Copt.), 181.
hebes (Cass.), 120.
hebræa (Copt.), 201.
hectica (Copt.), 198.
Helianthi (Phys.), 98.
Hemisphærota, 15.
hemisphærica (Cass.), 145.
hemispila (Om.), 89.
hepatica (Cass.), 120.
herbacea (Cass.), 137.
herbea (Cass.), 127.
—— (Copt.), 185.
herbida (Char.), 153.
Heroina (Aspid.), 110.
heteropuncta (Cass.), 141.
hexastigma (Cass.), 127.
hieroglyphica (Cass.), 57.
—— (Copt.), 201.
hilaris (Copt.), 192.
Himatidium, 11.
hispoides (Del.), 1.
Histrio (Pœcil.), 72.
Hoepfneri (Chel.), 73.
Hogbergi (Copt.), 165.
honesta (Chel.), 81.
Hoplionota, 2.
horrida (Hopl.), 4.
horrifica (Hopl.), 5.
Horsfieldi (Copt.), 191.
hospita (Lac.), 155.
Hottentotta (Cass.), 136.

humeralis (Bat.), 97.
humilis (Phys.), 102.
hyalina (Om.), 87.
Hybosa, 148.
hybrida (Cass.), 119.
hypocrita (Copt.), 199.

icterica (Aspid.), 113.
—— (Cass.), 130.
ignava (Taur.), 21.
ignita (Mes.), 44.
ignobilis (Chel.), 75.
illepida (Mes.), 44.
illota (Cass.), 120.
illudens (Bat.), 97.
illustris (Cyrt.), 48.
—— (Mes.), 57.
Illyrica (Cass.), 122.
Imatidium, 11.
immaculata (Copt.), 184.
immerita (Copt.), 185.
immunda (Copt.), 182.
immunita (Copt.), 182.
immutabilis (Cass.), 122.
imperialis (Cal.), 35.
—— (Chel.), 82.
impicta (Aspid.), 113.
implexa (Mes.), 44.
impluviata (Mes.), 43.
impressa (Lac.), 155.
impressipennis (Pœcil.), 65.
impudens (Mes.), 40.
impura (Copt.), 197.
impurum (Him.), 12.
impustulata (Cyrt.), 50.
—— (Pri.), 8.
inæqualis (Mes.), 49.
inanis (Isch.), 115.
incarnata (Mes.), 63.
incerta (Copt.), 168.
incita (Copt.), 187.
inclusa (Chel.), 83.
incomta (Hopl.), 7.
inconstans (Copt.), 164.
incorrupta (Copt.), 184.
inculta (Copt.), 183.
incurva (Copt.), 179.
indecens (Hyb.), 148.
indecorata (Copt.), 197.
Indica (Aspid.), 115.
indigacea (Mes.), 58.
indigesta (Chel.), 80.
indistincta (Asp.), 105.
inedita (Copt.), 186.
inepta (Dol.), 28.
infaceta (Chel.), 74.
—— (Chel.), 76.
infima (Chel.), 85.
inflata (Chel.), 74.
ingenua (Copt.), 197.
ingrata (Phys.), 101.
innocens (Om.), 88.
innocua (Asp.), 105.
innotata (Cass.), 131.
inornata (Mes.), 43.
inquinata (Aspid.), 114.
—— (Cass.), 147.
inscripta (Copt.), 187.
insculpta (Copt.), 184.
insidiosa (Bat.), 96.
insignis (Chel.), 81.
insubida (Copt.), 180.
insularis (Cass.), 136.
interjecta (El.), 72.
intermedia (Copt.), 201.
—— (Desm.), 25.
interrupta (Cass.), 119.
intricata (Lac.), 154.
—— (Pœcil.), 67.
—— (Porph.), 16.
intumida (Copt.), 192.
inuncta (Aspid.), 113.
invasa (Pœcil.), 71.
involuta (Copt.), 168.
Iris (Copt.), 192.
irregularis (Cass.), 129.
irrorata (Can.), 26.
—— (Om.), 91.
Ischyrosonyx, 115.
Isparetta (Asp.), 105.

Jamaicensis (Dol.), 111.
—— (Eur.), 103.
Javanica (Copt.), 161.
Jekeli (Mes.), 52.
Jousselini (Copt.), 172.

jucunda (Mes.), 57.
Judaica (Copt.) 179.
jugata (Om.), 88.
—— (Taur.), 21.
Justini (Copt.), 157.
Juvenca (Chel.), 84.

Kiesenwetteri (Bat.), 96.
Klugii (Chel.), 83.
Kollari (Mes.), 59.
Kunzei (Bat.), 96.

labilis (Cass.), 131.
Laccoptera, 153.
Lacordairei (Mes.), 61.
lacrymosa (Cass.), 129.
lacunosa (Mes.), 40.
læta (Copt.), 163.
lævis (Cass.), 135.
languida (Cass.), 127.
—— (Phys.), 100.
lanuginosa (Dol.), 30.
lata (Cass.), 126.
—— (Mes.), 57.
Lataniæ (Porph.), 16.
lateralis (Cass.), 49.
—— (Cyrt.), 52.
—— (Mes.), 51.
lateraria (Copt.), 180.
latevittata (Copt.), 164.
laticollis (Desm.), 25.
—— (Mes.), 37.
latipennis (Mes.), 38.
Leayana (Cal.), 2.
Leayanum (Imat.), 2.
Lebasii (Mes.), 50.
lenis (Mes.), 41.
leopardina (Copt), 175.
Leprieuri (Copt.), 171.
leprosa (Psalid.), 157.
Lerouxii (Bat.), 97.
leucophæa (Cass.), 203.
limbata (Aspid.), 112.
—— (Plat.), 1.
limbipennis (Aspid.), 110.
limbiventris (Cass.), 144.
limpida (Copt.), 183.
lineata (Mes.), 62.
lineatocollis (Mes.), 59.
lineigera (Om.), 89.
lineola (Cass.), 138.
—— (Him.), 13.
liquida (Copt.), 165.
liriophora (Cass.), 206.
litigiosa (Cass.), 145.
liturata (Copt.), 173.
livida (Cass.), 205.
—— (Copt.), 164.
lobata (Aspid.), 111.
—— (Copt.), 189.
longicornis (C.), 33.
longula (Chel.), 79.
Lucasi (Copt.), 158.
lucida (Cass.), 133.
—— (Mes.), 43.
lucidula (Copt.), 182.
luctifera (Copt.), 191.
luctuosa (C.), 33.
—— (Hopl.), 6.
ludicra (Copt.), 187.
lugens (Dol.), 28.
lugubrina (Copt.), 181.
lugubris (Copt.), 199.
lunata (Cass.), 49.
lunifera (Copt.), 173.
lunulata (Om.), 92.
lurida (Cass.), 124.
lutarella (Phys.), 98.
lutea (Cal.), 17.
luteicollis (Cass.), 121.
luteipennis (Mes.), 62.
luteocincta (Bas.), 97.
luteola (Char.), 149.
lutescens (Cass.), 128.
lyncea (Cass.), 120.

Macleayi (Cass.), 117.
macularia (Pœcil.), 66.
maculata (Cass.), 122.
maculatissima (Cass.), 117.
—— (Mes.), 63.
maculicollis (Om.), 90.
maculipennis (Cass.), 121.
—— (Hopl.), 5.
—— (Pri.), 8.
maculipes (Phys.), 101.

maculiventris (Phys.), 102.
maculosa (Copt.), 198.
Madagascarica (Aspid.), 109.
Maerkeli (Pri.), 10.
magica (Copt.), 105.
malachitica (Taur.), 22.
mamillata (Copt.), 174.
Mannerheimi (Mes.), 49.
mansueta (Copt.), 166.
marculenta (Copt.), 183.
margaritacea (Cass.), 136, 145.
marginalis (Char.), 150.
marginata (C.), 33, 73.
—— (Hopl.), 6.
—— (Om.), 90.
margineguttata (Cyrt.), 63.
marginella (Char.), 153.
marginicolle (Him.), 14.
—— (Im.), 13.
marginicollis (Isch.), 116.
marginipennis (Cal.), 18.
marginiventris (Desm.), 25.
Marylandica (Cass.), 205.
matronula (Ep.), 2.
maura (Copt.), 158.
media (Copt.), 196.
MEGAPYGA, 7.
melanophthalma (Cass.), 145.
melanota (Copt.), 196.
Mellyi (Mes.), 56.
mendax (Copt.), 164.
mera (Cass.), 147.
meridionalis (Cass.), 140.
MESOMPHALIA, 36.
metallica (Cass.), 22.
—— (Desm.), 25.
—— (Dol.), 30.
meticulosa (Copt.), 185.
Mexicana (Phys.), 100.
micans (Aspid.), 114.
—— (Cass.), 204.
microcera (Copt.), 176.
miliaris (Asp.), 106.
—— (Cass.), 155.
militaris (Dor.), 93.
miniata (Char.), 148.
—— (Dol.), 29.
—— (Porph.), 17.
minuta (Cass.), 136.
mirabilis (Dol.), 32.
mirifica (Dol.), 30.
misella (Mes.), 64.
mitis (Dol.), 28.
modesta (Copt.), 198.
modica (Mes), 50.
mœsta (Copt.), 187.
mollis (Cass.), 136.
Molucana (Hopl.), 6.
Mongolica (Cass.), 138.
monoceros (Bat.), 93.
—— (Dor.), 94.
monstrosa (Desm.), 23.
Moori (Cass.), 124.
morbillosa (Cass.), 204.
Morio (Copt.), 201.
Mouffleti (Cass.), 120.
mucronata (Bat.), 94.
Mulsanti (Porph.), 17.
multicava (Desm.), 24.
multiguttata (Chel.), 83.
multinotata (Lac.), 155.
multipunctata (C.), 80.
multisignata (Copt.), 188.
munda (Copt.), 180.
murræa (Cass.), 122.
mutabilis (Cass.), 125, 136.
mutata (Aspid.), 114.
mutilata (Aspid.), 115.
myops (Copt.), 167.
mysticæ (Copt.), 195.

nana (Cass.), 135.
Natalensis (Aspid.), 113.
navicula (Cass.), 135.
nebulosa (Cass.), 139, 140.
—— (Sel.), 86.
neglecta (Copt.), 165.
Nepalensis (Lac.), 156.
nervosa (Pœcil.), 66.
nigra (Bat.), 93.
—— (Cass.), 126.
nigriceps (Cass.), 144.
nigricollis (Chel.), 83.
nigricornis (Cal.), 17.
nigripes (Cass.), 124.
nigritarsis (Dol.), 34.

nigritarsis (Phys.), 101.
nigriventris (Cass.), 132.
nigro-ænea (Cass.), 27.
nigro-annulata (Copt.). 195.
nigro-cincta (Char.), 152.
nigro-dorsata (Cass.), 119.
nigro-lineata (Om.), 88.
nigro-maculata (Cass.), 205.
—— (Copt.), 164.
nigro-notata (Cass.), 121.
nigro-picta (Pœcil.), 68.
nigro-punctata (C.), 107, 206.
—— (Dol.), 29.
nigro-signata (Isch.), 116.
nigro-sparsa (Cass.), 121.
nigro-vittata (Cass.), 120.
Nilotica (Cass.), 121.
nimbatum (Him.), 12.
nitida (Copt.), 188.
nitidicollis (Phys.), 103.
nitidula (Mes.), 65.
nobilis (Cass.), 134.
nobilitata (Copt.), 195.
nodipennis (Cass.), 50.
nodosa (Bat.), 94.
—— (Poly.), 24.
nodulosa (Cass.), 116.
normalis (Om.), 86.
Notosacantha, 3.
Novæguineensis (Aspid.), 113.
novemdecimnotata (Lac.), 155.
novemmaculata (Cass.), 124.
nubilosa (Copt.), 158.
nucleata (Char.), 149.
nudicollis (Mes.), 40.
nupta (Om.), 87.
Nympha (Copt.), 173.

obducta (Chel.), 76.
obesa (Copt.), 193.
oblita (Mes.), 43.
obliterata (Cass.), 164.
oblonga (Cass.), 134.
—— (Isch.), 115.
obovata (Cass.), 120.
obscura (Cass.), 133.
obscuripennis (Cass.), 124.
obsoleta (C.), 62, 140.
obsoleta (Mes.), 60.
obtusa (Psalid.), 157.
obtusata (Cass.), 132.
ocellata (Cass.), 123.
ochracea (Hopl.), 6.
ochroleuca (Hopl.), 7.
octoguttata (Cass.), 141.
octomaculata (Pri.), 9.
octonotata (Pri.), 8.
octopunctata (Pri.), 9.
octopustulata (Pœcil.), 70.
—— (Pri.), 10.
ocularis (Char.), 151.
oculata (Copt.), 166.
odiosa (Hopl.), 6.
Œanthe (Cass.), 133.
Olivieri (Cass.), 117.
Omaspides, 91.
Omocera, 19.
Omoplata, 86.
Omoteina, 93.
opaca (Mes.), 51.
operosa (Copt.), 179.
opposita (Cass.), 141.
optata (Copt.), 198.
opulenta (Copt.), 168.
orbicularis (Asp.), 106.
orbifera (Char.), 149.
ordinata (Pœcil.), 66.
orientalis (Asp.), 107.
—— (Del.), 111.
ornata (Cass.), 56, 133.
—— (Copt.), 162.
—— (Pœcil.), 69.
orthogonia (Chel.), 75.
ovalis (Phys.), 98.
ovata (Phys.), 101.
Oxynodera, 26.

pacifica (Chel.), 79.
pæminosa (Cass.), 146.
pagana (Copt.), 181.
palacea (Copt.), 193.
Palleago (Asp.), 104.
pallens (Bas.), 97.
—— (Hopl.), 6.
pallescens (Copt.), 184.
—— (Hopl.), 3.

palliata (Dol.), 33.
pallida (Cass.), 132, 140, 204, 205, 206.
pallidicollis (Cass.), 138.
pallidicornis (Pri.), 11.
pallidipennis (Om.), 88.
pallido-guttata (Mes.), 57.
pallidula (Cass.), 141.
—— (Copt.), 197.
pallipes (Cass.), 13.
palmarum (Porph.), 16.
palmata (Copt.), 200.
pantherina (Chel.), 82.
—— (Pœcil.), 69.
papulosa (Copt.), 201.
paradoxa (Hopl.), 4.
parallela (Bat.), 95.
—— (Hopl.), 7.
Paramariboensis (Cass.), 54.
pardalina (Copt.), 175.
—— (Pœcil.), 69.
parellina (Cal.), 36.
partita (Chel.), 79.
parummaculata (Cass.), 121.
—— (Chel.), 76.
parvula (Cass.), 135.
Patagonica (Chel.), 80.
patruelis (Copt.), 198.
pauxilla (Cass.), 132.
pavonia (Pœcil.), 69.
pavonina (Cass.), 44.
pectoralis? (C.), 105.
pellucida (Copt.), 184.
peltata (Chel.), 79.
—— (Copt.), 200.
peltoides (Isch.), 115.
peregrina (Chel.), 76.
perforata (C.), 86.
—— (Dol.), 34.
—— (Desm.), 26.
perplexa (Del.), 193.
personata (Chel.), 80.
pertusa (Mes.), 64.
—— (Pol.), 22.
Peruana (Spil.), 19.
Peruviana (Desm.), 23.
—— (Dol.), 32.
Philippinensis (Lac.), 156.
phlyctæna (Cass.), 53.
Phœbe (Copt.), 187.
physodes (Copt.), 176.
PHYSONOTA, 97.
picea (Lac.), 154.
picta (Copt.), 158.
picticollis (Phys.), 98.
pictilis (Mes.), 64.
pictipennis (Pœcil.), 66.
picturata (Mes.), 60.
pilosella (Bas.), 97.
piperita (Cass.), 131.
placida (Copt.), 192.
plagiata (Copt.), 176.
—— (Mes.), 55.
plagicollis (Mes.), 62.
plagosa (Can.), 27.
plana (Phys.), 102.
PLATYAUCHENIA, 1.
PLATYCYCLA, 104.
platynota (Desm.), 24.
plebeja (Copt.), 203.
plecta (Copt.), 202.
pleurosticha (Mes.), 46.
plicata (Copt.), 200.
—— (Hopl.), 5.
plicatula (Char.), 152.
plumbea (Mes.), 50.
PŒCILASPIS, 65.
polita (Copt.), 160.
POLYCHALCA, 19, 22.
polysticha (Chel.), 80.
Pontifex (Aspid.), 110.
porcata (Pœcil.), 66.
porosa (Copt.), 195.
PORPHYRASPIS, 15.
porphyrio (Call.), 14.
porracea (Copt.), 188.
postica (Copt.), 158.
posticata (Copt.), 171.
præcincta (Copt.), 186.
præcox (Cass.), 118.
prætextata (Chel.), 79.
præusta (Copt.), 164.
prasina (Cass.), 125, 127.
primaria (Mes.), 49.
princeps (Chel.), 82.
—— (Dol.), 31.

Prioptera, 8.
procera (Can.), 26.
procincta (Copt.), 165.
profana (Copt.), 171.
prolixa (Phys.), 101.
promiscua (Copt.), 161.
propinqua (Copt.), 178.
proxima (Copt.), 169.
—— (Pœcil.), 72.
Psalidonota, 156.
puberula (Dol.), 33.
—— (Mes.), 54.
pubescens (Chel.), 73.
—— (Dol.), 30.
pudens (Cass.), 145.
pudibunda (Cass.), 133.
pudica (Copt.), 183.
puella (Copt.), 167.
—— (Om.), 90.
pugionata (Bat.), 94.
pugnax (Bat.), 94.
pulchella (Cass.), 134, 135.
—— (Hopl.), 6.
pulchra (Spil.), 18.
pullula (Copt.), 186.
pulvinata (Cass.), 137.
pumila (Cass.), 205.
pumilio (Cass.), 135.
punctaria (Copt.), 174.
punctata (Cal.), 18.
—— (Cass.), 56, 120, 175.
punctatissima (Copt.), 193.
punctatostriata (Char.), 153.
puncticollis (Cass.), 137.
—— (Chel.), 79.
puncticosta (Asp.), 104.
punctigera (Chel.), 79.
punctipennis (Chel.), 76.
—— (Copt.), 186.
—— (Taur.), 21.
punctulata (Phys.), 100.
Punctum (Aspid.), 110.
punicea (Copt.), 200.
—— (Mes.), 49.
pupillata (Cass.), 137.
—— (Copt.), 169.
purpurata (Copt.), 180.
purpurea (Copt.), 167.
pusilla (Cass.), 136.
pustulata (Copt.), 175.
pygmæa (Char.), 153.
pyramidata (Mes.), 39.

quadra (Copt.), 159.
quadrata (C.), 64.
—— (Copt.), 202.
quadraticollis (Mes.), 37.
quadricollis (Cyrt.), 37.
quadricornis (Hopl.), 4.
quadriguttata (Cass.), 141.
quadri-impressa (Pri.), 8.
quadri-lineata (Phys.), 100.
quadrilobata (Aspid.), 111.
quadrimaculata (Cass.), 54, 107.
—— (Lac.), 156.
quadrinodosa (Desm.), 25.
quadrinotata (Copt.), 158.
quadrioculata (Copt.), 174.
quadriplagiata (Cass.), 141.
—— (Mes.), 51.
quadripunctata (Cass.), 53.
quadripustulata (C.), 91.
quadriradiata (Aspid.), 113.
quadriramosa (Aspid.), 113.
quadriremis (Del.), 105.
quadrisignata (Copt.), 164.
—— (Pri.), 10.
quadrituberculata (Hopl.), 4.
quadrivittata (Copt.), 178.
quatuordecimmaculata (C.), 72.
quatuordecimmaculatum (Him.), 12.
quatuordecimnotata (Lac.), 155.
quatuordecimplagiata (Pœcil.), 70.
quatuordecimpunctata (Chel.), 78.
—— (Cass.), 204.
quatuordecimpustulata (Bot.), 70.
quatuordecimsignata (Mes.), 59.
—— (Pœcil.), 70.
quinquefasciata (Asp.), 105.
—— (Del.), 109.
—— (Pœcil.), 68.
quinquemaculata (Cass.), 142.
quinque-punctata (Chel.), 64.

radiata (Copt.), 189.
ramosa (Copt.), 189.
Redtenbacheri (Mes.), 48.
regalis (Cal.), 35.
Reicheana (Cal.), 2.
Reichei (Taur.), 21.
repanda (Copt.), 167.
resplendens (Call.), 15.
reticularis (Cass.), 202.
—— (Mes.), 56.
reticulata (Copt.), 202.
retifera (Copt.), 202.
retiformis (Cass.), 56.
retipennis (Mes.), 40.
retusa (C.), 29.
—— (Desm.), 25.
rimosa (Cass.), 143.
rorida (Cass.), 133.
rosea (Cass.), 128.
roseocincta (Copt.), 159.
roseomarginata (Cass.), 146.
roseopicta (Copt.), 172.
roseosignata (Copt.), 177.
rotalis (Copt.), 195.
rotundata (Char.), 153.
rubescens (Mes.), 60.
rubicunda (Copt.), 189.
rubicundula (Mes.), 47.
rubiginosa (Can.), 27.
—— (Cass.), 125.
rubra (Call.), 15.
rubricatum (Him.), 14.
rubricollis (Pœcil.), 67.
rubro-cincta (Char.), 150.
—— (Copt.), 201.
rubro-delineata (Pœcil.), 66.
rubro-discoidalis (Copt.), 166.
rubro-dorsata (Aspid.), 114.
rubro-fasciata (C.), 33.
rubro-guttata (Pœcil.), 70.
rubro-lineata (Copt.), 195.
rubro-marginata (Hopl.), 5.
rubro-notata (Phys.), 100.
rubro-ornata (Copt.), 192.
rubro-picta (Copt.), 177.
rubro-plagiata (Mes.), 60.
rubro-reticulata (Mes.), 57.
rubro-testacea (Copt.), 193.
rubro-vittata (Copt.), 178.
rubrum (Him.), 14.
ruderaria (Desm.), 26.
rufa (Copt.), 124.
rufescens (C.), 63.
ruficeps (Chel.), 83.
rufina (Call.), 14.
rufipennis (Chel.), 83.
rufiventre (Him.) 13.
rufomarginatum (Him.), 13.
rufo-reticulata (Pœcil.), 66.
rufo-signata (Mes.), 60.
rufovirens (Cass.), 128.
rufula (Call.), 15.
ruginosa (Lac.), 154.
rugipennis (Copt.), 191.
rugosa (Cass.), 142.
—— (Oxyn.), 27.
—— (Porph.), 16.
rugulosa (Cass.), 129.
Russica (Cass.), 138.
russula (Chel.), 76.
rustica (Copt.), 194.
rutila (Copt.), 189.

Sahlbergi (Copt.), 188.
salebrosa (Desm.), 23.
Salicorniæ (Cass.), 134.
sanguinea (Porph.), 15.
sanguineum (Im.), 15.
sanguinolenta (Cass.), 122.
—— (Cass.), 125, 128.
—— (Copt.), 192.
sanguinosa (Cass.), 125.
Saundersi (Mes.), 56.
scabra (Copt.), 201.
scabriuscula (Char.), 151.
scabrosa (Taur.), 22.
scalaris (Copt.), 161.
scapularis (Om.), 89.
scapulosa (Om.), 89.
scenica (Copt.), 188.
Schaumi (Mes.), 59.
Schœnherri (Aspid.), 109.
Schüppelii (Chel.), 82.
scitula (Chel.), 85.
Scolymi (Cass.), 127.
scoparia (Mes.), 54.

scoriosa (Can.), 27.
scripta (Cass.), 122.
scrobiculata (Mes.), 39.
sculpturata (Lac.), 156.
scutellaris (Cass.), 144.
sedecimmaculata (Cass.), 119.
—— (Pœcil.), 69.
sedecimpunctata (C.), 70.
sedecimpustulata (Pœcil.), 70.
sedecimstillata (Chel.), 78.
sejuncta (Copt.), 184.
seladonia (Cass.), 125.
selecta (Ep.), 2.
Selenis, 85.
sellata (Copt.), 167.
—— (Spil.), 18.
semiannulata (Copt.), 168.
semicirculare (Him.), 12.
semicircularis (Cass.), 12.
semifasciata (Copt.), 172.
semifasciatum (Him.), 12.
semiglobosa (Cass.), 137.
semilineata (Om.), 89.
seminulum (Char.), 150.
semiviridis (Hopl.), 3.
semivittata (Copt.), 168.
separata (Copt.), 173.
—— (Phys.), 99.
septemdecimpunctata (Copt.), 160.
septemnotata (Copt.), 162.
sepulchralis (Copt.), 200.
seraphina (Cass.), 147.
serena (Copt.), 168.
seriata (Mes.), 61.
sericata (Dol.), 34.
Serinus (Mes.), 52.
serricosta (Hopl.), 7.
sertata (Copt.), 202.
Servula (Copt.), 195.
setifera (Cass.), 135.
severa (Copt.), 199.
sexguttata (Copt.), 173.
sexlunata (Mes.), 61.
sexmaculata (Copt.), 160.
—— (Mes.), 39.
—— (Pri.), 8.
sexmaculosa (Mes.), 38.
sexnotata (Cass.), 161.
—— (Copt.), 160.
sexplagiata (Mes.), 51.
sexpunctata (Cass.), 163.
—— (Copt.), 162.
—— (Copt.), 185.
sexpustulata (Mes.), 44.
sexsignata (Mes.), 43.
sexstillata (Mes.), 42.
Sibirica (Cass.), 147.
signata (Cass.), 138.
signaticollis (Cal.), 36.
—— (Chel.), 84.
—— (Copt.), 198.
signatifera (Copt.), 196.
signatipennis (Cass.), 120.
signifer (Cass.), 181.
Signoreti (Copt.), 170.
silacea (Aspid.), 109.
Silbermanni (Mes.), 45.
similata (Copt.), 167.
—— (Mes.), 44.
similis (Cass.), 126.
simplex (Copt.), 194.
smaragdina (Cass.), 22.
—— (Mes.), 51.
—— (Taur.), 20.
Smithii (Chel.), 77.
sobrina (Om.), 90.
socia (Aspid.), 114.
—— (Chel.), 74.
sodalis (Mes.), 64.
Solani (Cass.), 118.
Solieri (Om.), 88.
Sommeri (Mes.), 42.
sordida (Copt.), 182.
Soror (Char.), 149.
spadicea (Cass.), 128.
sparsa (Sel.), 86.
speciosa (Cass.), 122.
spectabilis (Asp.), 104.
spectrum (Lac.), 154.
specularis (Om.), 92.
speculum (Copt.), 165.
Spencei (Phys.), 99.
Spergulæ (Cass.), 135.
Sphæropalpus, 1.
sphærula (Cass.), 136.

Spilophora, 18.
spilota (Cass.), 120.
spinifex (Sel.), 85.
spinipes (Phys.), 102.
spinosa (Bat.), 95.
splendidula (Cass.), 131, 135.
spreta (Char.), 150.
spurca (Dol.), 29.
squalida (Om.), 89.
St. Crucis (Aspid.), 111.
Stevensi (Mes.), 41.
stictica (Aspid.), 112.
stigma (Copt.), 159.
stigmata (Cyrt.), 42.
stigmatica (Cass.), 128.
—— (Copt.), 199.
stigmatilis (Phys.), 99.
stillata (Copt.), 174.
stolata (Aspid.), 109.
stolida (Bas.), 97.
straminea (Del.), 193.
striato-punctata (Copt.), 184.
strigata (Dol.), 33.
strigula (Cass.), 206.
striolata (Cass.), 138.
stulta (Copt.), 193.
stupidula (Copt.), 179.
Sturmii (Chel.), 74.
subcincta (Del.), 111.
subcornuta (Desm.), 25.
subcruciata (Aspid.), 112.
subfasciata (Cyrt.), 63.
subferruginea (Cass.), 140.
subirrorata (Copt.), 175.
sublævis (Mes.), 40.
sublineata (C.), 62.
sublucida (Om.), 88.
submaculata (Copt.), 164.
subnervosa (Pœcil.), 66.
subnotata (Copt.), 160.
—— (Mes.), 44.
subplagiata (Copt.), 163.
subpunctata (Chel.), 75.
subpustulata (Cyrt.), 42.
subreticulata (Cass.), 136.
—— (Mes.), 44.
subrugosa (Copt.), 183.
—— (Mes.), 61.
subtincta (Copt.), 194.
subvittata (Copt.), 178.
succincta (Copt.), 172.
succinea (Copt.), 185.
Suffriani (Cass.), 141.
sugillata (Aspid.), 109.
sulcata (Cass.), 116.
sulcipennis (Phys.), 101.
sulphurago (Cass.), 133.
sulphurea (Cass.), 129.
supposita (Cass.), 204.
surda (Copt.), 166.
suspecta (Copt.), 193.
suspiciosa (Mes.), 62.
suturalis (C.), 87.
—— (Mes.), 62.
suturella (Pœcil.), 71.
Swartzii (Chel.), 75.
syrtica (Cass.), 129.

tabida (Copt.), 202.
tædiosa (Cass.), 143.
tæniata (Hopl.), 7.
tantilla (Char.), 152.
tarsata (Dol.), 33.
Tauroma, 19.
Taurus (Taur.), 21.
tecta (Aspid.), 109.
tenebrosa (Bat.), 96.
—— (Disc.), 27.
tenella (Copt.), 185.
tenera (Cass.), 145.
tenuicula (Cass.), 130.
ternata (Copt.), 162.
tersa (Om.), 87.
tessellata (C.), 70.
—— (Char.), 151.
testacea (Cass.), 205.
testaceomarginata (Chel.), 79.
testudinea (Cass.), 142.
testudinaria (Cass.), 206; (Copt.), 189.
—— (Phys.), 100.
Testudo (Cass.), 146.
tetrastigma (Cass.), 53.
tetrica (Cass.), 117.
texta (Pœcil.), 66.
textilis (Mes.), 45.

thalassina (Copt.), 153.
—— (Mes.), 58.
thoracica (Cass.), 125.
—— (Cass.), 126.
thoracicum (Him.), 13.
THYREOMORPHA, 3.
tigrina (C.), 29.
—— (Cass.), 118, 139.
—— (Mes.), 61.
timefacta (Cass.), 131.
tomentosa (Mes.), 54.
tosta (Cass.), 146.
trabeata (Copt.), 182.
translucida (Phys.), 102.
transversa (C.), 91.
transversalis (Cyrt.), 63.
—— (Om.), 87.
tredecimmaculata (Chel.), 78.
tredecimnotata (Copt.), 161.
tredecimpunctata (Lac.), 155.
tredecimsignata (Copt.), 161.
trepida (Copt.), 203.
trichroa (Om.), 88.
tricolor (Cass.), 205.
—— (Char.), 149.
tricolorata (Om.), 88.
trifasciata (Om.), 92.
trigemina (Spil.), 18.
trilineata (Cass.), 123.
trimaculata (Copt.), 159.
—— (Spil.), 18.
trimaculatum (Im.), 18.
trisignata (Copt.), 162.
tristigma (Mes.), 49.
tristis (Disc.), 28.
—— (Porph.), 16.
tristriata (Cass.), 137.
trivialis (Om.), 90.
trivittata (Copt.), 177.
Troberti (Del.), 108.
Troglodytes (Cass.), 129.
truncata (Bat.), 95.
—— (Taur.), 21.
tuberculata (C.), 73.
—— (Copt.), 188.
—— (Del.), 114.
Tucumana (Chel.), 81.
tumidula (Mes.), 38.
tumulus (Mes.), 45.
Tunisiensis (Cass.), 134.
turgida (Phys.), 99.
turpis (Desm.), 24.
turrifera (Bat.), 95.
turrita (Mes.), 38.

umbrata (Cass.), 134.
undecimmaculata (Chel.), 77.
undecimnotata (Cass.), 147.
——(Copt.), 160.
undecimpunctata (Copt.), 159.
unicolor (Cass.), 133.
uniformis (Copt.), 183.
unimaculata (Cass.), 141.
urbana (Chel.), 75.
urticæ (Cass.), 135.
usta (Copt.), 180.

valida (Copt.), 178.
vana (Copt.), 176.
variabilis (Cass.), 123.
—— (Chel.), 81.
varians (Aspid.), 158.
—— (Chel.), 81.
—— (Copt.), 178.
—— (Mes.), 63.
varicolor (Copt.), 158.
varicornis (Copt.), 185.
varicosa (Can.), 26.
variegata (Dol.), 29.
variolaris (Mes.), 39.
variolosa (C.), 74.
—— (Desm.), 22.
venosa (Cal.), 35.
—— (Sel.), 86.
ventralis (Copt.), 159.
ventricosa (Copt.), 190.
venusta (Chel.), 83.
verecunda (Mes.), 51.
vermiculata (Chel.), 83.
verrucata (Copt.), 176.
verrucosa (Mes.), 46.
versicolor (Copt.), 192.
vesicularis (Copt.), 175.
vestita (Cyrt.), 38.
vetula (Mes.), 40.
vetusta (Deloy.), 191.

vibex (Cass.), 126.
vicina (Deloy.), 181.
—— (Mes.), 58.
vidua (Mes.), 41.
vigens (Copt.), 194.
vigintimaculata (Cass.), 121.
vigintisexnotata (Lac.), 155.
vilis (Copt.), 186.
villica (Copt.), 171.
vinula (Copt.), 165.
virescens (Copt.), 197.
virgata (Copt.), 172.
virginalis (Copt.), 176.
Virgo (Copt.), 183.
virgulata (Copt.), 182.
viridana (Cass.), 128.
viridicærulea (Taur.), 19.
virididorsata (Asp.), 106.
viridimaculata (Ep.), 2.
viridinotata (Copt.), 175.
—— (Hopl.), 4.
viridipennis (Cass.), 129.
viridis (Cass.), 125, 127, 144.
viridisignata (Bat.), 97.
viriditincta (Hopl.), 4.
viridula (Cass.), 135.
vitellina (Copt.), 194.
vitrea (Eur.), 104.
vitreata (Copt.), 191.
vittata (Cass.), 123.
vitticollis (Phys.), 98.
vittifera (Pœcil.), 72.
vittigera (Copt.), 194.
vittipennis (Copt.), 177.
vittula (Cass.), 138.
vivida (Copt.), 203.
voluptaria (Copt.), 173.
vulnerata (Mes.), 42.

Wahlbergi (Asp.), 105.
Waterhousei (Copt.), 172.
Wesmaeli (Copt.), 169.
Westermanni (Asp.), 107.
—— (Pri.), 8.
Westwoodii (Asp.), 105.
Whitei (Pri.), 11.
Wollastoni (Chel.), 73.

xanthocera (Porph.), 17.
xanthoptera (Pœcil.), 68.
xerampelina (Phys.), 103.

Zebra (Bot.), 68.
Zelleri (Copt.), 174.
Zetterstedti (Copt.), 180.
Zona (Copt.), 165.
zonata (Copt.), 195.

1756

SMITHSONIAN INSTITUTION LIBRARIES
3 9088 00239236 3
nhent QL577.B82X
pt. 9 Catalogue of coleopterous insects